

**PATTERN OF DEGENERATIVE LUMBAR
RETROLISTHESIS IN BASRAH****Thamer A Hamdan*, Mubder A Mohammed Saeed# & Yas khudair
Hadood@**

*FRCS, FRCP, FICS, FACS, American Board (Orth.), Professor of Orthopedic Surgery. #FICMS, (Orth.) Assistant Prof. of Orthopedic Surgery. College of Medicine, University of Basrah, Basrah, Iraq. @MB,ChB, Postgraduate Arab Board Medical Specialization (Orth.)

Abstract

Although retrolisthesis is not a well-known condition by many medical specialists dealing with back problems and it has been regarded as a radiological incidental finding with no clinical significance, a growing prove is now evolving stating that retrolisthesis could be a cause of many backache complaints and a sequelae of an altered spine biomechanics.

Objectives: to study and analyze the various biomechanical characteristics of retrolisthesis and its relationship with various radiological parameters of the lumbar spine and other patient's factors. Patients and Method: Forty patients, twenty six males, and fourteen females with an age range from 40–66 years with radiological evidence of significant lumbar spine retrolisthesis (slip > 3 millimeters) were evaluated clinically and radiologically by plain radiography and MRI in Basra General Hospital and Ibn AL-Bittar Private Hospital, during the period from the 1st of August 2014 to the 1st of March 2015. After a thorough history and physical examination, various radiological parameters were obtained including the lumbar lordosis, sacral slop, pelvic incidence, pelvic tilt, all those measurements were done digitally. A statistical analysis was made via IBM SPSS ver.17 and the results were compared with that of similar studies. Results: The retrolisthesis was found to be more common with more slip distance in males than in females, males=26 (65%), females=14 (35%), P-value=0.026, the mean of slip in mm in males was (4.002) and in females was (3.71). The patients had a mean BMI equal to (26.025=overweight). The most common level at which retrolisthesis occur in this study was the L5-S1 (40%) then L4-L5 (22.5%).

All the radiological parameters (the lumbar lordosis, sacral slop, pelvic incidence, and pelvic tilt) were lower than the known normal values in healthy subjects. Conclusion: It seems that retrolisthesis is not just an incidental finding, it may be a kind of a compensatory reaction for an abnormal spine biomechanics. The males are affected more frequently than females with a more slip distance.

The L5-S1 followed by the L4-L5 are the most common sites in both sexes. The degenerative spinal disease is the main cause of retrolisthesis in all patient's groups regardless of sex or age.

Introduction

Listhesis was defined as displacement (backward or forward) of one vertebra relative to the one below 3 mm or greater. Less than 3 mm displacement is considered to be within the normal range. The selection of 3 mm as the cut point for the definition of disease was based on the fact that this is the criterion commonly

used in orthopedic clinical practice¹ This 3-mm cutoff corresponded to a slip of 8% that was used as the lower limit to define retrolisthesis.

Retrolisthesis has historically been regarded as an incidental finding, one that does not cause any symptoms and is considered to be of little or no clinical significance. The literature has found a possible association between retrolisthesis

and increased back pain and impaired back function²⁻⁵.

Retrolisthesis is found mainly in the cervical spine and lumbar region but can also be often seen in the thoracic spine⁶.

Why retrolisthesis occur? Is it merely a result of a cause an effect scenario? Or it is a compensatory response for an altered spine biomechanics and an attempt from the spine to stay as close as possible to its previous normal alignment?

In this study, we will try to explain the various biomechanical characteristics associated with the retrolisthesis by analyzing some of the important biomechanical parameters in the affected spines, so that, we may figure out the real cause(s) of the retrolisthesis and whether it is by the chance phenomenon in its occurrence or an important compensatory strategy of the spine in response to some of its affections.

In a closed, related mechanical system, there is an important mechanical concept which states that any alteration in a part is followed by a suitable and related change in another part in the form of movement or change of its position, velocity or direction in any form.

This also can be applied to the spine as a whole, and the spinal motion segment in particular, emerging a big question whether retrolisthesis is the end result of a far or near spinal biomechanical change which follow the above mentioned mechanical fact.

Patients and Methods

This is a descriptive study done to disclose some of the relevant characteristics of retrolisthesis in selected patients' sample.

The sample in this study comprised of 40 patients; 26males 65% and 14 females 35%, age: 40-66 years, with back pain and a proved plain radiological diagnosis of retrolisthesis in the lumbar spine, they were collected from the attendance of Basra General Hospital and Ibn AL-Bittar Private Hospital in Basra, starting from

the 1st of August 2014 till the 1st of March 2015.

All the patients were evaluated clinically and radiographically after applying the inclusion and exclusion criteria as below:

The Inclusion Criteria:

The age is more than 18 years old.

Radiological proof of lumbar spine retrolisthesis (defined as a backward slip > 8% or 3 millimeter)

The Exclusion criteria were as follow:

Any case following major back trauma, old or recent.

Spinal or paraspinal Infections or Tumors, old or recent.

Postoperative cases, old or recent.

Obvious congenital and/or Developmental anomalies, including scoliosis.

All patients were subjected to full clinical evaluation including history, physical examination and plain radiographic examinations to the Lumbar spine and pelvis (both AP and Lateral views in standing position, fixed not dynamic films) with MRI scanning of the lumbosacral spine. All the examined parameters were documented in the patient's questionnaires.

After that the parameters were collected and analyzed statistically using the (IBM SPSS version 17.0) software, and a significant P value was set at ≤ 0.05 .

Obtaining the measurements of the various radiological parameters were done on the digital photos of the patient's plain radiographies of a size (17*14 inches) in lateral views to the lumbosacral spine while the patient was standing with a film-tube distance equal to (72) inches, then the calculation was made after correction of the magnification effect associated with the plain radiographic technique by a special computer software; (Image J) ®, as followed:


The pelvic incidence (PI)⁷ is defined as the angle subtended by a line connecting the axis of the femoral heads to the midpoint of the endplate of S1 and a perpendicular line to the endplate of S1 at its midpoint. Figure (1). The sacral slope

(SS)⁷ is defined as the angle between the superior endplate of S1 and the horizontal plane. Figure (1)

The pelvic tilt (PT)⁷ is defined as the angle between a line joining the midpoint

of the superior endplate of S1 and the axis of the femoral heads with the vertical plane. Figure (1)

Figure 1: The pelvic incidence (PI), the sacral slope (SS) and the pelvic tilt (PT), showing the way of making the measurements.



The retrolisthesis Slip

Measurement of retrolisthesis by Iguchi method⁸. A line is drawn along the inferior end plate of the vertebra that suspected to be slipped backward. Two lines are erected perpendicular to this line to pass through the adjacent posterior corners of the vertebral bodies. The distance between the points at which these 2 lines intersect the end plate line is the amount of retrolisthesis as demonstrated in figure 2.

The Lumbar lordosis⁹

This is measured on the erect lateral radiograph of the lumbo-sacral spine, as

the angle between the superior end-plates of L1 and S1, using the Cobb's method. (Figure 3)

The other parameters; namely the patient's age, gender, smoking habit, body mass index, the presence of backache, radiculopathy with the duration of his/her complaint and MRI findings,. All were

grouped for each patient in an Excel® file using the Microsoft Office® 2013 and statistically analyzed using the (IBM SPSS version 17.0) software.

Figure 3: The method of measuring the angle of lumbar lordosis using the Cobb's method.



Results

The different demographic parameters for the patients were summarized in table I.

Table I: The demographic parameters

Gender	Males		Females		P-value
	26 (65%)		14 (35%)		0.026
	Minimum	Maximum	Mean	SD	
Age	40	66	50	±6.63	
BMI	19.60	36.85	26.02	±4.05	
Duration of pain in months	4	72	18.4	±13.99	
	Smoking Habit				
Never	30				
Former	8				
Current	2				

The presenting symptoms for the patients were ranged from backache, and radiculopathy, in isolation or a combined presentation as shown below in table II.

Table II: The frequency of the presenting complaints

The Presenting Complaint	Backache	Radiculopathy	
		Rt	Lt
Number of Cases	40 (100%)	19 (47.5%)	12 (30%)

Radiological findings of the retrolisthetic levels in form of narrowing of the intervertebral disc

space, the osteophytes and the end plate sclerosis were all summarized in table III.

Table III: Radiological findings of retrolisthetic levels.

Radiological findings of retrolsthetic levels	No. of retrolsthetic level.
Narrowing of intervertebral space	36(83%)
Osteophytes	30(69%)
End plate sclerosis	18(41%)

The MRI findings of the retrolsthetic levels in form of disc T2 signal loss (signifying disc dehydration), the posterior degenerative changes (the

facet joint arthropathy and the ligamentum flavum hypertrophy) and modic changes (I, II, III) were summarized in table V.

Table IV: MRI findings of retrolsthetic levels

MRI findings of retrolsthetic levels	No. of retrolsthetic level.
Disc T2 signal loss	33(76%)
Posterior degenerative changes	13(30%)
Modic changes	11(25%)

The patient’s radiographic parameters that was used in this study were measured by the computer and the results as shown below in table V.

Table V: The radiographic parameters

	Minimum	Maximum	Mean	SD
Retrolstthesis slip in mm	3	5	4.25	±0.74
Pelvic incidence	250	680	36.350	±8.65
Sacral slope	110	530	25.950	±8.97
Pelvic tilt	50	150	10.270	±3.04
Lumbar Lordosis	250	630	36.220	±8.21

The measurements of the specific characteristics of each retrolsthetic level and its main features in the form

of the level it affect and the magnitude of the slippage in millimeters were as shown below in table VI.

Table VI: The characteristics of retrolstthesis according to the lumbar spine level

	Frequency of retrolisthesis	Minimum slip mm	Maximum slip mm	Mean of Slip	SD	P Value
L1 – L2	2 (5 %)	3	3	3	0	0.93
L2 – L3	8 (20 %)	4	5	4.16	0.675	0.04
L3 – L4	8 (20 %)	3	5	4.24	0.546	0.16
L4 – L5	9 (22.5 %)	4	5	4.25	0.485	0.23
L5 – S1	16 (40 %)	3	5	4.56	0.765	0.04

The sex of patients and its relation to the most common level it affects were

summarized as shown below in table VII.

Table VII: The Frequency of retrolisthesis in each level in the patients.

		In all Patients		Males		P-Value	Females		P-Value
L1-L2	Frequency of Retrolisthesis	2	5%	1	2.5%	0.01	1	2.5%	0.01
L2-L3	Frequency of Retrolisthesis	8	20%	5	12.5%	0.056	3	7.5%	0.052
L3-L4	Frequency of Retrolisthesis	8	20%	5	12.5%	0.056	3	7.5%	0.052
L4-L5	Frequency of Retrolisthesis	9	22.5%	7	17.5%	0.06	2	5%	0.064
L5-S1	Frequency of Retrolisthesis	16	40%	11	27.5%	0.03	5	12.5%	0.043
Total		43*	107.5%	29	75%		14	32.5%	

*There are 43 level with retrolisthesis in 40 patients, because 3 patients have multilevel retrolisthesis.

The sex of patients and its relation to the mean of the posterior slippage were

summarized as shown below in table VIII.

Table VIII: The mean of the posterior slippage in each level in the patients.

		In all Patients	Males	P-Value	Females	P-Value
L1-L2	Mean of slip mm	3	3	0.00	3	0.00
L2-L3	Mean of slip mm	4.16	4.04	0.46	3.9	0.43
L3-L4	Mean	4.24	4.14	0.35	3.7	0.34

L4	of slip mm					
L4-L5	Mean of slip mm	4.25	4.27	0.45	4.02	0.54
L5-S1	Mean of slip mm	4.56	4.58	0.03	3.93	0.04

The radiological parameters of patients and its relation to the sex as shown below in table IX.

Table IX: The characteristics of radiological parameters

	In all Patients	Males	P-Value	Females	P Value
Mean of Pelvic incidence	36.35	37.65	0.76	38.65	0.75
Mean of Sacral slope	25.95	28.95	0.65	31.68	0.68
Mean of Pelvic tilt	10.27	13.95	0.75	15.56	0.76
Mean of lumbar lordosis	36.22	10.45	0.03	9.37	0.76

The different radiographic parameters of the patients were compared with normal values as shown in table X.

Table X: The comparison of the different radiographic parameters in normal values and in cases of retrolisthesis¹⁰.

The Parameter	The Normal Mean*	The readings in this study
Lumbar lordosis(°)	66.36	36.22
Sacral slope(°)	41.18	25.95
Pelvic tilt(°)	11.96	10.27
Incidence(°)	53.13	36.35

The types of treatment of the patients were summarized as below in table XI.

Table XI: Treatment of retrolisthesis.

No. of patients	Type of Treatment		
34	Conservative Treatment		
6	operative Treatment	4	Laminectomy
		2	laminectomy +discectomy
		0	Instrumentation

Discussion

In this study, we shed a light on the increasing awareness among spine surgeons that retrolisthesis could be more than an incidental radiological finding and it could play a real role in the pathophysiology of back pain in the affected patients.

Why and how retrolisthesis cause symptoms? A study done by O'Brian in 1983 showed that retrolisthesis can cause narrowing of the disc space when the annulus fibrosus bulges posteriorly¹¹. Concurrently, there can be a relative translation of the superior articular process of the vertebra caudal to the mobile segment in the direction of the intervertebral foramen. This can cause a lateral stenosis that can produce painful radicular symptoms¹².

Studies on both white and African American women showed that retrolisthesis was associated with a higher likelihood of low back pain^{4,5}. Although once believed to be a benign finding, it is becoming more apparent that retrolisthesis can be a source of morbidity for patients.

In this study; being merely a descriptive type, we can't formulate a trusted hypothesis about the association between

retrolisthesis and back pain, this needs more big experimental study or randomized controlled clinical trials.

All patients share the same type of retrolisthesis, the partial type, this may reflect that other types of retrolisthesis are rare occurrence among patients of similar parameters. All patients were also having

degenerative spinal condition, of varying degrees.

We agree with other studies in regard the gender; retrolisthesis was seen in male^{13,14}.

Other study, as in that of Jeon C-H (2013)¹⁵ no statistically significant gender variation was noted in the patients suffering from back pain with a pure and significant retrolisthesis.

The prevalence of retrolisthesis did not vary by sex, age, race, smoking status, or education level when compared with individuals with normal sagittal of spine^{13,4}.

The age of the patients were ranged from (40) years to (66) years, although the inclusion criteria in this study involved any adult age group starting from age of 18 years, no age was shown to be less than 40 years; which may be regarded as the start line of degenerative process as the facet joint degeneration is relatively uncommon in persons under 40 years¹⁶, if we add to this fact that all the patients were having lumbar spine degeneration of varying degrees, we can figure out a significant relationship between the retrolisthesis and degeneration.

The average BMI of the all patients was (26.025, SD±4), this means that the predominate body habitus in retrolisthesis patients is the normal to overweight style. Our finding goes with that of Shen et al (2007)¹³, they found that retrolisthesis patients had a BMI with a mean equal to (28) with (SD±6.1).

We could explain this as the retrolisthesis is not only require some type or form of posterior weakening in order to occur, it

also need a more than normal or average posteriorly directed shearing force applied on the relevant vertebra, this is manifested by the net result of the overweight body mass. Our conclusion is that retrolisthesis is a disease of the over weighted persons.

The pelvic incidence (PI), sacral slop (SS), the pelvic tilt (PT), and the lumbar lordosis in our patients were all statistically lower than that of normal subjects and patients with anterolisthesis in other studies which were taken as a comparison with our results^{10,17}. When the lumbar spine is hyperlordosis, the contact force on the posterior joints and the intervertebral tilt will increase, thereby increasing the forward sliding force. By contrast, the contact force on the anterior intervertebral disc will increase with hypo lordosis, subsequently decreasing the intervertebral tilt.

As hypolordosis is related to a lower SS, and subsequently lower PI^{18,19}, backward displacement could occur in patients with a low PI for this reason.

In contrast, Degenerative anterolisthesis was reported to have a higher PI, SS, and lumbar lordosis than that in retrolisthesis^{20,21}.

The main level with more frequent retrolisthesis was the L5-S1, (16 patients, 40%) probably due to the high stress which is applied for this level (junctional level).

This was different than other studies were the retrolisthesis occur more commonly in higher lumbar spine levels, namely in L3-L4 (44.3%) followed by L2-L3 (35.7%) as stated in Jeon C-H (2013)¹⁵.

Treatment and what is its most accepted option, was also not a totally agreed point among all researchers, the choice depends on multiple factors in deciding which way is the best and most suitable for both the patient and his/her treating surgeon.

Before all, the conservative style was still, and will probably stay to be, the standard for retrolisthesis treatment, since there are no proofs that posterior vertebral slipping may in itself be a cause of low back pain¹⁰, also the surgical treatment may result in a more posterior slippage as a result of lowering the disc height after discectomy²².

Only 6 patients from the 40 patients were treated surgically in form of laminectomy (4 patients) and less commonly a laminectomy with discectomy (2 patients). No patient in this study was in need for fusion, this was due to the fact that a bilateral laminectomy and discectomy had not been carried out for any patient.

Conclusion

Retrolisthesis is commoner than it was assumed.

Partial retrolisthesis is commonest type The L5-S1 is the commonest level for retrolisthesis followed by L4-L5 and then by other higher lumbar levels.

The degenerative spinal disease is the main cause of retrolisthesis in this study.

Retrolisthesis is more common in males; with more slip, than in females.

Retrolisthesis is associated with less lumbar lordosis than in the normal populations.

References

1. Kevin K. Kang, Michael S. Shen, Wenyan Zhao, et al, Retrolisthesis and lumbar disc herniation: a postoperative assessment of patient function. *Spine J.* 2013; 13: 367–372
2. Deyo RA, Bass JE. Lifestyle and low-back pain. The influence of smoking and obesity. *Spine J* 1989; 14:501–506
3. Videman T, Battie M. A critical review of the epidemiology of idiopathic low back pain. *Eur Spine J;* 1996:637–641
4. Vogt MT, Rubin D, Valentin RS, et al. Lumbar olisthesis and lower back symptoms in elderly white women. *The Study of Osteoporotic Fractures.* *Spine J* 1998; 23:2640–2647
5. Vogt MT, Rubin DA, Palermo L, et al. Lumbar spine listhesis in older African American women. *Spine J* 2003; 3:255–261
6. (6)Peter Robb, Retrolisthesis, Health - Naturally Benalla, Bobinawarra & Mornington Victoria, Australia, updated Nov 2013, website: <http://www.headbacktohealth.com/Retrolisthesis.html>, last seen in 12:41 pm 30/1/2015

7. Legaye J, Duval-Beaupere G, Hecquet J, et al. Pelvic incidence: a fundamental pelvic parameter for three-dimensional regulation of spinal sagittal curves. *Eur Spine J* .1998; 7:99–103
8. Iguchi, Wakami T, Kurihara A, et al. Lumbar multilevel degenerative spondylolisthesis: radiological evaluation and factors related to anterolisthesis and retrolisthesis. *J Spinal Disord Tech* .2002; 15:93–99
9. Berlemann U, Jeszenszky D, Buhler D, et al. Mechanisms of retrolisthesis in the lower lumbar spine. A radiographic study. *ActOrthop Belg*. 1999; 4:427–77
10. Boulay C, Tardieu C, Hecquet J, et al. Sagittal alignment of spine and pelvis regulated by pelvic incidence: standard values and prediction of lordosis. *Eur Spine J*. 2006; 15:415-422
11. O'Brian JP. The role of fusion for chronic low back pain. *Orthop Clin North Am*. 1983; 14:1893–1896
12. Sihvonen T, Lindgren KA, Airaksinen O, et al. Movement disturbances of the lumbar spine and abnormal back muscle electromyographic findings in recurrent low back pain. *Spine*.1997; 22:289–295
13. Shen M, Razi A, Lurie JD, et al. Retrolisthesis and lumbar disc herniation: a preoperative assessment of patient function. *Spine J*. 2007; 7:406–413
14. Lai-Chang He, Wang YX, Gong JS, et al. Prevalence and risk factors of lumbar spondylolisthesis in elderly Chinese men and women. *Eur Radiol*. 2014; 24:441–448
15. Jeon C-H. Degenerative retrolisthesis: is it a compensatory mechanism for sagittal imbalance?? *.Bone Joint J* 2013; 95: 1244–1249
16. Gries NCBU, Moore RJ, Vernon-Roberts B., et al . Early histologic changes in lower lumbar discs and facet joints and their correlation. *Euro Spine J* 2000; 9:23–29.
17. Roussouly P, Pinheiro-Franco JL. Biomechanical analysis of the spino-pelvic organization and adaptation in pathology. *Eur Spine J* 2011; 20:609–618
18. Schwab F, Lafage V, Patel A, et al .Sagittal plane considerations and the pelvis in the adult patient. *Spine*.2009; 34:1828–1833
19. Mac-Thiong JM, Labelle H, Berthounaud E, et al. Sagittal spinopelvic balance in normal children and adolescents. *Eur Spine J* 2007; 16:227–234
20. Schuller S, Charles YP, Steib JP. Sagittal spinopelvic alignment and body mass index in patients with degenerative spondylolisthesis. *Eur Spine J* 2011; 20:713–719
21. Barrey C, Jund J, Perrin G, et al .Spinopelvic alignment of patients with degenerative spondylolisthesis. *Neurosurgery*2007; 61:981–986
22. F. Postacchini, C. Cinotti, O. L. Osti. Spinal fusion and disc prosthesis at primary surgery. 20. In: Franco Postacchini. *Lumbar Disc Herniation* .4th Edition. Springer Vienna. 1999:527-528