

***Theoretical study for some novelty Triazole derivatives by using
Ab Intio calculations(RHF -Model) .***

دراسة نظرية لبعض مشتقات التريازول باستخدام الطريقة التامة (RHF) .

**Manal. A. Mohammad Gehan H. Abdul- Ameer
Lemya S. Mahdi
Dep. Of chemistry, college of Education for pure science,
college of pharmacy ,
university of Karbala, Iraq.**


Abstract:-

RHF calculations, within Gaussian03 Program, have been carried out after complete optimization of geometry on X and Y disubstituted of (1H1,2,3,-triazole-4-Yl) ethoxy) methy)-1H-1,2,3-triazole), where X is H, COOH, COCH₃, and 2,3,4,6-Tetra-O-acetyl-β-D-glucopyranosyl. Y is C₇H₁₅. It was found that the compound R-sugar the more Stability than other compounds and all of them increasing dipole moment and hardness and decreasing in HOMO and LUMO energies. Geometrical parameters ,total energy ,electron charge ,Ionization energies , and the E⁰,H⁰,G⁰,A⁰,C_p,C_v and S⁰ thermodynamics functions are reported.

Key words: RHF study , Triazole ,thermodynamics functions.

الخلاصة:-

تضمنت هذه الدراسة اعتماد برنامج كاوس (Gaussian 03) لاستعمال طريقة الحساب التام على وفق طريقة نظرية دالة الكثافة (RHF) , وذلك لغرض حساب الأبعاد الهندسية (أطوال و زوايا التآصر) عند الشكل الهندسي المتوازن ، لمركب ثنائي التعويض (1H-1,2,3,-triazole-4-Yl) ethoxy) methy)-1H-1,2,3-triazole) , عندما تكون H , COOH , COCH₃ , and 2,3,4,6-Tetra-O-acetyl-β-D-glucopyranosyl = X , Y = C₇H₁₅.X , Y وكذلك الدوال الترموديناميكية (E⁰,H⁰,G⁰,A⁰) وكذلك تم حساب السعة الحرارية (S⁰ , C_p,C_v) , بعض الخصائص الفيزيائية وكذلك الشحنات . دلت الحسابات أن المركب R-sugar اكثر المشتقات استقرارية وله أعلى مقدار في كل من عزم ثنائي القطب ، الصلابة ، طاقة المدارين ، فرق الطاقة بين المدارين، جهد التأين وكذلك زيادة في جميع قيم الدوال الترموديناميكية .



X= H , COOH , COCH₃ , and 2,3,4,6-Tetra-O- acetyl-β-D-glucopyranosyl

Introduction.

Triazoles are five-membered heterocyclic aromatic rings, which contain two carbon and three nitrogen atoms. They are in isomeric forms 1,2,3-triazole and 1,2,4-triazole, and each of them is found in two tautomeric forms^[1].

Triazoles have received special attention in the medicinal fields because of low toxicity, high potency and broad range of biological activities such as antimicrobial^[2], antifungal^[3,4], antitumor^[5], and antitubercular activities^[6]

Click chemistry has been successfully approached in many different scientific fields, and its potential has played an important role in materials chemistry^[7], dendrimer build-up^[8,9], polymers^[10,11], nano particle synthesis^[12], peptide-based^[13] drug design and drug discovery^[14,15].

Click Chemistry is a biocompatible reaction and can take place in living cells thus it became a facile and efficient tool for universal modification of nucleosides^[16].


Glycoconjugates is the general classified for carbohydrates covalently linked with other chemical species such as proteins, peptides, lipids and saccharides^[17], thus an important role in many biological processes; glycoconjugates have been detected, contain cellular recognition, particularly in cases of inflammation^[18], tumor metastasis^[19], and immune response in bacterial and viral infections^[20].


The click chemistry has been extensively used on glycoconjugation chemistry, like the Synthesis of 1-glycopyranosyl-4-substituted-1,2,3-triazoles.

Glucose or galactose have many free hydroxyl groups which differ in the reactivity due to their position and configuration^[21]. Therefore, it is important to protect many hydroxyl groups in each carbohydrate and left just one single hydroxyl group in order to expose to the selected reagent before glycosylation for the preparation of glycoconjugates^[22,23].

COMPUTATIONAL


The compounds were investigated by conventional Ab Intio quantum theory and by HARTREE-FOCK (HF) using the Gaussian -03^[24,25].





1-(4-((2-(1-heptyl-1H-1,2,3-triazol-4-yl)ethoxy)methyl)-1H-1,2,3-triazol-1-yl)ethanone

R-COCH₃



(2R,3S,4S,5S)-2-(4-((2-(1-heptyl-1H-1,2,3-triazol-4-yl)ethoxy)methyl)-1H-1,2,3-triazol-1-yl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol

R-6-hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol.

Result and Discussion

Geometrical parameter

In this research calculated the geometry (bond lengths and bond angles) of the four molecules of derivatives 4- (2-(1H1,2,3, triazole (e-4-y1) ethoxymethyl)(1H-1,2,3- triazole) (R-H),(R-COOH) (RO-CH₃), R-CONH₂ , R-Suger using the Ab initio method of according to the Hartree – Fock method (RHF). According to the result calculated and recorded in the (table 1 and fig.1) Show that each the bonds (N₇-C_X) and (C₁₀ -N₉) in compound R-sugar has high value compared to other compounds studies this value may be return to the Ion pair in nitrogen atom and electron with drawing of substituted groups of the other compound. Where the bond (N₃-N₇) compound of (R-COOH) has high value compare with the other this value this caused by electron pair on (N) atom in this bound length that caused conjugated effect with substituted groups.

Also the change of the group substituted had effect on the value of the angles of the compounds studied in the research , have shown calculation in the (table 1 and fig.1). That the angle (∠C₆N₇N₃) showed that in R-OCH₃ and R-H larger than the others because the size of substituted group in the compounds when the angle (∠ N₄ N₃ N₇) the compound (R-OCH₃) has Larger value than the others because of the Ion pair on N with the electronegativity of the substituted groups.




Fig.(1): The geometric equilibrium for the 4- (2- (1H-1,2,3, - triazole (e-4-yl) ethoxymethyl (1H-1,2,3, triazole) derivatives.

Table (1) : Calculated geometric parameters (bond lengths in Angstrom length angles in degree) of the derivatives of(4-((2-(1H-1,2,3-triazol-4-ethoxy)methyl)-1-H-1,2,3,)Triazole calculated by (RHF)method.

| R-H | | R-COOH | | R-CO NH2 | | R-CO CH3 | | R-Sugar | |
|-----------|-----------------------|------------|-----------------------|------------|-----------------------|-----------|-----------------------|-----------|-----------------------|
| Para Geo | Bond Length and Angle | Para Geo | Bond Length and Angle | ParaGeo | Bond Length and Angle | Para Geo | Bond Length and Angle | ParaGeo | Bond Length and Angle |
| R(1-2) | 1.426 | R(1-2) | 1.423 | R(1-2) | 1.423 | R(1-2) | 1.424 | R(1-2) | 1.288 |
| R(3-4) | 1.284 | R(3-4) | 1.266 | R(3-4) | 1.227 | R(3-4) | 1.270 | R(4-5) | 1.354 |
| R(3-7) | 1.343 | R(3-7) | 1.370 | R(3-6) | 1.368 | R(3-7) | 1.363 | R(4-33) | 1.064 |
| R(4-5) | 1.369 | R(4-5) | 1.384 | R(4-5) | 1.473 | R(4-5) | 1.379 | R(5-12) | 1.455 |
| R(6-7) | 1.354 | R(6-7) | 1.374 | R(5-27) | 1.088 | R(6-7) | 1.373 | R(6-34) | 1.084 |
| R(6-17) | 1.062 | R(13-14) | 1.533 | R(6-23) | 1.380 | R(6-27) | 1.061 | R(6-35) | 1.085 |
| R(7-18) | 0.989 | R(13-19) | 1.531 | R(6-28) | 0.996 | R(7-22) | 1.411 | R(6-36) | 1.085 |
| R(8-9) | 1.282 | R(13-29) | 1.087 | R(7-8) | 1.288 | R(8-9) | 1.288 | R(7-8) | 1.533 |
| R(9-10) | 1.375 | R(14-15) | 1.533 | R(8-9) | 1.370 | R(9-10) | 1.369 | R(7-38) | 1.087 |
| A(4-3-7) | 107.2 | A(4-3-7) | 107.0 | A(4-3-6) | 120.9 | A(4-3-7) | 107.6 | A(2-3-4) | 107.1 |
| A(3-4-5) | 109.6 | A(3-4-5) | 110.5 | A(3-4-5) | 122.6 | A(3-4-5) | 110.1 | A(3-4-5) | 105.3 |
| A(3-7-6) | 111.0 | A(3-7-24) | 121.0 | A(7-6-27) | 125.5 | A(6-7-8) | 112.9 | A(6-7-8) | 112.9 |
| A(3-7-18) | 119.7 | A(6-7-24) | 129.1 | A(6-7-22) | 130.7 | A(6-7-37) | 109.4 | A(8-7-38) | 109.3 |
| A(7-6-17) | 124.4 | A(7-24-22) | 126.0 | A(7-22-23) | 121.9 | A(6-7-38) | 109.4 | A(7-8-9) | 113.2 |
| A(6-7-18) | 129.4 | | | | | | | A(7-8-39) | 109.2 |
| | | | | | | | | A(7-8-40) | 109.2 |

Physical properties.

Depending on the Ab initio of method of calculation according to the Hartree - Fock method (RHF), is calculate some physical properties of the molecules studied in this research ; Dipole moments (μ in Debye), energies (e V) of the high Occupied Molecular Orbital (E_{HOMO})and the Lower Unoccupied Molecular Orbital (E_{LUMO})and according Koopmans theorem (the negative E_{LUMO} is equal to the ionization potential) the calculation has been ionization energies (e V) , Also calculated the energy difference (ΔE , eV), And finally calculated (Molecular Hardness) Hardness (η) = $1/2(E_{HOMO} - E_{LUMO})$, (Electron Affinity) $E_A = - E_{LUMO}$ according Koopmans theorem[25-28]. Shown these results (table 2:) that compound R-CONH₂ has high value for each of (Dipole moments IP). And less value in E_{HOMO} (less the value of a negative energy), And the compound (R-COOH) has less value in ΔE (sense of activity of this compound is high) and (the compound R-Suger has high value hardness (η) Electron Affinity E_A).

Also, The MOPAC computational packages (semi-empirical method, AM1 model) employed to compute physical properties; heats of formation (ΔH_f^0 , kJ. mol⁻¹) [29,30] the results showed (Table 2:) for compound R-suger has lower heat of formation (more stability), Whereas the compound R-H has higher heat has of formation (less stability), Perhaps due this result to the effect of the group substitutes for the stability the compound .

Table 2: Calculated (ΔH_f^0 , (KJ mol⁻¹). (in Debye). Orbital energies (E_{HOMO}, E_{LUMO} , ΔE in ev), IP (in ev), E_A (in ev), and η -(in ev) for the derivatives of 4-(2-(1H1,2,3-triezo(e-4-y1) ethoxymethyl)-1H-1,2,3-triezoole.

| Comp. | R-H | R-COOH | R-CONH ₂ | R-COCH ₃ | R-Suger |
|----------------|----------|----------|---------------------|---------------------|-----------|
| ΔH_f^0 | 513.0275 | 590.7735 | 250.7611 | 232.6355 | -584.9466 |
| μ | 2.6706 | 6.1405 | 8.0186 | 3.3893 | 3.8743 |
| <i>HOMO</i> | -9.8073 | 9.7678 | -9.8345 | -9.51943 | -9.2527 |
| <i>LUMO</i> | 3.8393 | 2.3124 | 2.6518 | 2.8852 | 0.0714 |
| ΔE | 0.5015 | 0.4439 | 0.4588 | 12.4047 | 0.4896 |
| <i>IP</i> | 9.8073 | 9.7678 | 9.8345 | 9.51943 | 9.2527 |
| E_A | -3.839 | -2.3124 | -2.65177 | -2.8852 | -0.0714 |
| η | -6.8233 | -6.0401 | -6.2431 | -6.2044 | -4.6621 |

Thermodynamics functions


The fundamental vibration frequencies for the (R-H,R-COOH, R-COCH₃, R-CONH₂, R-Sugar) molecules along with the rotational constants, obtained in this study, were used to calculate the vibration and rotation contributions to the thermodynamic functions according to the statistical thermodynamic equations.

$$U_{vib}^0 = \sum_{i=1}^{3N-6} \frac{RTX_i}{e^{X_i} - 1}$$

$$X_i = \frac{1.44\bar{\nu}}{T}$$

$$S_{vib}^0 = R \sum_{i=1}^{3N-6} \left[\frac{X_i}{e^{X_i} - 1} - \ln(1 - e^{-X_i}) \right] \quad S_{rot}^0 = R \left[\frac{3}{2} + \ln \frac{8\pi^2 (8\pi^2 I_x I_y I_z)^{\frac{1}{2}} (KT)^{\frac{3}{2}}}{\sigma \hbar^3} \right]$$

These two contributions along with the others contribution, for the translation, electronic, and nuclear motions, were used to calculate E^0 , H^0 , S^0 , A^0 , and G^0 thermodynamic functions. Standard thermodynamic functions and heat capacity for the studied molecules listed Table: 3 looking at the calculation results show that each of the thermodynamic functions (G^0 , H^0 , A^0 , E^0 , S^0 , C_p , C_v) have the same gradient values (different the group substitutes)



The deferent thermodynamic functions values due to deferent substituted groups (X), viz, the nitrogen atom that has Ion pair and a high electro-negative in addition to its high size which makes restricted movement. (-COOH,-suger,-OCH₃,-ONH₂,-H).

Table 3: The calculated standard thermodynamics functions at 298. K of the derivatives of 4-(2-(1H1,2,3,-triazole(e-4-y1) ethoxy methyl)(1-H-1,2,3-traizole.

| COMP | R-H | R-COOH | R-COCH ₃ | R-CONH ₂ | R-Suger |
|-----------------------------|----------|----------|---------------------|---------------------|----------|
| E° KJ/mol | 150.465 | 1195.661 | 1313.085 | 1230.213 | 1672.369 |
| S° KJ/mol..K | 0.4554 | 0.7015 | 0.7854 | 0.6881 | 0.91985 |
| C _v KJ/mol..K | 0.1590 | 0.3269 | 0.3585 | 0.3243 | 0.4768 |
| H° KJ/mol | 2628.035 | 3973.321 | 3790.655 | 3707.783 | 4149.939 |
| C _p KJ/mol..K | 8.473 | 8.6409 | 8.6725 | 8.6383 | 8.7908 |
| A° KJ/mol | 2356.617 | 3255.137 | 322.557 | 3297.677 | 3601.709 |
| G° KJ/mol | 2492.326 | 3464.184 | 3556.606 | 3502.73 | 3875.824 |

The Charges

The Calculated for all charges atoms of the molecules studied according to the method (RHF), have shown calculation results of the charges (**Table 4:**) are each of the (N₇, O₁) has high negative value in compound (R-CONH₂) than the other's Perhaps the reason for this difference in electrical negative of substituted group . This scientific fact indicates that the compound (R-CONH₂) can be strong legend when it linked to metal complex formation , as a results founds that the amount of charge of these atoms

R-CONH₂> R-Sugar > R-H > R-COOH > R-COCH₃

Table:4 Calculated charge for the derivatives of 4-(2-(1H1,2,3,-triazole(e-4-y1) ethoxy methyl)(1-H-1,2,3-traizole calculated by (RHF)method .

| <i>R-H</i> | | <i>R-COOH</i> | | <i>R-CONH2</i> | | <i>R-COCH3</i> | | <i>R-Sugar</i> | |
|-------------|---------------|---------------|---------------|----------------|---------------|----------------|---------------|----------------|---------------|
| <i>Atom</i> | <i>Charge</i> | <i>Atom</i> | <i>Charge</i> | <i>Atom</i> | <i>Charge</i> | <i>Atom</i> | <i>Charge</i> | <i>Atom</i> | <i>Charge</i> |
| O1 | -0.7558 | O1 | -0.7556 | O1 | -0.7602 | O1 | -0.7435 | N1 | 0.0427 |
| C2 | 0.0957 | C2 | 0.1016 | C2 | 0.0994 | C2 | 0.0606 | N2 | -0.3571 |
| N3 | 0.0428 | N3 | 0.1231 | N3 | 0.1311 | N3 | 0.0745 | C3 | 0.0485 |
| N4 | -0.3556 | N4 | -0.3172 | N4 | -0.3294 | N4 | -0.2651 | C4 | 0.17668 |
| C5 | 0.0176 | C5 | -0.0028 | C5 | 0.0073 | C5 | -0.2177 | N5 | -0.7075 |
| C6 | 0.2039 | C6 | 0.3050 | C6 | 0.2771 | N6 | -0.7638 | C6 | -0.4514 |
| N7 | -0.6985 | N7 | -0.8599 | N7 | -0.8888 | N7 | 0.0418 | C7 | -0.2997 |
| N8 | 0.0517 | N8 | 0.0533 | N8 | 0.0539 | N8 | -0.3571 | C8 | -0.2903 |
| N9 | -0.3477 | N9 | -0.3566 | N9 | -0.3574 | C9 | 0.0478 | C9 | -0.2951 |
| C10 | 0.0374 | C10 | 0.0389 | C10 | 0.0386 | C10 | 0.1782 | C10 | -0.3114 |
| C11 | 0.1575 | C11 | 0.1765 | C11 | 0.1774 | N11 | -0.7077 | C11 | -0.2938 |
| N12 | -0.6999 | N12 | -0.7323 | N12 | -0.7321 | C12 | -0.2997 | C12 | -0.0341 |
| C13 | -0.2784 | C13 | -0.2997 | C13 | -0.4515 | C13 | -0.2903 | C13 | -0.2987 |
| C14 | 0.0589 | C14 | -0.2903 | C14 | -0.2997 | C14 | -0.2951 | C14 | 0.0699 |
| H15 | 0.1777 | C15 | -0.2946 | C15 | -0.2903 | C15 | -0.3113 | O15 | -0.7561 |
| H16 | 0.1777 | C16 | -0.3133 | C16 | -0.2946 | C16 | -0.2938 | C16 | 0.1016 |
| H17 | 0.2891 | C17 | -0.3082 | C17 | -0.3133 | C17 | -0.0339 | N17 | 0.0413 |
| H18 | 0.4203 | C18 | -0.0034 | C18 | -0.3083 | C18 | -0.4515 | N18 | -0.3504 |
| H19 | 0.2678 | C19 | -0.4515 | C19 | -0.0037 | C19 | -0.2957 | C19 | -0.0022 |
| H20 | 0.4209 | C20 | -0.2773 | C20 | -0.2776 | C20 | 0.0532 | C20 | 0.3046 |
| H21 | 0.2081 | C21 | 0.0563 | C21 | 0.0569 | O21 | -0.5737 | N21 | -0.7805 |
| H22 | 0.2081 | O22 | -0.4850 | C22 | 1.0856 | C22 | -0.5159 | O22 | -0.7228 |
| H23 | 0.1503 | O23 | -0.7282 | O23 | -0.5323 | C23 | 0.7638 | C23 | 0.4923 |
| H24 | 0.1503 | C24 | 1.0697 | N24 | -0.9487 | H24 | 0.1696 | C24 | 0.1695 |
| | | H25 | 0.1819 | H25 | 0.1835 | H25 | 0.1787 | C25 | 0.1258 |
| | | H26 | 0.1819 | H26 | 0.1835 | H26 | 0.2095 | O26 | -0.8082 |
| | | H27 | 0.3239 | H27 | 0.2911 | H27 | 0.1888 | O27 | -0.7867 |
| | | H28 | 0.2583 | H28 | 0.2592 | H28 | 0.4098 | O28 | -0.7679 |
| | | H29 | 0.1515 | H29 | 0.1540 | H29 | 0.2623 | C29 | 0.0427 |
| | | H30 | 0.1515 | H30 | 0.1491 | H30 | 0.1519 | C30 | 0.1445 |
| | | H31 | 0.1472 | H31 | 0.1491 | H31 | 0.1499 | C31 | 0.1519 |
| | | H32 | 0.1472 | H32 | 0.1515 | H32 | 0.1471 | O32 | -0.8040 |
| | | H33 | 0.1554 | H33 | 0.1515 | H33 | 0.1498 | H33 | 0.2629 |
| | | H34 | 0.1554 | H34 | 0.1474 | H34 | 0.1562 | H34 | 0.1537 |
| | | H35 | 0.1503 | H35 | 0.1474 | H35 | 0.1508 | H35 | 0.1491 |
| | | H36 | 0.1503 | H36 | 0.1555 | H36 | 0.1497 | H36 | 0.1503 |
| | | H37 | 0.1916 | H37 | 0.1555 | H37 | 0.1582 | H37 | 0.1518 |
| | | H38 | 0.1916 | H38 | 0.1505 | H38 | 0.1894 | H38 | 0.1498 |
| | | H39 | 0.1886 | H39 | 0.1505 | H39 | 0.1606 | H39 | 0.1472 |
| | | H40 | 0.1887 | H40 | 0.1916 | H40 | 0.2152 | H40 | 0.1499 |
| | | H41 | 0.1539 | H41 | 0.1916 | H41 | 0.1824 | H41 | 0.1562 |

References

- 1- A.Chawala and P.Kaur, *International Research Journal of Pharmacy IRJP*, 4(1), **2013**, 49-72.
- 2- V. L. Campo , R. Sesti-Costa, Z. A. Carneiro, J. S. Silva , S. Schenkman and I. Carvalho, *Bioorg. Med. Chem.*, 20 (1), **2012**, 145–156.
- 3- R.G. Lima-Neto, N.M. Cavalcante, R.M. Srivastava, F.M. Junior, A.G. Wanderley, R.P. Neves and J.V. dos Anjos, *Molecules*, 17 (5), **2012**, 5882-5892.
- 4- S. Pearson, W. Scarano and M.H. Stenzel, *Chem. Commun.*, **48(39)**, **2012**, 4695-4697.

- 5- L. Ou, S. Han, W. Ding, P. Jia, B. Yang, J. L. Medina-Franco, M. A. Giulianotti, J.Z. Chen, Y. Yu, *Mol. Diversity*, 15(4), **2011**, 927-946.
- 6- G.V. Kumar, Y. Rajendraprasad, B.P. Mallikarjuna, S.M. Chandrashekar and C. Kistayya , *Eur J MedChem.*, 45(5), **2010**, 2063–2074.
- 7- J.E. Hein and V.V.Foken, *Chem. Soc. Rev.*, 39(4),**2010**, 1302-1315.
- 8- W.H. Binder and R. Sachsenhofer, *Macromol. Rapid Commun.*,29(16),**2008**, 952-981.
- 9- J.A. Johnson, M.G. Finn, J.T. Koberstein and N.J. Turro,*Macromol. Rapid Commun.*, 29(12-13), **2008**,1052-1072.
- 10- D.Kushwaha and V.K. Tiwari,*J. Org. Chem.*, 78(16), **2013**, 8184–8190.
- 11- J. Lu, M.Shi and M.S.Shoichet, *Bioconjugatechem.*, 20(1), **2009**, 87-94.
- 12- H. Li, R. Aneja and I. Chaiken, *Molecules*, 18(8), **2013**, 9797-9817.
- 13- V.G.Nenajdenko, A.V.Gulevich, N.V.Sokolova, A.V.Mironov and E.S.Balenkova, *Eur.J. Org. Chem.*, 2010(8), **2010**, 1445-1449.
- 14- N.Li and W.H.Binder,*J. Mater. Chem.*, 21, **2011**, 16717-16734.
- 15- M.Reithofer, K.Chan, A.Lakshmanan, D.H.Lam, A.Mishra, B.Gopalan, M.Joshi, S. Wang and C.A.E. Hauser,*Chem. Sci.*, **2014**, Advance Article,DOI: 10.1039/C3SC51930A.J. Krim, M.Taourirte, C. Grünewald, I. Krstic and J.W. Engels, *Synthesis*, 45(3), **2013**, 396-405.
- 16- S.P.O. Assis, M.T. Silva, R. N. de Oliveira, and V.L.M. Lima,*The Scientific World Journal*, 2012,**2012**, 1-7.
- 17- B.L. Wilkinson, L.F. Bornaghi, T.A. Houston, A. Innocenti, D.Vullo, C.T. Supuranand S. A. Poulsen, *J. Med. Chem.*,50(7), **2007**, 1651 –1657.
- 18- A. Mohammed, A. Dawood, K. Ali and M. Al-Mosawi, *Asian J. Chem.*, 24(12),**2012**, 5592-5596.
- 19- H.M.I. Osborn, *Best Synthesis Methods: Carbohydrates*,Elsevier science Ltd., Oxford, UK, **2003**, p 10.
- 20- E.Bokor,C. Koppány, Z. Gonda, Z.Novák and L. Somsák,*Carbohydr. Res.* 351(7),**2012**, 42–48.
- 21- Y. Sawada, N. Nanboku, E. Yanase and S. Nakatsuka, *Heterocyclic Communications*, 16(1), **2011**, 21–24.
- 22- J. Guo and X.Ye,*Molecules*, 15(10), **2010**, 7235-7265.
- 23- W.J.Hehre,L.Radom,P.v,RSchleyer and J.A.Pople, *AbIntio Molecular Orbital Theory*.Wiley, New York(1989).
- 24- M.J. Frish, *Gaussian 03,RevisionB.03*, Gaussian Pittsburgh,PA., 2003.
- 25- Moore C. E. ,*"Ionization Potentials and Ionization Limits Derived from The Analyses of Optical Spectra"*, NSRDS-NBS, USA,**1970** .
- 26- Vanfleteren D., Neck D. V., Ayers P. W., Morrison R. C. and Bultinck P.,*The Journal of Chemical Physics*,**130**, **2009**,194104(1)-194104(10).
- 27- Hotop H. and Lineberger W. C., *Journal of Physical Chemistry*, 4, 1975,539-576.
- 28- Zhan C. G., J. A. Nicholsand ,D. A. Dixon, *Journal of Physical Chemistry A*, 107, **2003**. 4184-4195.
- 29- Pearson ,R.G., *"Chemical Hardness"*, wiley-VCH, Germany; **1997**.
- 30- Pearson R.G., *Proceedings of the National Academy of Sciences of the United States of America*, 83, **1986**.8440-8441.