

**Purely Quasi-Dedekind Modules And Purely Prime Modules**

**Tha'ar Younis Ghawi**

**Department of Mathematics , College of Education**

**University of AL-Qadisiya \ IRAQ**

**E-mail : thaar\_math83@yahoo.com**

**Abstract :-**

An  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called pure if  $IN = N \cap IM$  for every ideal  $I$  of  $R$ . In this paper we introduce the notion of purely quasi-invertible submodule and a purely quasi-Dedekind module, where an  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called purely quasi-invertible if,  $N$  is pure and  $Hom_R(M/N, M) = 0$ . And an  $R$ -module  $M$  is called purely quasi-Dedekind if, every nonzero pure submodule  $N$  of  $M$  is quasi-invertible ; that is  $Hom_R(M/N, M) = 0$ . Beside these, we also introduce the notion of purely prime module, where an  $R$ -module  $M$  is called purely prime module if  $ann_R M = ann_R N$  for all nonzero pure submodule  $N$  of  $M$ . We gave many properties related with this concepts. And we studied the relationships between these concepts and several other types of modules. In this paper  $R$  is a commutative ring with unity and  $M$  is a unitary  $R$ -module .

**0. Introduction:-**

Let  $R$  be a ring and  $M$  be a unital  $R$ -module. If  $N$  is a submodule of  $M$ , we write  $N \leq M$  and if  $N$  is an essential submodule of  $M$  then we write  $N \leq_e M$ , also if  $N$  is a direct summand of  $M$  then we write  $N \leq^{\oplus} M$ . Recall an  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called pure if  $IN = N \cap IM$  for every ideal  $I$  of  $R$  [5], [10], and  $N$  is called quasi-invertible if,  $Hom_R(M/N, M) = 0$  [14]. And an  $R$ -module  $M$  is called quasi-Dedekind if, each nonzero submodule of  $M$  is quasi-invertible [14]. And an  $R$ -module  $M$  is called prime module if  $ann_R M = ann_R N$  for all nonzero submodule  $N$  of  $M$  [8]. Ghawi Th.Y. in [11] introduced the concepts of essentially quasi-invertible submodules and essentially quasi-Dedekind modules as a

**Key Words : Purely quasi-invertible Submodules; Pure Submodules; Purely quasi-Dedekind Modules; Purely prime Modules .**

generalization of quasi-invertible submodules and quasi-Dedekind modules, where a submodule  $N$  of an  $R$ -module  $M$  is called essentially quasi-invertible if  $N \leq_e M$  and  $N$  is quasi-invertible and

$M$  is called essentially quasi-Dedekind if every essential submodule of  $M$  is quasi-invertible . This paper has been organized on three sections. In section 1, we generalized the concept of quasi-invertible submodule to a purely quasi-invertible submodule, where a submodule  $N$  of a module  $M$  is called purely quasi-invertible if  $N$  is a pure and quasi-invertible submodule. We give some basic properties of this class of submodules.

In section 2, we introduce the concept of a purely quasi-Dedekind module as a generalization to concept a quasi-Dedekind module, where an  $R$ -module  $M$  is called purely quasi-Dedekind if, every nonzero pure submodule of  $M$  is quasi-invertible . We prove that if  $M$  a purely quasi-Dedekind module with  $M/K$  is projective for all pure submodule  $K$  of  $M$  then  $M/N$  is a purely quasi-Dedekind module, for all  $N \leq M$  . Also, we show by an example a direct sum of purely quasi-Dedekind modules need not be a purely quasi-Dedekind module (see Ex 2.14) . On the other hand we give a condition under which the direct sum of purely quasi-Dedekind modules is a gain purely quasi-Dedekind ( see Prop 2.15) . Finally, in section 3, we introduce and study the concept purely prime module as a generalization of prime module, where an  $R$ -module  $M$  is called a purely prime module if  $ann_R M = ann_R N$  for all nonzero pure submodule  $N$  of  $M$  . We see that every prime module is a purely prime module, but the converse is not true. Also we give some equivalent formulas and results of this concept .

### 1. Purely Quasi-Invertible Submodules

Firstly, we recall that an  $R$ -submodule  $N$  of an  $R$ -module  $M$  is pure if,  $IN = N \cap IM$  for every ideal  $I$  of  $R$  [5], [10] . Mijbass A.S. in [14] introduced the following concept, an  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called quasi-invertible if,  $Hom_R(M/N, M) = 0$ . And an ideal  $J$  of a ring  $R$  is called quasi-invertible if  $J$  is a quasi-invertible  $R$ -submodule. In this section we introduce and study a generalization of the concept a quasi-invertible submodule namely " purely quasi-invertible " .

**Definition 1.1.** An  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called purely quasi-invertible if  $N$  is pure and  $Hom_R(M/N, M) = 0$ . And an ideal  $I$  of a ring  $R$  is called purely quasi-invertible if  $I$  is a purely quasi-invertible  $R$ -submodule . It is clear that every purely quasi-invertible submodule is a quasi-invertible submodule . The following example shows that the converse is false .

**Example 1.2.** Let  $R$  be an integral domain and let  $\bar{R} = R[x, y]$  be the polynomial ring of two independent variables  $x$  and  $y$  , then  $\bar{R}$  is also an integral domain . Let  $I = (x, y)$  is the ideal of  $\bar{R}$  generated by  $x$  and  $y$  , so by [14, Ex 1.3(1), P.6]  $I$  is quasi-invertible. But  $I$  is not pure of  $\bar{R}$  , thus  $I$  is not purely quasi-invertible; To see this: Let  $R = \mathbb{Z}$  ,  $\bar{R} = \mathbb{Z}[x, y]$ , let  $I = (x, y) = \{xf_1 + yf_2 : f_1, f_2 \in \bar{R}\}$  , thus by [14, Ex 1.3(1), P.6]  $I$  is quasi-invertible. Now, Let  $J = \{f \in \bar{R} : f(x, y) = a, a \in 2\mathbb{Z}\}$  then  $J \cap I = \{axf_1 + ayf_2 : f_1, f_2 \in \bar{R}\} \neq \{0\} = I \cap J$ ; that is  $I$  not pure, hence  $I$  is not purely quasi-invertible .

Remarks and Examples 1.3.

- 1) In any nonzero module  $M$ .  $0$  is not purely quasi-invertible, but  $M$  is a purely quasi-invertible submodule .
- 2) If  $N$  is a proper direct summand of an  $R$ -module  $M$  then  $N$  is pure by [21], but not quasi-invertible, since there exists  $0 \neq K \leq M$  such that  $M = K \oplus N$  and  $Hom_R(M/N, M) = Hom_R(K \oplus N/N, K \oplus N) = Hom_R(K, K \oplus N) \neq 0$ .

Recall that an  $R$ -module  $M$  is called semisimple if, every submodule of  $M$  is a direct summand of  $M$  [ 12, P.189] .

- 3) If  $M$  is a semisimple module, then  $M$  is the only purely quasi-invertible submodule of  $M$  ; since every proper submodule of  $M$  is direct summand; that is pure not quasi-invertible (see Rem.and.Ex 1.3(2)) .
- 4) Let  $M = Z_4$  as  $Z$ -module ,  $N = (\bar{2})$  is not a purely quasi-invertible submodule of  $Z_4$  as  $Z$ -module . In fact  $N$  is not quasi-invertible , since  $Hom_Z(Z_4/(\bar{2}), Z_4) \cong Z_2 \neq 0$ . Also,  $N$  is not pure, since  $\bar{2} = \bar{2} \cdot \bar{1} \in (\bar{2}) \cap 2(Z_2)$  but  $\bar{2} \notin 2(\bar{2})$  .
- 5) If  $N$  is a purely quasi-invertible  $R$ -submodule of an  $R$ -module  $M$ , then  $ann_R M = ann_R N$  .

**Proof.** Follows by [14, Prop 1.4, P.7] .  $\square$

However, the converse of (Rem.and.Ex 1.3(5)) is not true as the following example shows:

Consider  $Z$ -module  $Z \oplus Z_4$ , let  $N = 2Z \oplus Z_4 \leq Z \oplus Z_4$ , then  $ann_Z(Z \oplus Z_4) = ann_Z(2Z \oplus Z_2) = 0$  but  $N = 2Z \oplus Z_4$  is not purely quasi-invertible of  $Z \oplus Z_4$  as  $Z$ -module. In fact  $N$  is not pure, since  $(2, \bar{2}) = 2(1, \bar{1}) \in (2Z \oplus Z_4) \cap 2(Z \oplus Z_4)$  but  $(2, \bar{2}) \notin 2(2Z \oplus Z_4)$  .

- 6) Let  $I$  be an ideal of a ring  $R$  . If  $I$  is purely quasi-invertible then  $ann_R(I) = 0$  .

**Proof. Obvious .**  $\square$

The converse of (Rem.and.Ex 1.3(6)) is not true in general, consider the following example: Let  $R = Z$ , let  $I = 2Z$  then  $ann_Z(I) = ann_Z(2Z) = 0$ , but  $I$  is not pure of  $Z$ , since  $J = 4Z$  be an ideal of  $Z$  and  $J \cap I = (4Z) \cap (2Z) = 8Z \neq 4Z = (2Z) \cap (4Z) = I \cap JZ$  , so it is not purely quasi-invertible ideal of  $Z$ .

7) If  $M = M_1 \oplus M_2$  is an  $R$ -module and let  $K$  be a purely quasi-invertible in  $M_i$  for some  $i = 1, 2$ , then it is not necessarily that  $K$  is a purely quasi-invertible submodule of  $M$ ; For example: In the  $Z$ -module  $Z \oplus Z_2$ ,  $K = Z_2$  is a purely quasi-invertible submodule of  $Z_2$  as  $Z$ -module, but  $Z_2 \cong (0) \oplus Z_2$  which is not a purely quasi-invertible submodule of  $Z \oplus Z_2$  as  $Z$ -module, since  $\text{Hom}_Z(Z \oplus Z_2 / (0) \oplus Z_2, Z \oplus Z_2) = \text{Hom}_Z(Z, Z \oplus Z_2) \neq 0$ ; that is  $(0) \oplus Z_2$  not quasi-invertible.

Remark 1.4. We do not whether the intersection of purely quasi-invertible submodules is purely quasi-invertible.

Recall that an  $R$ -module  $M$  has the pure intersection property (briefly *PIP*) if, the intersection of any two pure submodules is again pure [3, def 2.1, P.33].

Now we can introduce the following result.

Proposition 1.5. Let  $M$  be an  $R$ -module has *PIP*. If  $N_1, N_2$  are purely quasi-invertible submodules of  $M$  then  $N_1 \cap N_2$  is also.

Proof. Since  $M$  has *PIP* then  $N_1 \cap N_2$  is pure in  $M$ . But it is easy to see that  $\text{Hom}(M/N_1 \cap N_2, M) \subseteq \text{Hom}(M/N_1, M) + \text{Hom}(M/N_2, M)$ . Hence  $\text{Hom}(M/N_1 \cap N_2, M) = 0$  and so that  $N_1 \cap N_2$  is a purely quasi-invertible submodule of  $M$ .  $\square$

Recall that an  $R$ -module  $M$  is called multiplication if, for each submodule  $N$  of  $M$ ,  $N = IM$  for some ideal  $I$  of  $R$ . Equivalently,  $M$  is multiplication if, for each submodule  $N$  of  $M$ ,  $N = [N : M].M$ , where  $[N : M] = \{r \in R : rM \subseteq N\}$  [19].

Corollary 1.6. Let  $M$  be a multiplication  $R$ -module. If  $N_1, N_2$  are purely quasi-invertible submodules of  $M$  then  $N_1 \cap N_2$  is also.

Proof. Follows by [3, Prop 2.3, p.33] and (Prop 1.5).  $\square$

However, the following results (1.5), (1.6) gives necessary conditions for make (Rem 1.4) is true.

Remark 1.7. Let  $M$  be an  $R$ -module and let  $N$  be a purely quasi-invertible submodule of  $M$ . If  $K \leq M$  such that  $K \cong N$  then it is not necessarily that  $K$  is a purely quasi-invertible submodule of  $M$ . We can give the following example show that.

Example 1.8. Let  $M = Z$  as  $Z$ -module, let  $N = Z$  be a submodule of  $M$ , then  $N$  is a purely quasi-invertible submodule of  $M$ , but  $K = 2Z \cong Z = N$  is not a purely quasi-quasi-invertible submodule of  $M$ . In the fact  $K = 2Z$  is not pure in  $M$ .

Remark 1.9. Let  $M_1, M_2$  be  $R$ -modules and let  $f : M_1 \longrightarrow M_2$  be  $R$ -homomorphism. If  $N$  is a purely quasi-invertible submodule of  $M_1$  then not necessary that the image of  $N$  is a purely quasi-invertible submodule of  $M_2$ . For example : Consider  $Z$ -modules  $Z_4, Z_6$ . Let  $f : Z_6 \longrightarrow Z_4$  be  $Z$ -homomorphism define by  $f(\bar{x}) = 2\bar{x}$  for all  $\bar{x} \in Z_6$ . Let  $N = Z_6$ , it is well known that  $N$  is a purely quasi-invertible submodule of  $Z_6$  as  $Z$ -module, but  $f(N) = f(Z_6) = \{\bar{0}, \bar{2}\} = \langle \bar{2} \rangle$  is not purely quasi-invertible submodule of  $Z_4$  as  $Z$ -module (see Rem.and.Ex 1.3(4)).

Recall that a nonzero  $R$ -module  $M$  is called a rational extension of the  $R$ -submodule  $N$  of  $M$  if, for all  $m_1, m_2 \in M, m_2 \neq 0$ , there exists an element  $r \in R$  such that  $rm_1 \in N$  and  $rm_2 \neq 0$  [20]. And recall that an  $R$ -module  $M$  is regular if for all  $a \in M$  and for all  $r \in R$ , there exists  $x \in R$  such that  $rxra = ra$ . Equivalently, every submodule of  $M$  is pure [7].

Proposition 1.10. Let  $M$  be a module over regular ring  $R$  and let  $N \leq M$ . If  $M$  is a rational extension of  $N$  then  $N$  is a purely quasi-invertible submodule of  $M$ .

Proof. Since  $M$  is a rational extension of  $N$  then by [14, Prop 3.3, P.14]  $N$  is a quasi-invertible submodule of  $M$ . On the other hand, since  $R$  is a regular ring then  $M$  is a regular  $R$ -module; that is every submodule of  $M$  is pure, thus  $N$  is a purely quasi-invertible submodule of  $M$ .  $\square$

Recall that an  $R$ -submodule  $N$  of an  $R$ -module  $M$  is called small (briefly  $N \ll M$ ) if, for all  $K \leq M$  with  $N+K = M$  implies  $K = M$  [12, P.106]. And recall that an  $R$ -submodule  $N$  of  $R$ -module  $M$  is called *SQI*-submodule if, for each  $f \in \text{Hom}_R(M/N, M)$  then  $f(\frac{M}{N})$  is a small in  $M$  [17, p.44].

Remark 1.11. It is clear that every quasi-invertible submodule is *SQI*-submodule, hence every purely quasi-invertible submodule is *SQI*-submodule. But the converse is not true in general, the following example shows.

Example 1.12. Let  $M = Z_4$  as  $Z$ -module and let  $N = \langle \bar{2} \rangle \leq M$ . Then  $N$  is *SQI*-submodule of  $Z_4$ , but it is known that  $N$  is not a purely quasi-invertible submodule of  $Z_4$  (See Rem.and.Ex 1.3(4)).

We end this section by the following theorem.

Theorem 1.13. Let  $M$  be a faithful multiplication over integral domain  $R$ . If  $N$  is a pure submodule of  $M$  then  $[N : M]$  is a purely quasi-invertible ideal of  $R$ .

Proof. Assume that  $N$  is a pure submodule of  $M$ . Since  $M$  be a faithful multiplication  $R$ -module, so by [4, Coro 1.2, P.65]  $[N : M]$  is a pure ideal of  $R$ . But  $R$  is an integral domain, hence by [14, Ex 1.3(1), P.6] every nonzero ideal of  $R$  is quasi-invertible, thus  $[N : M]$  is a quasi-invertible ideal of  $R$ . Hence  $[N : M]$  is a purely quasi-invertible ideal of  $R$ .  $\square$

## 2. Purely Quasi-Dedekind Modules

Recall that an  $R$ -module  $M$  is called quasi-Dedekind if, every nonzero submodule of  $M$  is quasi-invertible; that is  $Hom_R(M/N, M) = 0$  for all nonzero submodule  $N$  of  $M$  [14, P.24]. In this section we give generalization of the concept a quasi-Dedekind module namely "purely quasi-Dedekind module". We list some basic properties of purely quasi-Dedekind modules. Also we give a characterization of this concept. We study the relationships between a purely quasi-Dedekind modules with other related modules. We begin with the following definition :

**Definition 2.1.** An  $R$ -module  $M$  is said to be purely quasi-Dedekind if, every proper nonzero pure submodule of  $M$  is quasi-invertible. And a ring  $R$  is called purely quasi-Dedekind if  $R$  is a purely quasi-Dedekind  $R$ -module.

It is clear that every quasi-Dedekind  $R$ -module is a purely quasi-Dedekind  $R$ -module. But the converse may not be, as the following example shows :

**Example 2.2.** Consider  $Z$ -module  $Z_4$ , it is clear that  $Z_4$  is purely quasi-Dedekind, since  $Z_4$  as  $Z$ -module has no proper pure submodule. But it is not quasi-Dedekind, since  $(\bar{2}) \leq Z_4$  and  $Hom_Z(Z_4/(\bar{2}), Z_4) \cong Z_2 \neq 0$ .

### Remarks and Examples 2.3.

- 1) Every simple  $R$ -module is a purely quasi-Dedekind  $R$ -module.
- 2) Every nonzero semisimple and (not simple) module is not a purely quasi-Dedekind module. In particular  $Z_6$  as  $Z$ -module is semisimple and (not simple) but it is not purely quasi-Dedekind.
- 3) Every integral domain  $R$  is a quasi-Dedekind  $R$ -module [14, Ex 1.4(1), P.24], so it is a purely quasi-Dedekind  $R$ -module. But the converse need not be in general; For example: Let  $M = Z_4$  as  $Z_4$ -module, then  $Z_4$  is purely quasi-Dedekind, but  $Z_4$  is not an integral domain.
- 4)  $Z$  as  $Z$ -module is purely quasi-Dedekind.  $0, Z$  are the only pure submodules of  $Z$ .
- 5) Let  $M$  be a regular  $R$ -module. Then  $M$  is purely quasi-Dedekind if and only if  $M$  is quasi-Dedekind.

Proof. Clear .  $\square$

6) Let  $M$  be a module over regular ring  $R$ . Then  $M$  is purely quasi-Dedekind if and only if  $M$  is quasi-Dedekind .

Proof. Follows by (Rem.and.Ex 2.3(5)) and since every module over a regular ring is regular .  
 $\square$

7) If  $M$  is a purely quasi-Dedekind  $R$ -module then  $ann_R N = ann_R M$  for all nonzero pure submodule  $N$  of  $M$  .

Proof. Follows by ( Rem.and.Ex 1.3(5)) .  $\square$

Proposition 2.4. Let  $M$  be an  $R$ -module with  $\bar{R} = R/J$  , where  $J$  is an ideal of  $R$  such that  $J \subseteq ann_R M$  .  $M$  is a purely quasi-Dedekind  $R$ -module if and only if  $M$  is a purely quasi-Dedekind  $\bar{R}$  -module .

Proof. We have by [12, P.51]  $Hom_R(M/N, M) = Hom_{\bar{R}}(M/N, M)$  for all submodule  $N$  of  $M$  . Thus the result is obtained .  $\square$

Proposition 2.5. Let  $M$  be a uniform  $R$ -module with  $ann_R M$  is a maximal ideal of  $R$ , then  $M$  is a purely quasi-Dedekind  $R$ -module .

Proof. Follows by [11, Coro 1.2.10 and (Rem.and.Ex 1.2.2(5))] .  $\square$

Theorem 2.6. Let  $M$  be an  $R$ -module. If  $M$  is purely quasi-Dedekind then for all  $f \in End_R(M)$  and  $Ker f$  is a pure submodule of  $M$  implies  $f = 0$  .

Proof. Let  $f \in End_R(M)$  and  $Ker f$  is a pure submodule of  $M$  . Suppose that  $f \neq 0$ , define  $g : M/Ker f \longrightarrow M$  by  $g(m + Ker f) = f(m)$  for all  $m \in M$  . It is easy to see that  $g$  is Well-defined and  $g \neq 0$  ( since  $f \neq 0$ ) . Hence  $Hom_R(M/Ker f, M) \neq 0$  which is a Contradiction.  $\square$

Proposition 2.7. Let  $M$  be an  $R$ -module such that for all pure submodule  $N$  of  $M$ , and for all  $K \leq M$  such that  $N \leq K \leq M$  implies  $K$  is pure in  $M$  . If for all  $f \in End_R(M)$  ,  $Ker f$  is a pure submodule of  $M$  implies  $f = 0$  , then  $M$  is a purely quasi-Dedekind  $R$ -module .

Proof. Suppose that there exists  $0 \neq N \leq M$  ,  $N$  is pure such that  $Hom_R(M/N, M) \neq 0$ ; that is there exists  $R$ -homomorphism  $f : M/N \longrightarrow M$  and  $f \neq 0$  . Now, consider the following diagram :  $M \xrightarrow{\pi} M/N \xrightarrow{f} M$  , where  $\pi$  is the canonical projection map. Let  $\phi = fo\pi$  , so  $\phi \in End_R(M)$  , but  $N \subseteq Ker \phi$  and  $N$  is a nonzero pure submodule of  $M$ , thus  $Ker \phi$  is a

nonzero pure submodule of  $M$  (by hypothesis) . On the other hand  $\phi(M) = f(M/N) \neq 0$  which is a contradiction .  $\square$

We will need the following lemma for the proof next proposition .

Lemma 2.8. Let  $M_1, M_2$  be  $R$ -modules and let  $f : M_1 \longrightarrow M_2$  be  $R$ -epimorphism . If  $N$  is a pure submodule of  $M_2$  then  $f^{-1}(N)$  is a pure submodule of  $M_1$  .

Proof. Assume that  $I$  is an ideal of  $R$  , then  $I f^{-1}(N) = f^{-1}(IN) = f^{-1}(N \cap IM_2) =$

$f^{-1}(N) \cap f^{-1}(IM_2) = f^{-1}(N) \cap I f^{-1}(M_2) = f^{-1}(N) \cap I M_1$  , since  $f$  is epimorphism . Thus  $f^{-1}(N)$  is a pure submodule of  $M_1$  .  $\square$

Now, we can introduce the following proposition .

Proposition 2.9. Let  $M_1, M_2$  be  $R$ -modules such that  $M_1$  is isomorphic to  $M_2$  . Then  $M_1$  is purely quasi-Dedekind if and only if  $M_2$  is purely quasi-Dedekind .

Proof. Suppose that  $M_1$  is a purely quasi-Dedekind  $R$ -module. Since  $M_1 \cong M_2$  , so there exists  $f : M_1 \longrightarrow M_2$  be  $R$ -isomorphism. Let  $N$  be a nonzero pure submodule of  $M_2$ , thus

by above lemma  $f^{-1}(N)$  is a nonzero pure submodule of  $M_1$ , so  $\text{Hom}_R(M_1/f^{-1}(N), M_1) = 0$  .

But  $\text{Hom}_R(M_2/N, M_2) \cong (\text{Hom}_R(M_1/f^{-1}(N), M_1))$  , since  $M_1 \cong M_2$  . Thus  $\text{Hom}_R(M_2/N, M_2) = 0$  for all nonzero pure submodule  $N$  of  $M_2$  . Therefore  $M_2$  is purely quasi-Dedekind .

The proof of the converse is similarly .  $\square$

Remark 2.10. Let  $M$  be a purely quasi-Dedekind  $R$ -module and  $N \leq M$  then not necessary that  $M/N$  is a purely quasi-Dedekind  $R$ -module, as the following example shows .


Example 2.11. It is know that  $Z$  as  $Z$ -module is purely quasi-Dedekind, let  $N = 6Z \leq Z$  . But  $Z/6Z \cong Z_6$  is not a purely quasi-Dedekind as  $Z$ -module ( see Rem.and.Ex 2.3(2)) .

Now, we shall give a necessary condition under which the (Rem 2.10) is true .

Proposition 2.12. Let  $M$  be a purely quasi-Dedekind  $R$ -module with  $\frac{M}{K}$  is projective for all pure submodule  $K$  of  $M$ , then  $\frac{M}{N}$  is a purely quasi-Dedekind  $R$ -module for all  $N \leq M$  .



Proof. Let  $N \leq M$ . If  $N = 0$ , then nothing to prove. Now, let  $0 \neq N \leq M$ . Suppose that  $\frac{U}{N}$  is a pure submodule of  $\frac{M}{N}$ , then by ( Lemma 2.8)  $\pi^{-1}(\frac{U}{N})$  is a pure submodule of  $M$ , where  $\pi$  is the canonical projection map, so  $U$  is a pure submodule of  $M$ , hence  $\frac{M}{U}$  is projective by hypothesis. Assume that  $\frac{M}{N}$  is not purely quasi-Dedekind, thus there exists a nonzero  $R$ -homomorphism  $f : \frac{M/N}{U/N} \rightarrow \frac{M}{N}$ . But  $Hom_R(\frac{M/N}{U/N}, \frac{M}{N}) \cong Hom_R(\frac{M}{U}, \frac{M}{N})$ , so there exists  $R$ -homomorphism  $g : \frac{M}{U} \rightarrow M$  such that  $\pi \circ g = f$ .



$g \neq 0$  ( since  $f \neq 0$  ), thus  $Hom_R(\frac{M}{U}, M) \neq 0$ ,  $U$  is pure. Hence  $M$  is not a purely quasi-Dedekind  $R$ -module which is a contradiction. Therefore  $\frac{M}{N}$  must to be a purely quasi-Dedekind  $R$ -module.  $\square$

Remark 2.13. Let  $M$  be an  $R$ -module and  $N \leq M$ . If  $M/N$  is a purely quasi-Dedekind  $R$ -module then not necessary that  $M$  is a quasi-Dedekind  $R$ -module; For example: Consider  $Z$ -module  $Z_6$ ,  $N = (\bar{2}) \leq Z_6$ . Then  $Z_6/(\bar{2}) \cong Z_2$  is a purely quasi-Dedekind as  $Z$ -module, but  $Z_6$  is not a purely quasi-Dedekind as  $Z$ -module (see Rem.and.Ex 2.3 (1), (2)).

The following example shows the direct sum of purely quasi-Dedekind modules is not necessary that a purely quasi-Dedekind module.

Example 2.14. Each of  $Z_2, Z_3$  as  $Z$ -module is purely quasi-Dedekind (see Rem.and.Ex 2.3(1)), but  $Z_2 \oplus Z_3 \cong Z_6$  is not a purely quasi-Dedekind as  $Z$ -module.

Now, we gives a condition under which the direct sum of purely quasi-Dedekind modules is also purely quasi-Dedekind in the next proposition.

Proposition 2.15. Let  $M$  and  $N$  be a purely quasi-Dedekind  $R$ -modules with  $ann_R M + ann_R N = R$  then  $M \oplus N$  is a purely quasi-Dedekind  $R$ -module .

Proof. Assume that  $K$  is a pure submodule of  $M \oplus N$  . And since  $ann_R M + ann_R N = R$  then by same way of the proof of [1, Prop 4.2, Ch.1]  $K = K_1 \oplus K_2$ , where  $K_1 \leq M$  and  $K_2 \leq N$  .But

$K_1 \leq^{\oplus} K$  and  $K_2 \leq^{\oplus} K$  then by [21]  $K_1, K_2$  are pure in  $K$ , but  $K$  is pure in  $M \oplus N$  by hypothesis, then  $K_1$  is pure in  $M$  and  $K_2$  is pure in  $N$ ; to show this : Assume that there exists be an ideal  $I$  of  $R$  such that  $IK_1 \neq K_1 \cap IM$  and ( $IK_2 \neq K_2 \cap IN$  or  $IK_2 = K_2 \cap IN$  ) then

$IK = I(K_1 \oplus K_2) = IK_1 \oplus IK_2 \neq (K_1 \cap IM) \oplus (K_2 \cap IN) = (K_1 \oplus K_2) \cap I(M \oplus N)$   
 $= K \cap I(M \oplus N)$  which is a contradiction . So  $Hom_R(M/K_1, M) = 0$  and  $Hom_R(N/K_2, N) = 0$ , since  $M$  and  $N$  is purely quasi-Dedekind . On the other hand we have  $Hom_R(M \oplus N/K, M \oplus N) =$

$Hom_R(M \oplus N/K_1 \oplus K_2, M \oplus N) \subseteq Hom_R(M/K_1, M) \cap Hom_R(N/K_2, N) = 0$ . Hence  $M \oplus N$  is a purely quasi-Dedekind  $R$ -module .  $\square$

Recall that an  $R$ -module  $M$  is scalar if, for all  $f \in End_R(M)$  then there exists  $r \in R$  such that  $f(x) = rx$  for all  $x \in M$  [18, P.8] .

In the following proposition we shall study the endomorphism ring of purely quasi-Dedekind module .

Proposition 2.16. Let  $M$  be a scalar  $R$ -module with  $ann_R M$  is a prime ideal of  $R$ , then  $End_R(M)$  is a purely quasi-Dedekind ring .

Proof. Since  $M$  be a scalar  $R$ -module, then by [15, Lemma 6.2, P.80]  $End_R(M) \cong R/ann_R M$  ,

But  $ann_R M$  is a prime, so  $End_R(M)$  is an integral domain. Hence by (Rem.and.Ex 2.3(3))

$End_R(M)$  is a purely quasi-Dedekind ring .  $\square$

Corollary 2.17. If  $M$  is a scalar and prime  $R$ -module, then  $End_R(M)$  is a purely quasi-Dedekind ring .

Proof. It is clearly, since  $M$  is prime implies  $ann_R M$  is a prime ideal, so the result is obtained by ( Prop 2.16) .  $\square$

Proposition 2.18. Let  $M$  be a scalar faithful  $R$ -module .  $End_R(M)$  is a purely quasi-Dedekind ring if and only if  $R$  is a purely quasi-Dedekind ring .

Proof. Suppose that  $M$  is a scalar  $R$ -module, so  $End_R(M) \cong R/ann_RM$  by [15, Lemma 6.2, P.80], but  $M$  is a faithful, thus  $R/ann_RM \cong R$ , so  $End_R(M) \cong R$ . Hence we have on the result.  $\square$

Corollary 2.19. Let  $M$  be a finitely generated multiplication faithful  $R$ -module.  $End_R(M)$  is a purely quasi-Dedekind ring if and only if  $R$  is a purely quasi-Dedekind ring.

Proof. Since  $M$  is a finitely generated multiplication  $R$ -module, then by [16, The.3.2]  $M$  is scalar  $R$ -module; that is  $M$  is a scalar faithful  $R$ -module, thus by (Prop 2.18) the result is obtained.  $\square$

Recall that an  $R$ -module  $M$  is called quasi-prime if  $ann_R N$  is a prime ideal of  $R$  for each  $0 \neq N \leq M$  [2, def 1.2.1].

Proposition 2.20. Let  $M$  be a quasi-injective scalar and quasi-prime  $R$ -module then  $End_R(N)$  is a purely quasi-Dedekind ring for all  $0 \neq N \leq M$ .

Proof. Assume that  $0 \neq N \leq M$ . Since  $M$  is a quasi-injective scalar  $R$ -module, then by [18, Prop 1.1.16]  $N$  is a scalar  $R$ -module, thus  $End_R(N) \cong R/ann_R N$  by [15, Lemma 6.2, P.80]. But  $M$  is a quasi-prime  $R$ -module, so  $ann_R N$  is a prime ideal of  $R$ ; that is  $End_R(N) \cong R/ann_R N$  is an integral domain. Hence by (Rem.and.Ex 2.3(3))  $End_R(N)$  is a purely quasi-Dedekind ring.  $\square$

We end this section by the following two corollaries.

Corollary 2.21. If  $M$  is an injective scalar and quasi-prime  $R$ -module then  $End_R(N)$  is a purely quasi-Dedekind ring for all  $0 \neq N \leq M$ .

Proof. Obvious.  $\square$

Corollary 2.22. Let  $M$  be a quasi-injective scalar  $R$ -module and let  $0 \neq N \leq M$  be a faithful  $R$ -module. Then  $End_R(N)$  is a purely quasi-Dedekind ring if and only if  $R$  is a purely quasi-Dedekind ring.

Proof. Follows by [18, Prop 1.1.16] and (Prop 2.18).  $\square$

### 3. Purely Prime Modules

Recall that an  $R$ -module  $M$  is called prime if,  $ann_RM = ann_R N$  for all nonzero submodule  $N$  of  $M$  [8]. In this section we see that if  $M$  is purely quasi-Dedekind then  $ann_RM = ann_R N$  for all nonzero pure submodule  $N$  of  $M$  (Prop 3.2). This leads us to introduce many of important statement to this concept with other concepts in this section. We start this section with the following definition:

Definition 3.1. An  $R$ -module  $M$  is said to be purely prime if,  $ann_RM = ann_R N$  for all nonzero pure submodule  $N$  of  $M$ .

It is clear that every prime module is a purely prime module, but the converse need not be in general; for example :  $Z_4$  as  $Z$ -module is purely prime . In fact  $Z_4$  has no proper nonzero pure submodule as  $Z$ -module, but it is not prime as  $Z$ -module, since  $(\bar{2}) \leq Z_4$ ,  $ann_Z(\bar{2}) = 2Z \neq 4Z = ann_Z(Z_4)$ .

**Proposition 3.2.** Every purely quasi-Dedekind module is a purely prime module .

*Proof.* Follows by (Rem.and.Ex 2.3(7)) .  $\square$

**Proposition 3.3.** Let  $M$  be an  $R$ -module. Then  $M$  is a purely prime  $R$ -module if and only if  $M$  is a purely prime  $\bar{R}$ -module, where  $\bar{R} = R/ann_R M$  .

*Proof.*  $\Rightarrow$ ) Suppose that  $N$  is a nonzero pure  $\bar{R}$ -submodule of  $M$  . It is easy to see that  $N$  is a nonzero pure  $R$ -submodule of  $M$  . Let  $I$  be an ideal of  $\bar{R}$  , so it is also ideal of  $R$ , thus

$IN = N \cap IM$  hence  $N$  is a pure  $R$ -submodule of  $M$ , so that  $ann_R M = ann_R N$  . Now, it is clear that  $ann_{\bar{R}} M \subseteq ann_{\bar{R}} N$  , beside let  $r + ann_{\bar{R}} M \in ann_{\bar{R}} N$  then  $rN = 0$  ; that is  $r \in ann_R N = ann_R M$  ,

hence  $r + ann_{\bar{R}} M \in ann_{\bar{R}} M$  , therefore  $ann_{\bar{R}} M = ann_R N$  .

$\Leftarrow$ ) The proof is similarly .  $\square$

**Proposition 3.4.** Let  $M$  be a uniform regular  $R$ -module. Then the following statements are equivalent :

- 1)  $M$  is a prime  $R$ -module .
- 2)  $M$  is a purely prime  $R$ -module .
- 3)  $M$  is a purely quasi-Dedekind  $R$ -module .
- 4)  $M$  is a quasi-Dedekind  $R$ -module .

*Proof.*

(1)  $\Leftrightarrow$  (2): Clear .

(3)  $\Rightarrow$  (2): Follows by (Prop 3.2) .

(2)  $\Leftarrow$  (3): Suppose that  $M$  is purely prime, and since  $M$  is regular , so  $M$  is prime; that is  $M$  is prime uniform, thus by [14, The 3.11, P.37]  $M$  is quasi-Dedekind and hence  $M$  is purely quasi-Dedekind .

(3)  $\Leftrightarrow$  (4) : Follows by (Rem.and.Ex 2.3(5)) .  $\square$

Corollary 3.5. Let  $M$  be a multiplication uniform regular  $R$ -module. Then

(1)  $\Leftrightarrow$  (2)  $\Leftrightarrow$  (3)  $\Leftrightarrow$  (4)  $\Leftrightarrow$  (5)  $\Leftrightarrow$  (6)  $\Rightarrow$  (7)

- 1)  $M$  is a prime  $R$ -module .
- 2)  $M$  is a purely prime  $R$ -module .
- 3)  $M$  is a purely quasi-Dedekind  $R$ -module .
- 4)  $M$  is a quasi-Dedekind  $R$ -module .
- 5)  $End_R(M)$  is an integral domain .
- 6)  $End_R(M)$  is a quasi-Dedekind ring .
- 7)  $End_R(M)$  is a purely quasi-Dedekind ring .

**Proof.**

(1)  $\Leftrightarrow$  (2)  $\Leftrightarrow$  (3)  $\Leftrightarrow$  (4) : Follows by (Prop 3.4) .

(4)  $\Leftrightarrow$  (5) : Follows by [11, Prop 2.1.27] .

(5)  $\Leftrightarrow$  (6) : Follows by [11, Rem.and.Ex 1.1.2(7)]

(6)  $\Rightarrow$  (7) : Clear .  $\square$

Recall that an  $R$ -module  $M$  is monofrom if for each  $N \leq M$  and for each  $f \in Hom_R(N, M)$ ,  $f \neq 0$  implies  $Kerf = 0$  [22] .

Remark 3.6. Every monofrom module is a purely quasi-Dedekind module and hence it is a purely prime module .

The converse of above remark is not true in general; for example : Consider  $Z$ -module  $Z \oplus Z$  then it is known that is purely prime, since it is prime. But  $Z \oplus Z$  is not monofrom as  $Z$ -module.

Proposition 3.7. Let  $M$  be a uniform regular ring. Then the following statements are equivalent :

- 1)  $R$  is a monofrom ring .
- 2)  $R$  is an integral domain .
- 3)  $R$  is a quasi-Dedekind ring .

- 4)  $R$  is a purely quasi-Dedekind ring .  
 5)  $R$  is a purely prime ring .  
 6)  $R$  is a prime ring .

**Proof.**

(1)  $\Leftrightarrow$  (2)  $\Leftrightarrow$  (3) : Follows by [11, Coro 2.3.20] .

(3)  $\Leftrightarrow$  (4) : Clear .

(4)  $\Rightarrow$  (5) : Clear .

(5)  $\Rightarrow$  (4) : Assume that  $R$  is purely prime , and since  $R$  is regular, then  $R$  is prime. But  $R$  is uniform, so by [14, The 3.11, P.37]  $R$  is quasi-Dedekind, hence  $R$  is a purely quasi-Dedekind ring .

(5)  $\Leftrightarrow$  (6) : Clear .  $\square$

**Proposition 3.8.** Let  $M$  be an  $R$ -module. If  $M$  is embedded in each of its nonzero pure submodule then  $M$  is a purely prime  $R$ -module .

**Proof.** Suppose that  $N$  is a nonzero pure submodule of  $M$  . It is known that  $ann_R M \subseteq ann_R N$  .

On the other hand, let  $r \in ann_R N$  then  $rN = 0$ . But  $M$  is embedded in  $N$  (by hypothesis), so there exists a monomorphism  $f : M \longrightarrow N$  , thus  $f(rM) = rf(M) \subseteq rN = 0$  implies  $rM = 0$  (since  $f$  is monomorphism ), so  $r \in ann_R M$  and  $ann_R M = ann_R N$  . Hence  $M$  is a purely prime

$R$ -module .  $\square$

**Corollary 3.9.** Let  $M$  be a uniform regular  $R$ -module such that  $M$  is embedded in each of its nonzero pure submodule then  $M$  is a quasi-Dedekind  $R$ -module and hence it is a purely quasi-Dedekind  $R$ -module .

**Proof.** Follows by (Prop 3.8) and (Prop 3.4) .  $\square$

Recall that an  $R$ -module  $M$  is said to be weak cancellation if, for any two ideals  $A, B$  of  $R$  with  $AM = BM$  implies that  $A + ann_R M = B + ann_R M$  . And recall that an  $R$ -module  $M$  is cancellation if  $M$  is weak cancellation and faithful [6] .

Mijbass A.S. in [13, P.62 , P.63] introduce the following two results :

**Theorem 3.10.** Let  $M$  be an  $R$ -module and let  $N$  be a pure in  $M$  with  $ann_R N = ann_R M$  . If  $N$  is a weak cancellation  $R$ -module then  $M$  is a weak cancellation  $R$ -module .

**Corollary 3.11.** Let  $M$  be an  $R$ -module and let  $N$  be a pure in  $M$  with  $ann_R N = ann_R M$  . If  $N$  is a cancellation  $R$ -module then  $M$  is a cancellation  $R$ -module .

We end this section by the following two corollaries .

Corollary 3.12. Let  $M$  be a purely prime  $R$ -module and let  $N$  be a pure in  $M$ . If  $N$  is a weak cancellation  $R$ -module then  $M$  is a weak cancellation  $R$ -module .

Proof. Follows by (Th 3.10) .  $\square$

Corollary 3.13. Let  $M$  be a purely prime  $R$ -module and let  $N$  be a pure in  $M$ . If  $N$  is a cancellation  $R$ -module then  $M$  is a cancellation  $R$ -module .

Proof. Follows by (Coro 3.11) .  $\square$

## **References**

- [1] Abbas M.S. ,( 1991), On fully stable Modules , ph.D.Thesis , College of Science , University of Baghdad.
- [2] Abdul - Razak H. M ,( 1999), Quasi - Prime Modules and Quasi - Prime Submodules, M.Sc.Thesis, College of Education Ibn AL- Haitham ,University of Baghdad.
- [3] AL-Bahraany B.H. ,( 2000 ), Modules with the pure intersection property , Ph.D.Thesis , College of Science , University of Baghdad .
- [4] Ali M.M. and Smith D.J. ,( 2004), Pure submodules of Multiplication Modules, Contributions to Algebra and Geometry, Heldermann Verlag ,NO.1 , p.61-74 .
- [5] Anderson F.W. , Fuller K.R. ,( 1974), Rings and Categories of Modules , Springer -Verlag , Berlin , Heidelberg ,Newyork .
- [6] Atiyah M .F. , Macdonald I.G. ,( 1969), Introduction to commutative algebra, Addison Wesley, London .
- [7] Cheatham T.J. and Smith J.R. ,( 1976), Regular and Semisimple Modules, Pacific Journal Math ,NO.66 , p.315 - 323 .
- [8] Desale G. , Nicholson W.K. ,( 1981), Endoprimitive rings , J. Algebra ,NO.70 , p.548 - 560.
- [9] Faith C. ,( 1967), Lectures on injective Modules and quotient rings, Springer - Verlag , Berlin, Heidelberg , Newyork .
- [10] Fieldhouse D.J. ,( 1969), Pure theories, Math . Ann , NO.184, p.1-8 .
- [11] Ghawi Th.Y. ,( 2010), Some generalizations of Quasi - Dedekind Modules , M.Sc.Thesis ,College of Education Ibn AL- Haitham ,University of Baghdad .
- [12] Kasch F. ,( 1982), Modules and rings, Academic press , London .

- [13] Mijbass A .S.( 1992) , Cancellation Modules, MS.c.Thesis, College of Science, University of Baghdad .
- [14] Mijbass A .S.,( 1997), Quasi-Dedekind Modules, Ph.D.Thesis, College of Science, University of Baghdad.
- [15] Mohamed - Ali E. A . (2006) , On Ikeda - Nakayama Modules , Ph .D .Thesis, College of Education Ibn AL- Haitham , University of Baghdad .
- [16] Naoum A .G. ,( 1990), On the ring of endomorphisms of a finitely generated multiplication Modules, Periodica Math , Hungarica ,Vol .21(3), p. 249 - 255 .
- [17] Naoum A .G. ,Hadi I. M - A ., (2002), SQI Submodules and SQD Modules, Iraqi J.Sci , Vol.43.D, NO. 2 , P. 43 – 54 .
- [18] Shihab B . N. ,( 2004), Scalar Reflexive Modules, Ph .D.Thesis , College of Education Ibn AL-Haitham , University of Baghdad .
- [19] Smith P.F. , (1988), Some remarks on Multiplication Modules, Arch . Math, NO.50 , p. 223 – 235 .
- [20] Storrer H. H . , (1972), On Goldman's primary decomposition, Lecture notes in mathematics , Vol. 246, Springer-Verlag , Berlin, Heidelberg , Newyork .
- [21] Yaseen S.H. ,( 1985), F - Regular Modules, M.Sc.Thesis , University of Baghdad .
- [22] Zelmanowitz J. M . , (1986), Representation of rings with faithful polyform Modules , Comm. In Algebra, 14 (6) , p.1141 - 1169 .

## المقاسات شبه- ديديكاندية النقية و المقاسات الأولية النقية

ثامر يونس غاوي

العراق \ جامعة القادسية

كلية التربية - قسم الرياضيات

thaar\_math83@yahoo.com البريد الالكتروني :

### الخلاصة :-

يسمى المقاس الجزئي  $N$  من المقاس  $M$  على الحلقة  $R$  بالمقاس الجزئي النقي إذا كان  $IN = N \cap IM$  لكل مثالي  $I$  من الحلقة  $R$ . في بحثنا هذا قدمنا المقاس الجزئي شبه- معكوس النقي، حيث أن المقاس الجزئي  $N$  من المقاس  $M$  يسمى شبه- معكوس نقي إذا كان  $N$  مقاس جزئي نقي ومقاس جزئي شبه- معكوس أي  $Hom_R(M/N, M) = 0$ . يسمى المقاس  $M$  بأنه مقاس شبه- ديديكاندي نقي إذا كان كل مقاس جزئي غير صفري نقي  $N$  من  $M$  هو مقاس شبه- معكوس. من جانب آخر نحن أيضاً قدمنا مفهوم آخر من المقاسات يسمى المقاس الأولي النقي، حيث يسمى المقاس  $M$  على الحلقة  $R$  بأنه مقاس أولي نقي إذا كان  $M = N$  تالف  $N$  لكل مقاس جزئي غير صفري نقي  $N$  من  $M$ . لقد أعطينا العديد من الخواص الأساسية المتعلقة بهذه المفاهيم. كذلك درسنا العلاقات بين هذه المفاهيم وأنواع عديدة أخرى من المقاسات. في هذا البحث الحلقة  $R$  هي أبداً بمحايد و  $M$  مقاساً أحادياً على  $R$ .