

The effect of green tea and *Hibiscus sabdriffira* aqueous extract on some immunologic aspect in lab animals

Azhar – AL- Thahab*

الخلاصة

في مقارنة لدراسة تأثير المسد تخلص المائي لنبات الشاي الأخضر والكجرات في ض المعايير المناعية في الحيوان والمختبري جرع الحيوان والمختبري بكمية 5 مل من المسد تخلص المائي فموي لمدة أسبوعين، الاسد بوع الذامس تم قتل الحيوانات وتشريحها للحصول على اجزاء من الجهاز الهضمي للدراسة الموضوعية ولاجراء الاختبارات المناعية ومن النتائج اظهر مسد تخلص الشاي معنويًا أكثر مقارنة بمسد تخلص الكجرات في تحفيز الضد ألجهازية والموضوعية تحت مسد توى احتمالية $P \geq 0.005$ وان الكجرات زيادة في تركيز البروتين الكلي والمزاعي وحفزت النيلة مناعة خلوية متمثلاً باختبار تثبيط هجرة الخلايا البيضاء في الموضوع وغير تحفيزي في الجهازية .

Abstract

In this study compare between the effect of aqueous extract of green tea and *Hibiscus sabdroffira* on some immunological parameters in vivo . Five ml of watery extract of these was given in animal orally for three weeks after five weeks animals were killed and pieces of intestinal tract was collected to study local immunity .Green tea extract was shown higher significant at $P \geq 0.005$ in stimulation systemic and mucosal antibodies than *Hibiscus* extract . *Hibiscus sabdroffira* was shown increase in total protein and secretory immunoglobulins .Two plants were show effect on cellular immunity by increasing LIF mucosally but not systemically.

Introduction

Natural products play an important role in the field of new drugs researches and development . Since ancient times plants have been an exemplary source of medicine . In the last years there has been significant increase in the use of natural products in health care and there potential application in agriculture , pharmaceutical and food industry are being investigated .

*Babylon University- College of Science

Many literature mention the use of plants in treatment of various human aliment, so there are many plants which are having immunostimulatory whereas other have immunosuppressant activity (1).

Plants extracts are potentially curative , some of these extracts can boost the humoral and cell mediated immunity against bacteria, virus ,fungi , parasite (2) Some medicinal plants have been claimed to posses immunomodulatory activity (3). *Hibiscus sabdoffira* linn (Family Malvaceae) is a herb that is cultivated for leaf , fleshy calyx ,seed or fiber . It is taken as a common local drink . It is medicinal herb used in treatment hypertension , cardio protective , antiseptic , aphrodisiac , astringent , diuretic and digestive (4,5) . Green produced from the leaves of *Camellia sinensis* plant , it is one of the most widely consumed beverages in the world . It is contains several polyphenolic compounds , including EGCC – (-) – epigallocatechin (EGC) , epicatechin – 3-gallate (ECC) AND (-) - epicatechin EC -. Green tea has a beneficial effect on healthy , including a chemopreventive efficacy against various types of cancer (6,7) . In addition green tea has been shown to exert both antiangiogenic , antimutagen and anti-inflammatory (8) The aime of this study to compare between more popular two plants were used in many countries on some immunologic aspect in lab animals .

Materials and Methods

Animals

Rabbits were used as experimental animals . Healthy white Newzland rabbit (*Oryctolagus cuniculus*) about 1-15 kg. They hands at room temperature in labium condition during experimental condition

Tea extract

1gm of tea leaves and calyx *Hibiscus* with 10 ml of boiling tap water for 1hr at water both . Aqueous extract were then filtered and cooled at room temperature (9).

Immunization program

Animals were divided into 3 groups , 5 animals for each group. Each group were given 5 ml of plant extract orally and 5 ml of tap water as normal control daily for three weeks.

Blood samples

Five ml of blood was collected from each rabbits by using sterile disposable syringes from heart , 3 ml was put into AFMA disposable tubes without anticoagulant , then the serum was collected after centrifugation at 2500 rpm for 5 minutes and it was stored at -20C , other 2ml of blood was put in AFMA disposable to be with anticoagulant for LIF.

Mucosal extract . The groups of animal were killed and biopsies of appendix were opened by clean scissor and laid in clean Petri dishes and the mucosa were scraped with 10 ml formal saline and laid in clean tubes centrifuged at 3500 rpm / 30 minute pellet was used in LIF test , the supernatant was collected and equal volume of PEG 6% were added to supernatant and leave 30 minute to room temperature supernatant was remove and 1 ml of saline added to pellet to form mucosal immunoglobulin .

Immune function tests

Tube agglutination , total and immunoglobulin protein and LIF test was done as in (10,11,12).

Statistical analysis

All data obtained were subjected to student s' t-test analysis . Data were expressed as mean standard deviation .Values of $P \geq 0.005$ were considered significant (13) .

Results

Humoral

The rabbits were immunoprimered with plants extract showed specific mucosal and systemic antibodies . Systemic antibodies was higher in two extracts 640(3 times) . 320(one times) in green tea extract and 320(4 times) ,160(one times) in *Hibiscus* extract . Mucosal antibodies appeared 64(4 times) ,32(one times) in tea extract and 32(3 times) ,16(one times) in hibiscus extract , table (1) . Comparison between two plants extract was done by using T test ,

we found that variable in significant , tea extract higher significant than *Hibiscus* extract in p.05 ,table (2) . The effect of two plants extract on total and immunoglobulin protein by using Biurate method compare with control . *Hibiscus* extract showed higher significant and with tea extract is significant too, table(3)

Cellular

Two plant extract was shown to stimulate LIF test compare with control ,table (4).

Green tea was 0.42 while *H. sabdraffira* was 0.40 in mucosal but not stimulate in systemic .

Table(1): Specific mucosal and systemic antibodies in rabbits that administration tea and *Hibiscus* extract

Groups	systemic		mucosal	
	Titer	Frequency	Titer	Frequency
Green tea	640	4	64	4
	320	1	32	1
<i>H. sabdraffira</i>	320	4	32	4
	160	1	16	1
<i>control</i>	10	5	1	5

Table(2) : Titer antibodiess of green tea and *Hibiscus* extract in animals

Groups	Systemic		Mucosal	
	Mean ± Standar deviations	P ≥0.005	Mean ± Standard deviations	P ≥0.005
Green tea	560.000±16.000	0.006	56.000±16.000	0.006
<i>H.sabdraffira</i>	280.000±8.000	0.006	26.000±8.000	0.006

Table(3): Comparative of protein and immunoglobulin concentrations of tea and *Hibiscus* extract with control

Groups	Total protein mg/ml		Immunoglobulin concentrations mg/ml	
	Mean ± Standar deviations	P ≥0.005	Mean ± Standard deviations	P ≥0.005
Green tea	12.208 ± 5.999	0.02	2.91 ± 0.4849	0.001
<i>H.sabdraffira</i>	.234 ± 7.1712	0.001	4.4933 ± 17.927	0.000
<i>control</i>	30.78 ± 0.0447	0.000	3.677 ± 0.2543	0.000

Table(4 :Comparative in LIF between plants extract and control

Animals groups			
Systemic		Mucosal	
Mean±.standard deviations	P ≥0.005	Mean ± Standard deviations	P ≥0.005
Green tea 0.2250 ±0.01	0.003	0.425±0.04	0.003
<i>H. sabdroffira</i> 0.2250 ±0.01	0.003	0.4000±0.01	0.011
Control 0.9250 ± 0.01	0.000	0.8500 ± 0.01	0.000

Discussion

For each plants have been a good source of food and they provide essential nutritional values, medicinal properties and notable physiological effects to life. Plants extracts are potentially curative. Some of these extracts can boost the humoral and cell-mediated immunity against some kinds of microbes (2). In this study *Hibiscus* extract stimulated humoral response demonstrated in high titers and increased in protein and tea extract stimulated humoral response demonstrated by high titers but the protein is decreased.

Two plants extracts were effective on cellular response demonstrated by LIF there was no migration for T cells compared with control in systemic while in local was significant. Two plants contain some compounds that proved effective on immune system varies

between stimulator or suppressor . Delphinidin-3- sabu bioside (anthocyanin), flavons, glycosides, polysaccharides , protocatechuic acid , vitamin c and other substances , which are powerful antioxidant .The antioxidants prevent cells and tissue from oxidative damage , lipid peroxidation and also improve immunity response and productivity against stress ,including challenges of disease (14) . *Hibiscus* plant act as antiseptic ,sedative astringent. cholagogue,. Some polyphenolic compound ,flavonoids ,theaflavin ,catechins act as anti-inflammatory ,antimutagenic ,antiproliferative and antineoplastic activity (15). This study proposed two plants may be act as B dependent cells because they were stimulate higher antibodies titers and also possessed immunomodulating effect .

References

- 1- Jin-Ming,K.;Ngoh, G.;Lian.C;Tet,C.(2003).Resent advance in tradiotional plant drugs and orchids.Acta .Pharmacol.Sin.24:7-21.
- 2-Rehman,J.;Dillow,J.;Carter,S.;Cho,J.;Le.,B.Maisol,A.(1999).
Increased production of antigen specific immunoglobulin G and M fallowing in vitro treatment with medical plants .Immunol.Lett.68:391-396.
- 3- Hamant and Yaseen ,(2007) .Immunomodulatory Plants :Aphytopharmacological Review . Pharmacognosy Reviews .Vol.1,issue 2.
- 4- Okasha,M.; Abubakr ,M.; Bako,J.(2008) . Study the effect of aqueous *Hibiscus sabdroffira*Linn seed extract on serum prolactin level of lactating femal albinorats .European Journal of science research .ISSN 1450-216X Vol.4 .575-583
- 5- Wang ,C.;Wang ,J.and Lin,W.(200) .Protective effect of *Hibiscus* anthocyaninagainst ter-butyl hydroperoxidase induced hepatic toxicity in rats .Food.Chem.Toxicol.51:411-416
- 6- Shimizn,M.;Shirakami ,Y.; and Moriwaki ,H.(2008).Targeting reseptor tyrosinkinesis for chemoprevention by green tea catchin ,EGCC. Int.J.Mol.Sci .9:1034-1049.
- 7- Rodriguez,S.;Guno,W.;Liw,L.;Band,M.Paulson,E.;Meydani,M.(2006). Green tea catchin,epigallocatechin -3-gallate inhibits

- vascular endothelial growth factor angiogenic signaling by disrupting the formation of receptor complex. *Int.J.Cancer*.118:1635-1644
- 8- Han ,C.(1997). Screening of anticarcinogenic ingredients in tea polyphenol. *Cancer. Lett*.114:153-158
 - 9- Charles , O.; Michael , V.; Chukwuemeka, S.Damian,C.(2007). Adaptogenic potentials of *Camellia sinensis* leave, *Garcinia*
 - 10- Bishop, MC; Dben-con, J.; Fody Ep and et al (1985). *Clinical chemistry principles, procedure and correlation*, pp.189-182. The Murry printing company Philadelphia.
 - 11- Garvey, JS.; Cremev, NE. and Sussdrof, DH. (1977). *Method in immunology* 3th ed. PP53-267. Addison-Wesley publishing company Inc., Reading
 - 12- Soborg,M, (1969). In vitro migration inhibition of peripheral blood leucocytes in delayed type hypersensitivity. *Acta . Medica. Scand* 184:13-25.
 - 13- Dawed , K. M. and Al- Yas, ZA, (1990). *Statistical method in agricultural researches*. PP. 430-446. Al- Mous. Univ. (In Arabic)
 - 14- Minka, N.S ;Fayomi ,A.and Ayo.J.O.(2007) .Protective influence of calyces of *Hibiscus sabdariffa* against heat stress in laying hense during the hot –dry season. *Research Journal of Poultry Sciences* 1(1):7-11.
 - 15- Michiyo,T.;kara,I.; Zi-Jian,x.,andThoma,J.(2002) .Tea pigments inhibit the production of type 1 (TH1) and type 2 (TH2) helper T cell cytokine in CD4 T cell. *Phytotherapy Research*. 16,36-421