

Ministernotomy in Off-Pump Coronary Artery Bypass Surgery

Jabbar Jasim AL-Tai

ABSTRACT:

BACKGROUND:

Ministernotomy is a new surgical procedure which is decrease invasiveness of surgical treatment coronary arteries insufficiency and there are several advantages over standard procedure.

OBJECTIVE:

Patients with isolated lesions of the proximal left anterior descending artery (LAD) have been demonstrated to benefit more from surgical treatment than percutaneous coronary interventions (PCI). However, with the less invasiveness of PCI, the majority of the patients have been referred for this latter procedure. We report here on the inferior ministernotomy approach for the treatment of patients with single LAD lesions, with off-pump anastomosis of the left internal thoracic artery graft.

METHOD:

Fourteen patients, consecutively operated on using this technique with the LITA graft anastomosed to the LAD, were examined. The mean age of the patients was 56.7 ± 10.1 years. The length of the skin incision varied from 8 to 10 cm and only the distal sternum was split lengthwise. The anastomosis was facilitated with the use of an Octopus-3 stabilizer.

RESULTS:

All patients had satisfactory postoperative outcomes, the length of postoperative hospital stay ranged from 2 to 6 days (median 3 days). No ECG changes or enzymatic rises were seen in this series. One patient was re-admitted for wound infection.

CONCLUSION:

The ministernotomy approach allows safe accomplishment of off-pump LAD grafting, providing the long-term benefits of using the LITA.

KEY WORDS: ministernotomy, left internal thoracic artery (LITA), off-pump.

INTRODUCTION:

Recent studies have demonstrated that patients with isolated lesions of the proximal left anterior descending artery (LAD) achieve greater clinical benefits when submitted to surgical treatment using the left internal thoracic artery (LITA) ⁽¹⁻³⁾. Observational data, over the last decade, demonstrated that there is a lower incidence of cardiovascular events and greater survival when the LITA is utilized in coronary artery bypass grafting. However, with the reduced invasiveness of the percutaneous coronary intervention (PCI), the majority of patients with isolated injuries of the LAD have been referred for this type of procedure. The utilization of smaller operative incisions and doing without cardiopulmonary bypasses (CPB) might minimize the invasiveness and the surgical

aggression with real benefits for the patient.

The objective of this work is to report on the utilization of ministernotomy as a technical approach for the treatment of patients with single injuries to the LAD, using LITA grafts without the use of CPB.

AIM OF STUDY:

In this paper we are evaluating the technique of ministernotomy with Off-Pump coronary bypass surgery using the LITA graft, in the prognosis of the recovery.

PATIENTS AND METHODS:

In this series 14 consecutive patients had been operated in Ibn-alnafees for cardiovascular diseases hospital, from October 2009-April 2010, with single injuries of the LAD confirmed by coronary

Department of Cardiac Surgery, Ibn AL-Nafees Hospital for Cardiothoracic Diseases.

angiography were studied. All patients were submitted to coronary artery bypass grafting (CABG) using ministernotomies. LITA grafts were anastomosed to the LAD in all patients. Nine patients were male and five were female with ages ranging from 35 to 69 years old with a mean of 56.7 ± 10.1 years. The associated comorbidities were: prior myocardial infarction in eight patients (57.1%), diabetes mellitus in four (28.5%), peripheral arteries disease in one (7.1%), previous angioplasty of the right coronary artery with the implantation of a stent in one (7.1%) and

chronic obstructive pulmonary disease in one (7.1%) patient.

The ejection fraction measured by echocardiography, varied from 47% to 75% (mean $59.8\% \pm 9.6$). All the patients continue to be clinically followed up.

Technique

In ministernotomies, the cutaneous incision starts at the xiphoid appendix and extends for 7 to 10 cm in the cephalic direction (figure 1). The xiphoid appendix and the lower portion of the sternum are opened lengthways using an electric saw without additional bone incisions to the side (figure 2).



Figure 1 skin incision

The LITA is dissected from the thoracic wall using electric cautery close to its junction with the left subclavian artery, employing the skeletonization technique. Anticoagulation is achieved using a single dose of 400 IU/kg of heparin (figure 3). Subsequently, the pericardium is partially opened lengthwise and the heart is moved medially, using small compresses on the left ventricle lateral wall to expose the LAD. A tourniquet of 4-0

polypropylene thread is placed proximal to the site of the coronary arteriotomy. Anastomosis of the LITA with the anterior descending artery is achieved with a continuous suture employing a single 7-0 polypropylene thread without using CPB (figure 4).

The anastomosis is facilitated by using Octopus suction stabilizers. Figure 5; shows the cutaneous incision in a patient in the immediate postoperative period.



Figure 2:Ministernotomy.



Figure 3:LIMAharvasting.



Figure 4:LIMA to LAD anastamosi.



Figure 5:Incision immediate postoperative period.

RESULTS:

All the patients had a good evolution and were released from hospital after between 2 to 6 days of the postoperative period (median 3 days). No changes in the ECG or enzymatic increases were seen in this group. One patient was re-hospitalized due to infection of the surgical wound.

In the inter-operative period, all the patients tolerated the maneuvers of heart luxation to perform the anastomoses well. No patients required inotropic support or the use of vasoconstrictors

during or after the procedure. Additionally, there were no changes in the ECG or increases in the enzymatic during the postoperative period and no patients required mediastinal re-exploration for bleeding.

The length of follow up was 11 to 34 months (mean 22.6 months). Twelve patients are asymptomatic, one patient died from a stroke after 11 months of follow-up and one patient had a re-incidence of angina. For this latter patient, a control coronary

angiography showed severe stenosis of the middle segment of the LITA graft, probably due to a lesion caused during dissection.

DISCUSSION:

Cardiovascular surgeons have been trying hard to find new less-invasive techniques with less organic aggression over the last decade[9]. In particular, two aspects have been prioritized in this task: smaller incisions to reduce the surgical aggression and to avoid CPB(cardiopulmonary bypass) to diminish the systemic inflammatory response and risk of neurological events⁽¹⁰⁾.

CABG currently presents with low morbidity and mortality rates and provides late results that are safe and reliable; however, the surgical aggression caused by sternotomy and the use of CPB increase the necessity of longer hospital stays. Elimination of the CPB and smaller incisions may provide a quicker recovery with earlier hospital release giving a proportional reduction in hospital costs⁽¹¹⁾. Boylan MJ and colleagues published the same results⁽⁴⁾.

Grafting of the LAD by off-pump ministernotomy might still be a more effective alternative in cases of total occlusion, distal stenosis, small caliber vessels, selected cases of complex coronary trunk lesions where the results of PCI are still more restrictive than the studies of selected cases⁽⁵⁾.

The inferior ministernotomy approach offers some advantages, such as an easy access to the LAD and the possibility of dissection of the LITA under direct vision. It also allows planning of the extension of the utilization of this technique, as dissection of the right internal thoracic and gastroepiploic arteries for use as pedicled artery grafts is also possible. Additionally, access to the posterior descending artery is facilitated by this incision⁽¹²⁾.

Inferior ministernotomy has been used as 'J' or 'L' techniques⁽⁶⁾. These techniques use a second sternal incision to the left, generally in the 3rd intercostal space, which can increase the risk of injury to the LITA by the retractor and sternal pseudo-arthritis as in Calafiore AM, and colleagues study⁽⁷⁾.

The difficulties with the anastomoses of the LITA with the LAD through the left anterior ministernotomy encouraged us to develop the inferior ministernotomy. This latter technique presents advantages when compared to the left anterior ministernotomy, which was the first approach developed for minimally invasive CABG surgeries. The advantages include greater comfort

for the surgeon who works with a more familiar and routine incision, even though it is smaller; if there are complications the incision can be rapidly extended and converted to total sternotomy. Additionally, through this incision it is possible to have access to both internal thoracic arteries and also to the right gastroepiploic artery, which may be utilized in pedicled arterial grafts (although, without the necessity of a proximal anastomosis on the aorta) for revascularization of the three coronary arterial systems. For this, there is clearly a necessity to improve and develop new devices that allow greater access to the heart.

Skeletonization of the LITA is important to perform this technique, as it enables the attainment of longer pedicles which reach the site of the anastomosis with the LAD more comfortably and without tension. Additionally, skeletonization preserves the arterial vascular connections of the sternum, circumventing devascularization and the consequent risk of sternal necrosis⁽⁸⁾.

To avoid distal injury of the coronary artery, only proximal clamping is used. Another important aspect related to the quality of the surgery is the utilization of suction stabilizers, which immobilize the site of the coronary artery and allow better anastomosis techniques to be used. For this series, an Octopus 3® stabilizer (Medtronic, Inc) was used⁽¹³⁾.

With the incorporation of new devices that allow better exposure (both for the dissection of the internal thoracic arteries and the coronary arteries) better results and expand the use of the technique for multiple artery revascularization utilizing pedicled grafts only may be possible. In the only patient in this series who had recurrence of angina, the angiography demonstrated significant stenosis of the LITA, probably due to injury during dissection. An improved access to the operative field may contribute to avoid this complication⁽¹⁴⁾.

The advantage related to the less invasiveness of this technique is the possibility of earlier hospital release, with reduction in the hospitalization and a consequent reduction in costs. Additionally, there is a reduction in the necessity of blood transfusion; none of the patients in this series required transfusions. Biancari Fand colleagues demonstrate the same results⁽¹⁵⁾.

This technique combines the advantages of limited sternal incisions with the benefits of off-pump revascularization.

Thus, from prospective randomized studies of patients with injuries of the LAD in isolation,

where the superiority of the results with surgical treatment using LITA grafts is well known, it is possible to affirm that minimally invasive CABG surgery is the first-line choice giving safe and lasting results over the long term.

The length of follow up was 11 to 34 months (mean 22.6 months). Twelve patients are asymptomatic, one patient died from a stroke after 11 months of follow-up and one patient had a re-incidence of angina. For this latter patient, a control coronary angiography showed severe stenosis of the middle segment of the LITA graft, probably due to a lesion caused during dissection.

CONCLUSION:

Ministernotomy allows safe off-pump surgical CABG of the LAD with the long-term benefits of using the LITA.

REFERENCES:

1. Hueb WA, Soares PR, Oliveira SA, Arie S, Cardoso RH, Wajsbrot DB et al. Five-year follow-up of the medicine, angioplasty, or surgery study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty, or bypass surgery for single proximal left anterior descending coronary artery stenosis. *Circulation*. 1999;100:II107-13.
2. Goy JJ, Eeckhout E, Moret C, Burnand B, Vogt P, Stauffer JC et al. Five-year outcome in 2004;25:567-71. patients with isolated proximal left anterior descending coronary artery stenosis treated by angioplasty or left internal mammary artery grafting: as prospective trial. *Circulation*. 1999;99:3255-59.
3. Drenth DJ, Veeger NJ, Grandjean JG, Mariani MA, van Boven AJ, Boonstra PW. Isolated high-grade lesion of the proximal LAD: a stent or off-pump LIMA? *Eur J Cardiothorac Surg*.
4. Boylan MJ, Lytle BW, Loop FD, Taylor PC, Borsh JA, Goormastic M, Cosgrove DM. Surgical treatment of isolated left anterior descending coronary stenosis: comparison of left internal mammary artery and venous autograft at 18 to 20 years of followup. *J Thorac Cardiovasc Surg*. 1994;107:657-62.
5. Niinami H, Takeuchi Y, Ichikawa S, Suda Y. Partial median sternotomy as a minimal access for off-pump coronary artery bypass grafting: feasibility of the lower-end sternal splitting approach. *Ann Thorac Surg*. 2001;72:S1041-45.
6. Lichtenberg A, Klima U, Harringer W, Kim PY, Haverich A. Mini-sternotomy for off-pump coronary artery bypass grafting. *Ann Thorac Surg*. 2000;69:1276-77
7. Calafiore AM, Teodori G, DiGiammarco G, Vitolla G, Contini M. Minimally invasive coronary artery surgery: the last operation. *Semin Thorac Cardiovasc Surg*. 1997;9:305-11.
8. Henriquez-Pino JA, Gomes WJ, Prates JC, Buffolo E. Estudo anatômico da artéria torácica interna aplicado à cirurgia cardiovascular. *Rev Bras Cir Cardiovasc*. 1997;12:83-88.
9. Hirata N, Kanaya N, Yamazaki Y, et al. Pulmonary embolism caused by a carbon dioxide blower during off-pump coronary artery bypass grafting. *J Anesth* 2010;24:114-16.
10. Magee MJ, Hebert E, Herbert MA, et al. Fewer grafts performed in off-pump bypass surgery: Patient selection or incomplete revascularization? *Ann Thorac Surg* 2009;87:1113-18.
11. Halbersma WB, Arrigoni SC, Mecozzi G, et al. Four-year outcome of OPCAB no-touch with total arterial Y-graft: Making the best treatment a daily practice. *Ann Thorac Surg* 2009;88:796-801.
12. Valletly MP, Potger K, McMillan D, et al. Anaortic techniques reduce neurological morbidity after off-pump coronary artery bypass surgery. *Heart Lung Circ* 2008;17:299-304.
13. Joanna Chikwe, David Tom Cooke, Aaron Weiss. *Cardiothoracic Surgery*. 2nd Ed. United Kingdom: OXFORD university press, 2013; pp 304-5.
14. Larry R. Kaiser, MD, Irving L. Kron, MD, Thomas L. Spray, MD. *Mastery of Cardiothoracic Surgery*. 3rd Ed. USA: Wolters Kluwer Lippincott Williams & Wilkins, 2014; pp 508-9.
15. Biancari F, Mahar MA, Mosorin M, et al. Immediate and intermediate outcome after off-pump and on-pump coronary artery bypass surgery in patients with unstable angina pectoris. *Ann Thorac Surg* 2008;86:1147-52.

