

# SUBCLINICAL LEFT VENTRICULAR DIASTOLIC DYSFUNCTION IN HYPERTENSIVE PATIENTS IN SLEMANI CITY USING 2D ECHOCARDIOGRAPHY

Omed Hamakarem <sup>a</sup>, Kamaran Hassan <sup>b</sup> and  
Dana Mohammed Saeed <sup>a</sup>



Submitted: 28/5/2018; Accepted: 12/8/2018; Published 15/10/2018

## ABSTRACT

### *Background*

Diastolic dysfunction refers to when the diastole part is abnormal. The ventricles do not properly relax and become stiff meaning they cannot fill with blood properly. This causes blood to “dam up” in other parts of the body. The pressure in the ventricles then increases as blood from the next heartbeat tries to enter. Diastolic dysfunction is a common problem, with many people aged older than 70 years having the condition

### *Objectives*

To detect subclinical left ventricular diastolic dysfunction in hypertensive patients with preserved ejection fraction using 2D echocardiography.

### *Methods*

This is a cross-sectional study that was carried on 80 hypertensive patients referred to slemani cardiac hospital and medical emergency outpatient clinic for evaluation and treatment of hypertension and 80 age and sex-matched healthy volunteers as a control group. All subjects underwent conventional echocardiographic examination and assessment of diastolic dysfunction by 2D echocardiography according to updated European guidelines for diastolic dysfunction

### *Results*

A total of one hundred sixty (160) individuals were included in the final analysis. Mean age was  $55.03 \pm 7.67$ SD years. Males accounted for 51.9% (n=83) of the studied sample and 48.1% (n=77) were females. Fifty percent (n=80) were hypertensive (cases) and the other 50% (n=80) were non-hypertensive (controls).hypertensive patients have significantly higher rates of diastolic dysfunction (22 cases vs 12) and LVH (19 vs 5) than non-hypertensives, they also have higher mean lateral annulus velocity ( $8.83 \pm 2.41$  vs  $10.21 \pm 2.62$  ) and TR velocity ( $2.28 \pm 0.37$  vs  $2.11 \pm 0.11$ ) but smaller mean LA volume.

### *Conclusion*

Impairment of diastolic function detected by 2D echocardiography was significantly higher in hypertensive patients (with and without LVH) compared to nonhypertensive patients. Early detection of diastolic dysfunction in hypertensive patients is necessary to prevent diastolic heart failure which is the major type of heart failure in hypertensive patients.

**Keywords:** *Hypertension, 2D echocardiography, Diastolic dysfunction, Heart failure.*

---

<sup>a</sup> Department of Medicine, College of Medicine, University of Sulaimani.

Correspondence: [omid.karim@univsul.edu.iq](mailto:omid.karim@univsul.edu.iq)

<sup>b</sup> Kurdistan Board of Medical Specialties Candidate.

## INTRODUCTION

The global prevalence of hypertension defined as an average systolic BP of 140 mmHg or greater, a diastolic BP of 90mmHg or greater, or the use of antihypertensive medication was estimated to increase by 60% to a total of 1.56 billion by 2025, which is 29% of the worldwide adult population <sup>(1)</sup>. During the same period, 75% of the world's hypertensive population will be residing in the third world countries<sup>(1, 2)</sup>. According to the Framingham Study, hypertension accounts for about a quarter of heart failure cases<sup>(3)</sup>.

Diastolic heart failure (HF) is a progressive disorder characterized by impaired left ventricular (LV) relaxation, increased LV stiffness, increased interstitial deposition of collagen, and modified extracellular matrix proteins. Diastolic HF, also referred to as HF with normal ejection fraction currently accounts for 40% to 50% of all HF cases and has a prognosis, which is as ominous as that of systolic HF <sup>(4)</sup>. With life expectancy increasing, HF is growing into a major health problem. Because the process of myocardial remodeling starts before the onset of symptoms, recent HF guidelines <sup>(5)</sup>.

Doppler echocardiography is the most practical method for assessing filling patterns and myocardial relaxation and for estimating LV filling pressures at rest and with exertion by recording flow velocities from the atrioventricular valves, central veins, and myocardial tissue <sup>(6, 7, 8)</sup>. Diastolic dysfunction and diastolic heart failure are not synonymous terms<sup>(9)</sup>.

In the last 10 years or so, it has become apparent that approximately half of the patients who present with classic signs and symptoms of heart failure appear to have a normal ventricular function, typically defined by the finding of an ejection fraction of >50% on echocardiography. This group has been variously described as having diastolic dysfunction, diastolic heart failure, or "heart failure with a normal ejection fraction (HFNEF)<sup>(9)</sup>.

A major reason for the paucity of randomized controlled trials in heart failure patients is the difficulty in defining and measuring diastolic function. Although hemodynamic data obtained by heart catheterization can be used to measure diastolic function, the invasive nature of this assessment limits its applicability to most patients. Therefore, Doppler echocardiography is the method of choice in routine clinical practice to assess for diastolic dysfunction<sup>(10)</sup>.

LV diastolic dysfunction is a progressive condition and is characterized by an increased resistance dependence on ventricular preload <sup>(11)</sup>. Diastolic HF is thought to be responsible for as many as 74% of cases of HF in hypertensive patients<sup>(12)</sup>. Echocardiography is an alternative technique to cardiac catheterization in the evaluation of patients with diastolic dysfunction which include M-mode, 2-D, and Doppler echocardiography studies <sup>(13)</sup>.

The aim of the study is to detect the prevalence of subclinical left ventricular diastolic dysfunction in hypertensive patients with preserved ejection fraction using 2D echocardiography.

## PATIENTS AND METHODS

This is a cross-sectional study that was carried from January 2017 to January 2018 on 80 hypertensive patients referred to Slemani cardiac hospital and medical emergency outpatient clinic for evaluation and treatment of hypertension and 80 age and sex-matched healthy volunteers as a control group. They were enrolled in the study after obtaining their written informed consent. Our cases were all known cases of hypertension. Control subjects had no detectable cardiovascular risk factors and not receiving any medications.

Exclusion criteria included patients with ejection fraction <50% or sign and symptoms of heart failure, known case of diabetes mellitus, valvular lesions, atrial fibrillation, any rhythm disturbance, known case of thyrotoxicosis, COPD, morbid obesity, age>80, <40 and ischemic heart disease.

The following data were collected:

- Complete and detailed medical history: With attention to Hypertension, thyrotoxicosis, COPD, DM and family history of premature coronary artery disease.
- Full clinical examination including body surface area <sup>(14)</sup>, heart rate, rhythm, heart, and chest auscultation, systolic and diastolic blood pressure. Hypertension was classified based on JACC (Journal of the American college of cardiology) guidelines for classification of hypertension<sup>(15)</sup>.
- Assessment of LV diastolic functions with conventional echocardiography.

For diastolic function assessment, Echocardiography

was done by a single operator using the GE Vivid 7 echocardiography machine. Four variables were evaluated when determining whether LV diastolic function is normal or abnormal <sup>(16)</sup>. LV diastolic function is normal if more than half of the available variables do not meet the cutoff values for identifying abnormal function. LV diastolic dysfunction is present if more than half of the available parameters meet these cutoff values. The study is inconclusive if half of the parameters do not meet the cutoff values <sup>(16)</sup>. The presence of several abnormal findings, as well as cutoff values with high specificity for the myocardial disease, is recommended to decrease false positive diagnoses of diastolic dysfunction.

The four recommended variables and their abnormal cutoff values are annular e0 velocity (septal  $e0 < 7$  cm/sec, lateral  $e0 < 10$  cm/sec), average E/e0 ratio  $> 14$ , LA maximum volume index  $> 34$  mL/m<sup>2</sup>, and peak TR velocity  $> 2.8$  m/sec. On the basis of the writing group's collective expert opinion, average E/e0 ratio is recommended for simplification. Although E/e0 ratio may be obtained at septal or lateral annulus, and different values exist because of the normally higher lateral annular velocities, an average E/e0 ratio  $> 14$  is used throughout this document and is consistent with recent studies in normal subjects <sup>(16)</sup>.

Data analysis was done by computerized statistical software; Statistical Package for Social Sciences (SPSS) version 24. Descriptive statistics presented as (mean  $\pm$  standard deviation) and frequencies as percentages. Multiple contingency tables conducted and appropriate statistical tests performed, Chi-square used for categorical variables and independent t-test was used to compare between means. In all statistical analysis level of significance (p-value) set at  $\leq 0.05$  and the result presented as tables and/or graphs.

## RESULTS

A total of one hundred sixty (160) individuals were included in the final analysis. Mean age was  $55.03 \pm 7.67$ SD years. Males were accounted for 51.9% (n=83) of studied sample and 48.1% (n=77) were females. Fifty percent (n=80) were hypertensive (cases) and the other 50% (n=80) were non-hypertensive (controls). These values along with other socio-demographic features are shown in table 1

Both cases and controls are relatively well-matched in terms of numbers, ages (the hypertensive group is older by a mean of 3.59 years) and gender as shown in table 2 below.

Regarding hemodynamic measures of both groups, hypertensive patients have significantly higher rates of diastolic dysfunction (22 cases vs 12) and LVH (19 vs 5) than non-hypertensives, they also have higher mean lateral annulus velocity ( $8.83 \pm 2.41$  vs  $10.21 \pm 2.62$ ) and TR velocity ( $2.28 \pm 0.37$  vs  $2.11 \pm 0.11$ ) but smaller mean LA volume as shown in table 3.

The rate of Diastolic dysfunction and its parameters are significantly related to the grades of HT as illustrated in table 4.

Again hemodynamic measures and hence diastolic dysfunction are significantly related to the BMI as shown in table 5.

Regarding the relation of age to E/E' ratio in both groups, increasing age causes increase in E/E' ratio, (Moderate Positive correlation) and P values are significant (less than 0.05), interestingly, correlation coefficient (R) in non-hypertensive patients is greater than correlation coefficient in hypertensive patients, this means that age has lesser effect on E/E' when a patient has hypertension and vice-versa!. See figure 1 and 2.

**Table 1. Socio-demographic and clinical characteristics of the participants.**

<b>Variable</b>	<b>N (%)</b>
<b>Age (mean <math>\pm</math> SD) years</b>	55.03 $\pm$ 7.67
<b>Sex</b>	
Female	77 (48.1)
Male	83 (51.9)
<b>BMI</b>	
18.50 – 24.90	103 (64.4)
25.0 – 29.90	51 (31.9)
30.0 – 34.90	6 (3.8)
<b>Smoker</b>	
No	111 (69.4)
Yes	49 (30.6)
<b>Alcoholic</b>	
No	148 (92.5)
Yes	12 (7.5)
<b>Hypertension</b>	
No	80 (50)
Yes	80 (50)

**Table 2. Basic characteristics of hypertensive and non-hypertensive patients with comparison**

<b>Variable</b>	<b>HTN N</b>	<b>Non-HTN N</b>	<b>P Value</b>
<b>Age (mean <math>\pm</math> SD)</b>	56.82 $\pm$ 8.88	53.23 $\pm$ 5.72	0.003
<b>Sex</b>			
Female	41	36	0.429
Male	39	44	
<b>BMI</b>			
18.50 – 24.90	57	46	0.022
25.0 – 29.90	23	28	
30.0 – 34.90	0	6	
<b>Smoke</b>			
No	56	55	0.864
Yes	24	25	
<b>Alcoholic</b>			
No	74	74	1.000
Yes	6	6	

Table 3. Comparison of cardiac hemodynamic measures in both groups.

Variable	HTN N	NO HTN N	P Value
<b>Diastolic dysfunction</b>			
No	58	68	0.053
yes	22	12	
<b>Left ventricular hypertrophy</b>			
No	61	75	0.002
Yes	19	5	
<b>E/E' (mean ± SD)</b>	11.05±3.13	11.07±2.39	0.959
<b>LA volume(mean ± SD)</b>	29.16±5.27	31.41±1.65	0.000
<b>E' lateral (mean ± SD)</b>	8.83±2.41	10.21±2.62	0.001
<b>E'septal (mean ± SD)</b>	7.71±2.81	7.92±1.23	0.533
<b>TR Velocity (mean ± SD)</b>	2.28±0.37	2.11±0.11	0.0001

Table 4. Cardiac hemodynamic measures according to the grades of hypertension.

HT Grades	Diastolic dysfunction N No/Yes	LVH N No/Yes	E/E' (mean ± SD)	E' Lateral (mean ± SD)	E' septal (mean ± SD)	LA volume (mean ± SD)	TR Velocity (mean ± SD)	P-value
<b>Grade 0</b>	9/0	9/0	9±1.93	11.44±0.52	8.88±2.31	28.38±3.02	2.13±0.17	0.000
<b>Grade 1</b>	39/10	41/8	10.64±2.54	8.59±2.28	7.52±2.66	28.22±5.22	2.22±0.35	
<b>Grade 2</b>	10/12	11/11	11.17±3.84	9.35±2.61	8.64±3.41	28.82±4.72	2.39±0.51	

Table 5. Cardiac hemodynamic measures according to Body mass index.

BMI	Diastolic dysfunction No/Yes	LVH* No/Yes	E/E' (mean ± SD)	E' Lateral (mean ± SD)	E' septal (mean ± SD)	LA volume (mean ± SD)	TR Velocity (mean ± SD)	P-value
<b>18.5–24.9</b>	90/13	91/12	10.67±2.44	9.57±2.27	7.9±2.14	29.13±4.07	2.18±0.27	0.000
<b>25–29.9</b>	36/15	39/12	11.38±3.17	10.07±2.65	7.86±2.27	32.06±3.12	2.21±0.32	
<b>30–35</b>	0/6	6/0	14.96±0.25	4.0±0.0	6.0±0.0	35.1±0.0	2.3±0.0	

\*LVH: left ventricular hypertrophy




Figure 1. Correlation between E/E' with age in hypertensive patients.




Figure 2. Correlation between E/E' with age in non-hypertensive patients.

## DISCUSSION

Abnormalities of LV diastolic function are common in cases of systemic hypertension in whom the left ventricular systolic performance is still normal.

In this study hypertensive patients have significantly higher rates of diastolic dysfunction (22 cases vs 12) and LVH (19 vs 5) than non-hypertensives, they also have higher mean lateral annulus velocity ( $8.83 \pm 2.41$  vs  $10.21 \pm 2.62$ ) and TR velocity ( $2.28 \pm 0.37$  vs  $2.11 \pm 0.11$ ) but smaller mean LA volume. The rate of Diastolic dysfunction and its parameters are significantly related to the grades of HT.

The prevalence of LV diastolic dysfunction in hypertensives in this study was 27.5% (22 out of the 80). However, the prevalence in the normotensive controls was 15% (12 out of 80). A recent study in Asia on the prevalence of DD by Masliza et al <sup>(17)</sup> found a lower incidence of 18.6% (32 out of 198) in their newly diagnosed with systemic hypertensives and 6 (3.4%) among the controls. another study Kingue et al <sup>(18)</sup> in Black African untreated hypertensive population in Cameroon (67%). Balogun et al <sup>(19)</sup> reported a prevalence of 46% in their study on hypertensives. The selection, inclusion criteria and the characteristics of those patients might have accounted for the higher prevalence in their study. The reported prevalence



of LVDD depends on several factors, such as the characteristics of the population under study, imaging techniques, and the criteria applied to diagnose or to grade LVDD.

Relationships between LV diastolic function and overweight and obesity as classified by the WHO expert consultation definition for Asians were analyzed. This study suggests that BMI has an independent correlation with parameters of LV diastolic function. In our study population, the overweight and obese groups had more impaired diastolic function compared with the normal-weight group, and both of these weight groups were independent predictors of LV diastolic dysfunction.

Many studies have demonstrated that overweight and obesity are important risk factors for the development of heart failure <sup>(20-23)</sup>. Most studies use the most popular WHO BMI criteria of  $\geq 25$  kg/m<sup>2</sup> for overweight and  $\geq 30$  kg/m<sup>2</sup> for obesity <sup>(24)</sup>, which were both calculated based on Western populations. However, Asian populations have a lower mean BMI than Western populations, in addition to a higher percentage of body fat and a greater risk for cardiovascular disease at a given BMI <sup>(25, 26)</sup>.

In the I-PRESERVE echocardiographic substudy, LV mass and LA size remained independently associated with an increased risk of morbidity and mortality <sup>(27)</sup>. In other study LV hypertrophy, septal E/e' ratio, and TR peak velocity were predictive of outcome beyond clinical and laboratory characteristics. Because LV hypertrophy, elevated filling pressures, and elevated PASP frequently coexist, a greater number of these abnormalities is associated with a higher risk for incident hospitalization for heart failure <sup>(28)</sup>. LVDD develops early in most cardiac diseases and leads to the elevation of LV filling pressures. Therefore, echocardiographic measurements of diastolic function provide important prognostic information.

In this study Regarding the relation of age to E/E' ratio in both groups, increasing age causes increase in E/E' ratio, (Moderate Positive correlation) and P values are significant (less than 0.05), interestingly, correlation coefficient (R) in non-hypertensive patients is greater than correlation coefficient in hypertensive patients, this means that age has lesser effect on E/E' when a patient has hypertension and vice-versa! Whether this difference is important or not is not well known. This finding needs further studies.

## REFERENCES

1. Kearney PM; Whelton M et al, Global burden of hypertension: analysis of worldwide data. *Lancet* 2005; 365:217-23
2. Kearney PM, et al. Worldwide prevalence of hypertension: a systematic review. Department of Epidemiology, Tulane University School of Public Health and Tropical Medicine, New Orleans, Louisiana 70112, USA. 2004 Jan; 22(1):11
3. Vasan RS et al Residual lifetime risk for developing hypertension in middle-aged women and men: the Framingham heart study. *JAMA* 2002; 287:1003-10]
4. Paulus WJ, Tschöpe C, Sanderson JE, Rusconi C, Flachskampf FA, Rademakers FE, Marino P, Smiseth OA, De Keulenaer G, Leite-Moreira AF, Borbély A, Edes I, Handoko ML, Heymans S, Pezzali N, Pieske B, Dickstein K, Fraser AG, Brutsaert DL. How to diagnose heart failure: a consensus statement on the diagnosis of heart failure with normal left ventricular ejection fraction by the Heart Failure and Echocardiography Associations of the European Society of Cardiology. *Eur Heart J*. 2007; 28: 2539-2550.
5. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini D, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW: ACC/AHA 2005 guideline update for the diagnosis and management of chronic heart failure in the adult. A report of the American College of Cardiology/American Heart Association Task Force on practical guidelines. 2005.
6. European Study Group on Diastolic Heart Failure. How to diagnose diastolic heart failure. *European Heart Journal*, 1998; 19:990-1003.
7. Thomas JD, Garcia MJ, Greenberg NL. Application of color Doppler M-mode echocardiography in the assessment of ventricular diastolic function: potential for quantitative analysis. *Heart Vessels* 1997; 12(suppl) 135-137.
8. Adebisi AA, Aje A, Ogah OS, et al. Left ventricular function parameters in Hypertensives. *J Natl Med Assoc*. 2005 97(1)41-5.
9. Zile MR; Baicu CF; Bonema D. Diastolic heart failure: definitions and terminology; *Prog Cardiovasc Dis*. 2005 Mar-Apr;47(5):307-13
10. Rohde LE; Palombini DV; Polanczyk CA; Goldraich LA; Clausell N A. Hemodynamically oriented echocardiography-based strategy in the treatment of congestive heart failure. *J Card Fail*. 2007 Oct;13(8):618-25

11. Fischer M, et al, Prevalence of left ventricular diastolic dysfunction in the community: Results from a Doppler echocardiographic-based survey of a population sample. *Eur Heart J* 2003; 24: 320-328.
12. Vasan RS; Benjamin EJ; Levy D. Prevalence, clinical features and prognosis of diastolic heart failure: an epidemiologic perspective. *J Am Coll Cardiol* 1995 Dec;26(7):1565-74 failure of the Frank Starling mechanism.
13. Thomas JD, Garcia MJ, Greenberg NL. Application of color Doppler M-mode echocardiography in the assessment of ventricular diastolic function: potential for quantitative analysis. *Heart Vessels* 1997; 12(suppl) 135-137
14. Du Bois D, Du Bois EF (1989) A formula to estimate the approximate surface area if height and weight be known. 1916. *Nutrition* 5: 303-311.
15. Whelton pk, et al. 2017 High blood pressure guideline. *journal of American college of cardiology* .2017;11.006
16. Caballero L, Kou S, Dulgheru R, Gonjilashvili N, Athanassopoulos GD, Barone D, et al. Echocardiographic reference ranges for normal cardiac Doppler data: results from the NORRE Study. *Eur Heart J Cardiovasc Imaging* 2015;16:1031-41
17. 2005 Young Investigator's Award Winner: Assessment of diastolic function in newly diagnosed hypertensives. Masliza M, Daud SM, Khalid. *Ann Acad Med Singapore*. 2005 Dec; 34(11):684-5
18. Kingue S, Mbango GF, Ouankou M. Echocardiographic study of left ventricular hypertrophy in 98 black hypertensives. *Trop Cardiol*. 1993;19:51-55.
19. Slama M, Music D, Varagic J, Frohlich ED. Diastolic dysfunction in hypertension. *Curr Opin Cardiol*. 2002;17:368-373.
20. Russo C, Jin Z, Homma S, Rundek T, Elkind MS, Sacco RL, Di Tullio MR. Effect of obesity and overweight on left ventricular diastolic function: a community-based study in an elderly cohort. *J Am Coll Cardiol*. 2011;57:1368-1374.
21. Bae HK, Choi HS, Sohn S, Shin HJ, Nam JH, Hong YM. Cardiovascular screening in asymptomatic adolescents with metabolic syndrome. *J Cardiovasc Ultrasound*. 2015;23:10-19.
22. Loefer LR, Rosamond WD, Poole C, McNeill AM, Chang PP, Folsom AR, Chambless LE, Heiss G. Association of multiple anthropometrics of overweight and obesity with incident heart failure: the Atherosclerosis Risk in Communities study. *Circ Heart Fail*. 2009;2:18-24.
23. Ávila-Vanzzini N, Machain Leyva CZ, Rodríguez Castellanos LE, Arias Godínez JA, Ruiz Esparza ME, Herrera Bello H. Excessive weight and obesity are associated to intra-ventricular asynchrony: pilot study. *J Cardiovasc Ultrasound*. 2015;23:86-90.
24. Kossaiy A, Nicolas N. Impact of overweight and obesity on left ventricular diastolic function and value of tissue Doppler echocardiography. *Clin Med Insights Cardiol*. 2013;7:43-50.
25. Deurenberg P, Deurenberg-Yap M, Guricci S. Asians are different from Caucasians and from each other in their body mass index/body fat per cent relationship. *Obes Rev*. 2002;3:141-146.
26. Chang CJ, Wu CH, Chang CS, Yao WJ, Yang YC, Wu JS, Lu FH. Low body mass index but high percent body fat in Taiwanese subjects: implications of obesity cutoffs. *Int J Obes Relat Metab Disord*. 2003;27:253-259.
27. Zile MR, Gottdiener JS, Hetzel SJ, McMurray JJ, Komajda M, McKelvie R, et al., Investigators IP. Prevalence and significance of alterations in cardiac structure and function in patients with heart failure and a preserved ejection fraction. *Circulation* 2011;124:2491-501
28. Shah AM, Claggett B, Sweitzer NK, Shah SJ, Anand IS, O'Meara E, et al. Cardiac structure and function and prognosis in heart failure with preserved ejection fraction: findings from the echocardiographic study of the Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist (TOPCAT) Trial. *Circ Heart Fail* 2014;7:740-51