

RESEARCH PAPER

Tinnitus and vertigo in CKD patients in basra hemodialysis center

Haider Kadhem Saeed ¹, Ahmed M. Alabbasi ²

¹ MBChB, F.I.B.M.S, Department of Otolaryngology, Saddam Teaching Hospital, Alzahra Medical College, Basra, Iraq.

² F.R.C.S.F.I.B.M.S, Department of Surgery, College of Medicine, University of Basra, Basra, Iraq.

Received: 05.11.2020

Accepted: 15.05.2021

Abstract

as the relation between hearing impairment and CKD been discussed thoroughly in many literatures, the relationship if present between CKD, hemodialysis, tinnitus and vertigo is not clear in regard to their prevalence, incidence, relation to disease severity and other co-morbid conditions. **Objective:** inspect the relation between hemodialysis and the development of tinnitus and/or vertigo in patients with chronic kidney disease undergoing hemodialysis in addition to the effect of other factors associated with CKD as increase blood urea and creatinine levels, the effect of hypertension and Diabetes mellitus, and the effect of other electrolytes mainly Na level.

Methods: A prospective study conducted in Basra hemodialysis center and otolaryngology departments in Basra Teaching Hospital, Basra, Iraq. Fifty-nine patients with CKD on regular hemodialysis of variable duration enrolled in this study. The statistical analysis done using SPSS 22.

Results: patients mean age was 41.8 ± 9.2 years (range: 17–50 years). Male to Female ratio was 1.27:1. 16.95% of patients had tinnitus only, 16.95% had vertigo only, 30.51% had combination of both tinnitus and vertigo and only 35.59% didn't develop symptoms. There was significant relation between the duration of hemodialysis and the development of tinnitus and vertigo (P value = 0.047). No significant relation was found between age, sex, serum electrolytes, blood urea and serum creatinine.

Conclusion: Tinnitus and/or vertigo are common in patients with CKD on hemodialysis as 64.41% were affected either with one symptom or with both and they were affected by the duration of the disease.

Keywords: CKD, Tinnitus, Vertigo

Corresponding to:

Ahmed M. Alabbasi, F.R.C.S.F.I.B.M.S, Department of Surgery, College of Medicine, University of Basra, Basra, Iraq

✉ ahmed.mhoder@uobasrah.edu.iq

Introduction

Chronic kidney disease (CKD) is characterized by progressive and irreversible loss of kidney function over the period of months and years. Diabetes and hypertension are the main causes in most countries and diabetes accounts for 30–50% of the patients with CKD ¹. As CKD progresses, the formation and accumulation of uremic toxin will lead to adverse biological effects. Uremic

toxin can contribute to inflammation, immune dysfunction, vascular disease and platelet dysfunction ².

The general increase in life span of patients with CKD wither they are on renal replacement therapy or not had led to development of late manifestation hearing system impairment ³. Tinnitus is a perception of sound in the absence of external auditory stimulus and often has a negative effect upon the quality of life. Vertigo is a delusional perception of movement either of one's own body, such as swaying or rotation, or of the environment, or both ^{4,5}.

In addition to antigenic similarity, the cochlea and kidney have similar physiological mechanisms, namely, the active transport of fluid and electrolytes achieved by the stria vascularis in the cochlea and the glomeruli in the kidney [6] 7.

It was previously confirmed that the cochlea is affected by the systemic metabolic, hydroelectrolytic and hormonal alterations that are associated with CKD [8].

Several variables may contribute to the etiopathogenetic mechanisms of tinnitus and vertigo in CKD including factors related to the severity and duration of the disease, electrolyte disturbances,

ototoxic drugs, age, comorbid conditions such as diabetes mellitus and hypertension, and hemodialysis [9-11]. MRI and MRA of the brain of patients with CKD brought light on etiological aspect of vertigo in these patients. Results indicate vascular etiology of vertiginous syndrome in patients on chronic program of hemodialysis [12].

Although the relation between hearing impairment and CKD has been discussed, the relationship if present between CKD, hemodialysis, tinnitus and vertigo is not clear in regards to their prevalence, incidence, relation to disease severity and other co-morbid conditions. The aims of our study are to inspect the relation between tinnitus and vertigo in CKD patients submitted to hemodialysis treatment, and to inspect any changes of the symptoms (tinnitus and vertigo) in association with the duration of CKD and hemodialysis, medical treatment, levels of blood urea, creatinine and other electrolytes.

Methods

Study design and setting

This prospective cross-sectional study was done at the Otolaryngology and Haemodialysis units in Basra Teaching Hospital from December 2014 to December 2016. Written informed consent was obtained from each patient before enrollment in the study following a detailed explanation of the objectives and protocol of the study. This study was approved by the Ethical Committee of the College of Medicine, University of Basra and the ethical committee in Basra Teaching Hospital.

Patients

Chronic kidney disease patients on regular hemodialysis for different durations were interviewed for this study. The inclusion criteria for the patients were a stage 5 CKD on hemodialysis regardless of the duration of the renal failure or the dialysis. The exclusion criteria included patients age >50 years, history of chronic use of ototoxic drugs, chronic suppurative otitis media, acoustic neuroma, childhood-onset hearing loss or hearing loss before the onset of chronic kidney disease, excessive or chronic noise exposure, otological trauma or surgery, renal transplantation, and conductive hearing loss. Eighty-one consecutive patients admitted to the hemodialysis unit were targeted. Fourteen patients were excluded according to the exclusion criteria. In addition, eight patients died during the follow-up period. The remaining 59 patients (33 males and 26 females) representing the final sample size enrolled in the study.

Data collection

The patients interviewed about their sociodemographic characteristics and medical history. Otological examination, pure tone audiometry and tympanometry were done to exclude conductive hearing loss, external and middle ear pathologies. The pure-tone audiometric examinations performed by an audiologist. Symptoms of Tinnitus and/or Vertigo evaluated regarding their duration, medication taken and their corresponding effect on the patient's life.

Statistical analysis

The data analyzed using the SPSS (SPSS Inc., Chicago, IL, USA) version 22. Chi-square test used to find the significance of differences in categorical parameters.

Results

Fifty-nine patients were enrolled in this study (33 [55.9%] men and 26 [44.1%] women). The mean age of the studied population was 41.8 ± 9.2 years (range: 17–50 years). Other sociodemographic findings are illustrated in table [1].

Of the 59 patients enrolled in this study, 10 patients (16.95%) complain from tinnitus, 10 patients (16.95%) complain from vertigo, 18 patients (30.51%) had both tinnitus and vertigo, and the remaining 21 patients (35.59%) didn't develop any tinnitus or vertigo symptoms during the time of the study, as shown in figure [1].

TABLE [1]: THE BASIC VARIABLES OF 59 PATIENTS		
	Variable	Total (59)
1	Mean age in years	41.8 +/- 9.2
2	Male / Female ratio	1.27:1
3	Duration of hemodialysis (years)	3.4±2.9
4	percentage of patients with HTN, DM, or both	94.9 %
5	Duration of CKD (years)	
	< 2 yrs.	27 (45.8%)
	2-5 yrs.	19 (32.2%)
	>5 yrs.	13 (22%)
6	Serum blood urea (mg/dL)	194.8±39.9
7	Serum creatinine (mg/dL)	9.5±1.5
8	Serum sodium (mmol/L)	141.7±8.8
9	Serum potassium (mmol/L)	4.7±0.7
10	Serum calcium (mg/dL)	4.1±0.6




Figure [1]: distribution of symptoms among the patients

Onset of development of tinnitus and/or vertigo symptoms with regards to the start of hemodialysis was also reviewed as shown in table [2].

A significant correlation was found between the duration of hemodialysis and the onset of symptoms development in the involved patients (P-value = 0.047).

No other significant correlations found between the presence of tinnitus and/or vertigo with age, gender, blood urea level, serum creatinine level, serum sodium, serum potassium, serum calcium, comorbidity like HTN, DM and medications.

This is shown in table [3].

TABLE [2]: TIME ONSET OF DEVELOPMENT OF TINNITUS AND/OR VERTIGO SYMPTOMS IN RELATION TO HEMODIALYSIS DURATION.

	Onset of symptoms development	Frequency	Percentage	Cumulative percentage
1	No tinnitus \ vertigo from the start of hemodialysis	21	35.59%	35.59%
2	Tinnitus \ vertigo developed during the 1 st year of hemodialysis	19	32.21%	67.79%
3	Tinnitus \ vertigo developed during 1-5 years of hemodialysis	15	25.42%	93.22%
4	Tinnitus \ vertigo developed after 5 years of hemodialysis	4	6.78%	100%

TABLE [3]: THE SIGNIFICANCE OF DIFFERENT VARIABLES TESTED IN THE STUDY

	The tested variables	P-value
1	Patients age (mean)	0.883
2	Gender	0.647
3	Blood urea level	0.643
4	serum creatinine level	0.634
5	serum sodium	0.285
6	serum potassium	0.125
7	serum calcium	0.458
8	comorbidity like HTN, DM	0.448
9	medications	0.498

Discussion

The effect of CKD and chronic hemodialysis on auditory system had been known since more than 80 years, as Alport had reported the presence of hereditary nephritis associated with sensorineural hearing loss and ocular abnormalities, which he describes as Alport's Syndrome in 1927¹³.

Since then, many studies had been conducted to explain the relationship between CKD, hemodialysis and auditory functions, those were concerned with the hearing and few studies explained their relation to the development of tinnitus and/or vertigo.

alterations in the peripheral and central nervous system, "uremic neuropathy," may be involved in auditory impairment associated with CKD said Albertazzi¹⁴ and Bazzi¹⁵. Di Paolo *et al.*¹⁶ reported a high incidence of nerve conduction dysfunction in patients with CKD. They found decreased conduction velocity in the sensory and motor units, with the sensory units being more compromised than the motor. Da Jung *et al.*²⁵ found CKD to be associated with vestibular dysfunction in the general population. Therefore, the participants with CKD may be closely monitored for vestibular dysfunction. Gaber *et al.*²⁶ found that both the auditory and vestibular pathways are affected in different stages of chronic kidney disease regardless of the disease stage.

This study found that 69.49% of the enrolled patients develop tinnitus, vertigo or both during their treatment with hemodialysis for their CKD. Only 30.51% didn't develop any history of tinnitus and/or vertigo during the study period. This finding was similar to the finding of Sun Kang *et al.*²⁷, Gabr *et al.*²⁶ and which found 68.2% and 60% respectively had either tinnitus or vertigo. Neslihan *et al.*²⁹ found that 15.95% had develop tinnitus alone . While Maria Köping *et al.* [17,18] and Juliana *et al.*¹⁹ found that 58.8%, 45.6% and 71.4% respectively did not develop any symptoms of tinnitus and/or vertigo (Juliana *et al.*¹⁹ said only 20% had tinnitus, 8.6% had combination of tinnitus and light dizziness and no patient develop true vertigo. This might be attributed to the restricted number of patients taken by their study (35) and variable disease duration (from 1 month to 17 year).

In this study, significant positive relationship was found between the duration of hemodialysis and the occurrence of tinnitus and/or vertigo

symptoms among our patients (p value= 0.047). this was similar to the finding of Shih CP et al [20] (P value = 0.001), Peyvandi et al²⁸ and Da Jung Jung et al²⁵ while Juliana et al¹⁹, Bazzi [15] and Neslihan et al²⁹ stated that the duration of dialysis treatment is not a risk factor for tinnitus and vertigo. Again, this might be attributed to the little number of patients enrolled in the study and more frequent home and hospital dialysis sessions.

In this study, we didn't found any significant correlations between the presence of tinnitus and/or vertigo with age, gender, blood urea level, serum creatinine level, serum sodium, serum potassium, serum calcium, comorbidity like HTN, DM and medications. This was similar to the finding of Agarwal *et al*¹⁰, Yassin et al²¹ and Reddy *et al.*²². While, Bazzi¹⁵ and Gatland²³ report that High blood urea and electrolyte disturbances, particularly sodium, have been suggested as possible factors that contribute to hearing system deterioration in CKD.

Conclusion

Despite the limited number of patients enrolled, this study found that both tinnitus and/or vertigo is common in patients with CKD on hemodialysis (up to 69% in this study). The duration of hemodialysis has a significant adverse effect on both auditory and vestibular systems.

Conflict of interest: The authors declare no conflict of interest.

Financial Support: None declared.

References

1. Webster AC, Nagler EV, Morton RL, Masson P. Chronic Kidney Disease. *Lancet*. 2017;389: 1238–1252. doi: 10.1016/S0140-6736(16)32064-5.
2. Consortium CKDP. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet*. 2010; 375: 2073–2081. doi: 10.1016/S0140-6736(10)60674-5.
3. Orendorz-Fraczkowska K, Makulska I, Pospiech L, Zwonlinska D. The influence of Haemodialysis on hearing organ of children with chronic renal failure. *Otolaryngol Pol*. 2002;56(5):597-602.
4. McCormack A, Edmondson-Jones M, Somerset S, Hall D. A systematic review of the reporting of tinnitus prevalence and severity. *Hear Res*. 2016; 337: 70–79. doi: 10.1016/j.heares.2016.05.009.
5. Neuhauser HK. Epidemiology of vertigo. *Curr Opin Neurol*. 2007; 20:40–46.
6. Cosgrove D, Samuelson G, Meehan DT, Miller C, McGee J, Walsh EJ, et al. Ultrastructural, physiological, and molecular defects in the inner ear of a gene-knockout mouse model for autosomal alport syndrome. *HearRes* 1998; 121:84-98.
7. Ozturan O, Lam S. The effect of hemodialysis on hearing using pure-tone audiometry and distortion-product otoacoustic emissions. *ORL J Otorhinolaryngology Relat Spec* 1998; 60:306-13.
8. Pirodda A, Cicero AF, Borghi C. Kidney disease and inner ear impairment: A simpler and closer pathogenic analogy? *Intern Emerg Med* 2012;7(Suppl 2): S93-5.
9. Stavroulaki P, Nikolopoulos TP, Psarommatis I, Apostolopoulos N. Hearing evaluation with distortion-product otoacoustic emissions in young patients undergoing haemodialysis. *Clin Otolaryngol Allied Sci* 2001; 26:235-42.
10. Agarwal MK. A study of otorhinolaryngological manifestations in patients of chronic renal failure. *Indian J Otolaryngol Head Neck Surg* 1997; 49:316-20.
11. Thodi C, Thodis E, Danielides V, Pasadakis P, Vargemezis V. Hearing in renal failure. *Nephrol Dial Transplant* 2006; 21:3023-30.
12. Alajbegovic A1, Resic H, Merhemic Z, Rasic S, Bratic M., The vertigo syndrome, magnetic resonance and magnetic angiography of the head in patients on a chronic hemodialysis program, *Med Arh*. 2001;55(4):227-9.
13. Gleeson MJ. Alport's syndrome: audiological manifestations and implications. *J Laryngol Otol* 1984; 98(5): 449-65.

14. Albertazzi A, Cappelli P, Di Marco T, Maccarone M, Di Paolo B. The natural history of uremic neuropathy. *Contrib Nephrol* 1988; 65:130-7.
15. Bazzi C, Pagani C, Sorgato G, Albonico G, Fellin G, D'Amico G, et al. Uremic polyneuropathy: A clinical and electrophysiological study in 135 short- and long-term haemodialyzed patients. *Clin Nephrol* 1991; 35:176-81.
16. Di Paolo B, Di Marco T, Cappelli P, Spisni C, Del Rosso G, Palmieri PF, et al. Electrophysiological aspects of nervous conduction in uremia. *Clin Nephrol* 1988; 29:253-60.
17. Köping M, Shehata-Dieler W, Cebulla M, et al. Cardiac and renal dysfunction is associated with progressive hearing loss in patients with Fabry disease. *PLoS One*. 2017;12(11): e0188103. Published 2017 Nov 21. doi: 10.1371/journal.pone.0188103
18. Köping, Maria et al. "Characterization of vertigo and hearing loss in patients with Fabry disease." *Orphanet journal of rare diseases* vol. 13,1 137. 15 Aug. 2018, doi:10.1186/s13023-018-0882-7
19. Juliana Altavilla van Petten Machado¹, Gustavo Magalhães Tôrres¹, Alano Nunes Barcellos¹, Licianny Tupinambá Valle², Ludmilla Teixeira², Marconi Teixeira Fonseca³, Incidence of Otologic Symptoms in Patients on Regular Dialysis Treatment, international archive of ORL, Year: 2006, Vol. 10, Num. 1 - Jan/Mar
20. Shih CP, Lin HC, Chung CH, et al. Increased risk of tinnitus in patients with chronic kidney disease: A nationwide, population-based cohort study. *PLoS One*. 2017;12(8): e0183192. Published 2017 Aug 15. doi: 10.1371/journal.pone.0183192
21. Yassin, A., Badry, A., & Fatt-Hi, A. (1970). The relationship between electrolyte balance and cochlear disturbances in cases of renal failure. *The Journal of Laryngology & Otology*, 84(4), 429-435. doi:10.1017/S0022215100072030
22. Reddy EK, Surya Prakash DR, Rama Krishna MG. Proportion of hearing loss in chronic renal failure: Our experience. *Indian J Otol* 2016; 22:4-9.
23. Gatland D, Tucker B, Chalstrey S, Keene M, Baker L. Hearing loss in chronic renal failure-hearing threshold changes following haemodialysis. *J R Soc Med* 1991; 84:587-9.
24. Krajewska (Wojciechowska), J., Krajewski, W. & Zatoński, T. Otorhinolaryngological dysfunctions induced by chronic kidney disease in pre- and post-transplant stages. *Eur Arch Otorhinolaryngol* 277, 1575–1591 (2020)
25. Da Jung Jung, Kyu-Yup Lee, Jun Young Do & Seok Hui Kang (2017) Chronic kidney disease as a risk factor for vestibular dysfunction, *Postgraduate Medicine*, 129:6, 649-652.
26. Gabr, T., Kotait, M., & Okda, H. (2019). Audiovestibular functions in chronic kidney disease in relation to haemodialysis. *The Journal of Laryngology & Otology*, 133(7), 592-599.
27. Sun-Myoung Kang, Hyun Woo Lim & Hoon Yu (2018) Idiopathic sudden sensorineural hearing loss in dialysis patients, *Renal Failure*, 40:1, 170-174.
28. Peyvandi, A., Roozbahany, N.A. Hearing Loss in Chronic Renal Failure Patient Undergoing Hemodialysis. *Indian J Otolaryngol Head Neck Surg* 65, 537–540 (2013).
29. Neslihan BİNEKÇİ, M. Volkan AKDOĞAN, Cihat Burak SAYIN, A. Fuat BÜYÜKLÜ, Tinnitus in Hemodialysis Patients: Cross-Sectional, Clinical Trail, *ournal of ear nose throat and head neck surgery*, KBB ve BBC Dergisi 2018;26(3):69-74.