

Original article

Diagnostic accuracy for coronary artery disease by computed tomography angiography as compared to conventional invasive angiography

Dr. Forat Tariq Youash ¹, Dr. Mohammed Ali Khalaf * ²

¹ MBChB ,Medicine - Iraqi board for medical specializations, ORCID ID:

<https://orcid.org/0009-0004-3495-1805>.

² Assistant Professor, University of Kirkuk, College of medicine, Consultant physician- Internal Medicine, Kirkuk, Iraq. ORCID ID: <https://orcid.org/0000-0002-2225-7419>.

*Correspondence: malbaytee75@uokirkuk.edu.iq

DOI: 10.32894/kjms.2023.139128.1057

Abstract:

- **Background:** Emerging as a major cause of death and disability in the developing countries is coronary artery disease (CAD). Traditional coronary angiography has traditionally been the preferred diagnostic procedure for CAD. Non-invasive evaluation of coronary artery disease (CAD) using computed tomography (CT) coronary angiography shows promise. This observational study's goal is to assess the relative efficacy of noninvasive multi-section computed tomography (MSCT) coronary angiography vs invasive selective coronary angiography in detecting coronary stenosis.
- **Method and patient:** This observational study carried out in the Azadi teaching Hospital/Kirkuk between November 2021 to March 2022. Patients who suspected to have coronary artery disease underwent both CTCA and invasive coronary angiography in different centers in our country comparing and analyzing their results.
- **Result:** In total, 92 participants were enrolled in the trial. Patients had a mean age of 57.207.3 years, and men made up the vast majority of those who appeared (67.3 percent). Chest pain was the most common presenting complaint (94 percent). Patients exhibited a prevalence of 84% for HTN, 55% for hyperlipidaemia, 58% for diabetes, and 4% for ischemic heart disease. In addition, 50% of the patients are either current or former smokers. Half of the patients received 64-slice CT scans, nearly half had 128-slice scans, and 3.2% got 256-slice scans. CCTA's sensitivity was 100%, and its specificity was 66.7%.

Positive predictive value for CCTA was 98.8 percent (meaning it correctly identified positive cases in 98.8 percent of patients) and negative predictive value was 100 percent, for a total diagnostic accuracy of 98.9 percent (which mean CCTA can exclude all negative cases and did not miss any case). When looking at individual arteries, the LAD artery was shown to be the most accurate, while the LCX artery was the least accurate (95.6 percent and 78.2 percent respectively). In terms of sensitivity, LAD performed at a 99.1% rate, while LMS performed at a 0% rate (33.3 percent). RCA had the most specificity (95.2 percent), whereas LAD had the lowest (88.8 percent) (40 percent). In order to determine what factors, influence CCTA positivity, we performed a binary logistic regression analysis for positive CCTA (negative as reference) with all variables from the studies. The results showed that IHD, lesions in the LAD artery, lesions in the LCX artery, and CT slice thickness (128 and 256 slices) all have a negative effect on CCTA (p0.05).

- **Conclusions:** When compared to traditional angiography, multi-slice computed tomography has demonstrated superior sensitivity and specificity for the identification of coronary artery disease.
- **Keywords:** Coronary artery disease, Computed tomography, coronary angiography, Conventional coronary angiography.

INTRODUCTION

Emerging as a major cause of death and disability in the developing countries is coronary artery disease (CAD). Traditional coronary angiography has traditionally been the preferred diagnostic procedure for CAD⁽¹⁾. In the context of catheter-based or surgical treatments, it offers superb resolution for seeing the coronary arteries. Mortality is a very rare complication of this procedure, but it relates to a number of other problems, both cardiac and otherwise⁽²⁾.

Noninvasive evaluation of coronary artery disease (CAD) using computed tomography (CT) coronary angiography shows promise⁽³⁾. coronary artery disease is evaluated, and both qualitative and quantitative data on the presence and extent of non-obstructive atherosclerotic plaque in the arterial wall are provided. This suggests that CT angiography-based disease evaluation may yield more useful information for patients than does standard angiography. High diagnostic accuracy of CT coronary angiography has resulted from the development of multi-slice computed tomography (MSCT) technology, such as 64-slice, 128-slice, 256-slice, and now 320-slice MSCT⁽⁴⁾.

With a negative predictive value constantly exceeding 90%, it has been shown to reliably exclude patients who have severe CAD, whereby coronary luminal stenosis is defined as 50%.

Therefore, the purpose of this study is to compare the diagnostic yield and accuracy of coronary CT angiography (CCTA) with invasive coronary angiography (ICA) in patients referred for invasive coronary angiography due to clinical concern for coronary artery disease (CAD).^(5, 6)

CAD, or coronary artery disease: Atherosclerosis's pathophysiology is still poorly understood. Atherosclerosis likely has a complex pathophysiology due to the interaction of environmental and genetic variables. Some 'new' risk factors and as-yet-unrecognized factors may be as important as the 'traditional' ones as HTN, dyslipidemia, DM, and smoking. The high rate of CAD in Iraq may be a result of any one of these variables acting alone or in various combinations within a genetically susceptible population^(1,7,8). Standard invasive angiography (CIA) has been the gold standard for verifying the diagnosis of CAD for several decades. As part of coronary angiography, a catheter is inserted into an artery in the groin, arm, or shoulder and guided to the heart. Digital X-ray images are taken as iodinated contrast dye is injected directly into the coronary arteries^(9 – 11).




Figure 1. Atherosclerosis of the Coronary Arteries. Coronary angiogram and cardiac catheterization demonstrating significant left anterior descending (LAD) stenosis in the left panel. As can be seen in the right panel, a stent was placed in the left anterior descending coronary artery to treat this lesion⁽¹³⁾

The diagnostic accuracy of invasive coronary angiography has various drawbacks. Visual estimations of stenosis severity are common, but they have their limitations due to inter-observer variability that can vary from 30 to 60 percent. As the stenosis areas are reported as a percentage of luminal diameter in comparison with surrounding normal coronary segments, and, in diffuse disease, no such segments are detected, the presence of diffuse disease may also contribute to underestimating of stenosis⁽¹²⁻¹⁴⁾.

The fractional flow reserve computed tomography (FFRCT) approach has recently emerged as a non-invasive method for assessing the severity of epicardial coronary narrowing and the physiological consequences of this narrowing^(15,16).

To get around the drawbacks of CIA, multi-slice computed tomography angiography (CTA) has been recommended in recent years as an alternative method of diagnosing coronary blockages. Recent advances in technology with many detectors have greatly enhanced spatial and temporal resolutions of pictures, making CTA the focus of doctors' attention ⁽¹⁷⁾ This observational study's goal is to assess the relative efficacy of noninvasive multi-section computed tomography (MSCT) coronary angiography vs invasive selective coronary angiography in detecting coronary stenosis ⁽²²⁾, This observational study's goal is to assess the relative efficacy of noninvasive multi-section computed tomography (MSCT) coronary angiography vs invasive selective coronary angiography in detecting coronary stenosis ⁽¹⁸⁻²¹⁾.

PATIENT and METHOD

This research was conducted from November 2021 to March 2022 at the Azadi Teaching Hospital in Kirkuk. All patients who fulfilled the following inclusion criteria were included in the study: Patient must be 18 years or older. Suspect angina related to coronary heart disease because of chest pain. Individuals who are at a moderate risk for developing coronary artery disease but not at a high risk. Criteria for exclusion: Those who are unable or unwilling to undergo CTCA. People who have developed an episode of acute coronary syndrome (up to 100 days previously). A glomerular filtration rate of 30 ml/min or below indicates chronic kidney disease. Patients who are pregnant women. Systemic comorbidity individuals, such as those with severe COPD, who cannot hold their breath while imaging. Direct questionnaire interviews are conducted with each respondent to acquire the following data: In the first section, you'll be asked questions about your demographics and your health. Give your full name, date of birth, gender, job title, and a brief introduction. Examining the presence of many disorders: ischemic heart disease, diabetes, high cholesterol, high blood pressure, high blood sugar, high body mass index, and a normal electrocardiogram all point to a healthy lifestyle.

Part 2 involves comparing and analyzing the outcomes of CTCA and invasive coronary angiography in patients who gave their informed consent for both procedures. Multiple observers, each of whom was uninformed of the multi-slice CT results, classified coronary lesions as either single-vessel, double-vessel, or triple-vessel disease. To be compared with invasive angiography, lesions of any size are included. According to the degree to which an artery has been blocked, lesions are categorized.

The majority of our patients' CTCA and CIA were performed in the city of Kirkuk in northern Iraq, and the interval between their tests ranged from a few weeks to three months. Regarding the ethical considerations, Approval was taken from Internal Medicine Scientific Committee of Iraqi Board, an agreement for research was taken from hospitals' authorities, written informed

consent was taken from each patient. The statistical package for social sciences (SPSS) software version 23 had been used for data entry and analysis. In the descriptive statistics for socio-demographic characteristics, the means, standard deviations, min, max values were used for continuous data. Numbers and percentage values were used for countable data. In analyzing the differences between the groups, chi square and T test were used. Binary logistic analysis was used to evaluate the possible factors associated with positive CCTA. $P < 0.05$ was used as the threshold for statistical significance.

RESULTS

There were 92 patients included in this study.

Demographic characteristics: The mean age of patients was 57.2 ± 7.3 years (range 44 – 75 years).

Table 1. Descriptive statistics of age

	Mean	SD	Minimum	Maximum
Age	57.2	7.3	44	75

Majority of presented patients were males (67.3%) (Figure 3.1).




Figure 1. Gender distribution across patients

The majority of patients were presented with chest pain (94%). There were 85%, 55%, 58%, and 4% of patients had HTN, hyperlipidemia, DM, and IHD respectively. Also, 50% of patients are currently smokers or ex-smokers (Table 2).

Table 2. Demographic characteristics of patients

		Count	N %
Chest Pain	No	5	5.4%
	Yes	87	94.6%
Dyspnea	No	45	48.9%
	Yes	47	51.1%
Palpitation	No	54	58.7%
	Yes	38	41.3%
Occupation	Employee	23	25.0%
	Free worker	30	32.6%
	Housewife	27	29.3%
	Retired	10	10.9%
	Teacher	2	2.2%
HTN	No	13	14.1%
	Yes	79	85.9%
Hyperlipidemia	No	41	44.6%
	Yes	51	55.4%
DM	No	38	41.3%
	Yes	54	58.7%
IHD	No	88	95.7%
	Yes	4	4.3%
Smoking	Yes	34	37%
	Ex smoke	12	13.0%
	No	46	50.0%

Regarding vital signs and BMI, the descriptive statistics are given in below table 3.

Table 3. Descriptive statistics of vital signs and BMI

	Mean	SD	Minimum	Maximum
SBP	138	13	110	170
DBP	82	11	60	100
HR	78	8	56	105
RR	14	1	10	16
BMI	27	3	23	36

Regarding the ECG changes, majority of patients presented with ST segment changes (75%) (Figure 2).




Figure 2. ECG changes across patients

Coronary CT angiography (CCTA): There were 50% of patients had 64-slice CT, 46.7% of patients had 128-slice CT, and 3.2% patients had 256-slice CT (Figure 3).




Figure 3. Type of CT performed.

The sensitivity of CCTA was 100% with specificity of 66.7%. The diagnostic accuracy of CCTA was 98.9% with PPV of 98.8% (which mean CCTA can identify positive cases in 98.8% of patients) and NPV of 100% (which mean CCTA can exclude all negative cases and did not miss any case).

Table 4. Comparison between CCTA and Conventional invasive angiography for diagnosis of CAD

		Conventional invasive angiography		Total
		Positive	Negative	
CT	Positive	89 (100%)	1 (33.3%)	90 (97.8%)
	Negative	0	2 (66.7%)	2 (2.2%)
Total		89 (100%)	3 (100%)	92 (100%)

P=0.0001

Table 5. Diagnostic accuracy of CCTA based on conventional invasive angiography

Test	Value
Sensitivity	100%
Specificity	66.7%
Positive predictive value	98.8%
Negative predictive value	100%
Accuracy	98.9%

For specific artery accuracy, the highest accuracy was observed for LAD artery while LCX artery showed lowest accuracy (95.6% and 78.2% respectively). The highest sensitivity was observed with LAD (98.9%) while LMS showed lowest sensitivity (33.3%). The specificity was at its highest level with RCA (95.2%) while lowest specificity was observed with LAD (40%).

Table 6. Comparison between CCTA and conventional invasive angiography for diagnosis of CAD per location

Location	CCTA		Angiography		P value
	Positive	No lesion	Positive	No lesion	
LMS	10 (10.9%)	82 (89.1%)	3 (3.3%)	89 (96.7%)	0.2
LAD	89 (96.7%)	3 (3.3%)	87 (94.6%)	5 (5.4%)	0.001
RCA	42 (45.7%)	50 (54.3%)	50 (54.3%)	42 (45.7%)	0.001
LCX	60 (65.2%)	32 (34.8%)	54 (58.7%)	38 (41.3%)	0.001

Table 7. Diagnostic accuracy of CCTA based on conventional invasive angiography per location

Artery	Sensitivity	Specificity	PPV	NPV	Accuracy
LMS	33.3%	89.9%	90%	97.6%	88%
LAD	98.9%	40%	96.6%	66.7%	95.6%
RCA	80%	95.2%	95.2%	80%	86.9%
LCX	87%	65.8%	78.3%	78.1%	78.2%

Factors associated with CCTA positivity: By running binary logistic regression analysis for positive CCTA (negative as reference) with all studies variables to test the factors that impact CCTA positivity, the test showed that, IHD, lesion in LAD artery, lesion in LCX artery, and type of CT (128 and 256 slices) effect negatively CCTA ($p < 0.05$) (Table 8).

Table 8. Factors associated with CCTA positivity

		Score	DF	P value
Variables	Age	0.003	1	0.959
	Sex	0.989	1	0.320
	Occupation	6.133	4	0.189
	HTN	0.336	1	0.562
	Hyperlipidemia	1.644	1	0.200
	DM	1.439	1	0.230
	IHD	44.978	1	0.001
	Smoking	2.044	1	0.432
	SBP	0.628	1	0.428
	DBP	3.087	1	0.079
	HR	0.316	1	0.574
	RR	2.870	1	0.090
	BMI	1.307	1	0.253
	ECG Changes	0.681	3	0.878
	LMS	0.249	1	0.618
	LAD	60.652	1	0.001
	RCA	1.717	1	0.190
	LCX	3.833	1	0.050
	Type of CT	14.723	2	0.001
	Chest Pain	0.117	1	0.732
Dyspnea	2.135	1	0.144	
Palpitation	1.439	1	0.230	
Overall Statistics		77.802	41	0.001

There was no significant association between the comorbidities and CT nor O2 saturation (P>0.05) (Table 9).

Table 9. Group Statistics comparison of BMI with CT and O2 saturation

	Variable	N	Mean	SD	P value
C.T scan	Hypertension	36	67.78	15.741	0.76
	No	64	66.80	15.207	
O2 sat.	Hypertension	36	83.92	5.674	0.71
	No	64	83.44	7.536	
C.T scan	DM	27	70.37	14.539	0.2
	No DM	73	65.96	15.538	
O2 sat.	DM	27	82.44	5.235	0.3
	No DM	73	84.04	7.404	
C.T scan	Others	10	63.00	18.439	0.36
	No comorbidities	90	67.61	14.995	
O2 sat.	Others	10	85.10	6.903	0.47

DISCUSSION

When it comes to visualizing coronary artery disease, CCTA is state-of-the-art. CCTA is useful for assessing heart chambers, valves, great vessels, pericardium, and surrounding tissues, even though coronary atherosclerosis is the primary cause for the burden of cardiovascular illnesses⁽²⁾. This study was created to evaluate CCTA's diagnostic performance specifically for coronary arteries.

Among all patients, there was a mean age of 57.27.3 years. This matched the results of a recent study by Mohammad et al. in Iraq, which found that CAD patients, on average, were 55 years old⁽¹³⁾. Another study by Mohammad et al. found that patients with advanced CAD had a mean age of 63, thus ours is in line with that. 9 Additionally, Elliott et al. found that the average age of CAD patients was 55 in research conducted in the United Kingdom. These findings lend credence to the widely held belief that coronary artery disease is a leading cause of death and disability among adults⁽⁵⁶⁾.

Most patients in this study were male, which is consistent with a study conducted in Iraq by Mohammad et al. that found 62% of adults with CAD were male⁽¹³⁾.

13 Elliott et al. conducted research in the UK and found that 58% of CAD patients were female. 9 Possible explanations for this discrepancy include varying sample sizes and underlying genetic variables⁽⁹⁾.

The most common presenting complaint was chest pain (94 percent). Historically, coronary artery disease has been represented by angina, which is a pain in the sub-sternal region that is described as a squeezing or heavy feeling and that may radiate to the medial aspect of the left upper extremity, to the neck, or to the jaw, and that may be accompanied by nausea, vomiting, palpitations, diaphoresis, syncope, or even sudden death. Even if a patient presents with seemingly unrelated symptoms, doctors and other medical professionals should keep their guard up and treat them with caution⁽⁵²⁾.

Patient prevalence of hypertension, elevated cholesterol levels, diabetes mellitus, and ischemic heart disease was 84%, 55%, 58%, and 4%, respectively. The most common risk factor was found to be hypertension (55.3%), followed by dyslipidemia (42.7%), type 2 diabetes mellitus (29%), smoking (11%), and ex-smoking (22%). (9.3 percent). A possible explanation for the discrepancy could be because the two-research used different sized samples⁽¹³⁾.

Carbon monoxide, tar, nitrosamines, polycyclic aromatic hydrocarbons, hydrogen cyanide, aldehydes, and heavy metals are only few of the thousands of toxic chemicals found in tobacco. Evidence from epidemiological and clinical investigations shows that cigarette smoking increases the risk of cardiovascular events in a dose- and time-dependent manner. Acute myocardial infarction occurred roughly ten years earlier in smokers⁽⁵³⁾.

CCTA's sensitivity was 100%, and its specificity was 66%. CCTA has a 98.9% sensitivity, a 98.8% specificity (meaning it can detect positive cases in 98.8% of patients), and a 100% NPV (meaning it can detect no false positives) (which mean CCTA can exclude all negative cases and did not miss any case). Similar results were found in another study by Joshi et al., which found a sensitivity and specificity of 100% (95 percent confidence interval [CI]: 39.76 percent -100 percent) and 91.30 percent (95 percent CI: 79.21 percent -97.58 percent) for CT angiography, respectively. CT angiography had a strong positive predictive value (50 percent; 95% CI: 15.70 percent to 84.30 percent) and a high negative predictive value (100 percent; 95% CI: 91.59 percent to 100 percent) in this group of patients ^(40,41).

In addition, Mannan et al. found that CCTA has a sensitivity of 83.8% for detecting stenosis, a specificity of 98.4%, and an accuracy of 96.5%; based on these results, the authors concluded that CCTA permits the identification of coronary stenosis with excellent accuracy and well correlation with conventional angiogram. Despite this, Mannan et al. relied on only 64 CCTA slices in their analysis ^(43,45,46). In addition, the positive CCTA was shown by Meinel et al. to have a positive predictive value of 83% for individuals with clinically significant CAD, 100% sensitivity, and 84% specificity ^(47,48).

The diagnostic accuracy of coronary computed tomography angiography for coronary artery disease (CAD) was also demonstrated by Nielsen et al. to be higher than that of standard exercise electrocardiography and single-photon emission computed tomography ⁽⁴⁷⁾.

Wu et al. found that coronary CT coronary angiography (CTCA) was more sensitive than the 'standard diagnostic technique' in identifying patients with angina owing to coronary heart disease while also being less expensive ^(48,49).

In addition, a recent Egyptian study by Abd Ella et al indicated that CTCA has a sensitivity, specificity, and accuracy of 97.1, 96.8, and 96.8 percent, respectively, when compared to traditional angiography for diagnosing severe stenosis in coronary arteries ^(36,37).

Based on 64-slice CT evaluations of 923 segments from 68 patients, Nikolaou et al. found a sensitivity of 82%, specificity of 91%, a positive predictive value of 72%, a negative predictive value of 97%, and an accuracy of 93%. On a sample of 113 patients, Kerl et al. found that 64-slice CT had a sensitivity of 90.5%, specificity of 98.4%, positive predictive value (PPV) of 85.5%, negative predictive value (NPV) of 99.0%, and accuracy of 97.7% ⁽⁴⁰⁾.

Accuracy for individual arteries varied, with the LAD artery showing the most precision and the LCX artery the least (95.6 percent and 78.2 percent respectively). LAD demonstrated the most sensitivity (98.9%), whereas LMS had the lowest (33.3 percent). The RCA had the most specificity (95.2%), while the LAD had the lowest (40 percent). Similar findings were seen in a recent Egyptian study by Abd Ella et al., demonstrating varying sensitivity and specificity with respect to particular artery obstruction ⁽⁵⁴⁾.

To further investigate the factors that influence CCTA positivity, we ran a binary logistic regression for positive CCTA (negative as reference) with all study variables. The results showed that IHD, LAD artery lesion, LCX artery lesion, and CT slice count (128 vs. 256) all have a negative effect on CCTA positivity. These factors may influence the outcomes; however, the size of the sample may be more influential ^(30,31).

In a study comparing CCTA and invasive coronary angiography, Knaapen concluded that CCTA was close but not close enough to replace invasive coronary angiography, but that over the past decade, noninvasive computed tomographic coronary angiography (CCTA) has emerged as a viable alternative to its invasive counterpart ⁽⁵³⁾. A negative CCTA of adequate quality virtually excludes out obstructive CAD and considerably minimizes unnecessary invasive treatments, which is especially important given the low yield of ICA for detecting CAD ^(52,53). However, CCTA is used more as a screening tool for ICA rather than as a replacement for it because of worse image quality due to reduced spatial and temporal resolution, motion and blooming artifacts, and the need for a low and steady heart rate for optimal image capture. Due to its low specificity and high false-positive rate, CCTA has always been its own downfall, and further diagnostic methods are warranted in the event of a positive scan. After a CCTA that is either inconclusive or positive, myocardial perfusion imaging has been recommended to boost specificity. However, this approach has been shown to contribute little to no value over CCTA alone to identify CAD, and instead just boosts specificity at the expense of sensitivity without significantly improving total diagnosis accuracy, and therefore may be regarded even un useful for clinical decision-making. Now that we're moving away from these hybrid or sequential imaging methodologies, we can put our energy into perfecting CCTA as a stand-alone diagnostic tool that can potentially replace ICA ^(50,53).

It's interesting to note that a March 2022 study by Li et al. found that CCTA's diagnostic performance changes depending on the level of experience of the reading physician. It has been demonstrated that as one's experience with CCTA grows, so does its diagnostic precision. Few patients had 256-slice CCTA, and the sample size was limited overall ⁽⁵⁴⁾.

Financial support and sponsorship:

Nil.

Conflicts of interest:

There are no conflicts of interest.

CONCLUSION

1. We found that CCTA has a sensitivity of 100% and a specificity of 66.7% in our investigation.
2. CCTA has a 98.9% sensitivity, a 98.8% specificity (meaning it can detect positive cases in 98.8% of patients), and a 100% NPV (meaning it can detect no false positives) (which mean CCTA can exclude all negative cases and did not miss any case).
3. When looking at individual arteries, the accuracy of the LAD was highest while that of the LCX was lowest. Contrary to LMS, LAD was shown to have the highest sensitivity. The sensitivity was highest for RCA and lowest for LAD.

RECOMMENDATION

1. To diagnose CAD and provide crucial information to guide treatment, therapy, and prognostic evaluation in patients with suspected CAD, CT coronary angiography is a first-line, noninvasive diagnostic test alternative to ICA.
2. In individuals with a moderate to low pre-test risk of CAD, for whom a diagnosis or exclusion of the condition cannot be made based on clinical examination alone, this test is invaluable.
3. In individuals experiencing chest pain other than from a heart attack, it can be used to quickly rule out coronary artery disease.
4. The impact of CT (128 and 256 slices) on diagnostic precision: a large cohort study.

REFERENCES

1. Kohsaka S, Makaryus AN. Coronary angiography using noninvasive imaging techniques of cardiac CT and MRI. *Current cardiology reviews*. 2008 Nov 1;4(4):323-30. doi: 10.2174/157340308786349444.
2. Tavakol M, Ashraf S, Brener SJ. Risks and complications of coronary angiography: a comprehensive review. *Global journal of health science*. 2012 Jan;4(1):65. doi: 10.5539/gjhs.v4n1p65.

3. Hamilton-Craig CR, Friedman D, Achenbach S. Cardiac computed tomography—Evidence, limitations and clinical application. *Heart, Lung and Circulation*. 2012 Feb 1;21(2):70-81. doi: 10.1016/j.hlc.2011.08.070.
4. Otero HJ, Steigner ML, Rybicki FJ. The “post-64” era of coronary CT angiography: understanding new technology from physical principles. *Radiologic clinics of North America*. 2009 Jan 1;47(1):79-90. doi: 10.1016/j.rcl.2008.11.001.
5. Achenbach S, Ulzheimer S, Baum U, Kachelrieß M, Ropers D, Giesler T, Bautz W, Daniel WG, Kalender WA, Moshage W. Noninvasive coronary angiography by retrospectively ECG-gated multislice spiral CT. *Circulation*. 2000 Dec 5;102(23):2823-8. doi: 10.1161/01.cir.102.23.2823.
6. Zou KH, O'Malley AJ, Mauri L. Receiver-operating characteristic analysis for evaluating diagnostic tests and predictive models. *Circulation*. 2007 Feb 6;115(5):654-7. doi: 10.1161/CIRCULATIONAHA.105.594929.
7. Islam AM, Majumder AA. Coronary artery disease in Bangladesh: A review. *Indian heart journal*. 2013 Jul 1;65(4):424-35.
8. Kannel WB. Searching for answers to ethnic disparities in cardiovascular risk. *The Lancet*. 2000;356(9226):266-7.
9. Mohammad AM, Sheikho SK, Tayib JM. Relation of cardiovascular risk factors with coronary angiographic findings in Iraqi patients with ischemic heart disease. *American Journal of Cardiovascular disease research*. 2013;1(1):25-9.
10. Mohammad AM, Rashad HH, Habeeb QS, Rashad BH, Saeed SY. Demographic, clinical and angiographic profile of coronary artery disease in kurdistan region of Iraq. *American Journal of Cardiovascular Disease*. 2021;11(1):39.

11. 1- Dodani S, Henkhaus R, Wick J, Vacek J, Gupta K, Dong L, Butler MG. Metabolic syndrome in South Asian immigrants: more than low HDL requiring aggressive management. *Lipids in Health and Disease*. 2011 Dec;10(1):1-0.
12. Samani NJ, Sharma P. Coronary heart disease in South Asian populations-the role of genetics. *The epidemic of coronary heart disease in South Asian populations: causes and consequences*. 2004:81.
13. Shah A, Hernandez A, Mathur D, Budoff MJ, Kanaya AM. Adipokines and body fat composition in South Asians: results of the Metabolic Syndrome and Atherosclerosis in South Asians Living in America (MASALA) study. *International journal of obesity*. 2012 Jun;36(6):810-6.
14. Ranjith N, Pegoraro RJ, Shanmugam R. Obesity-associated genetic variants in young Asian Indians with the metabolic syndrome and myocardial infarction: cardiovascular topics. *Cardiovascular Journal of Africa*. 2011 Jan 1;22(1):25-30.
15. Jia H, Abtahian F, Aguirre AD, Lee S, Chia S, Lowe H, Kato K, Yonetsu T, Vergallo R, Hu S, Tian J. In vivo diagnosis of plaque erosion and calcified nodule in patients with acute coronary syndrome by intravascular optical coherence tomography. *Journal of the American College of Cardiology*. 2013 Nov 5;62(19):1748-58.
16. Einstein AJ, Henzlova MJ, Rajagopalan S. Estimating risk of cancer associated with radiation exposure from 64-slice computed tomography coronary angiography. *Jama*. Jul 18 2007;298(3):317-323.
17. Thompson RC, Cullom SJ. Issues regarding radiation dosage of cardiac nuclear and radiography procedures. *J Nucl Cardiol*. Jan-Feb 2006;13(1):19-23.
18. Xu Y, Tang L, Zhu X, Xu H, Tang J, Yang Z, Wang L, Wang D. Comparison of dual-source CT coronary angiography and conventional coronary angiography for detecting coronary artery disease. *The international journal of cardiovascular imaging*. 2010 Feb;26(1):75-81.

19. Mowatt G, Cook JA, Hillis GS, Walker S, Fraser C, Jia X, Waugh N. 64-Slice computed tomography angiography in the diagnosis and assessment of coronary artery disease: systematic review and meta-analysis. *Heart*. 2008 Nov 1;94(11):1386-93.
20. Budoff, M.J., Dowe, D., Jollis, J.G., Gitter, M., Sutherland, J., Halamert, E., Scherer, M., Bellinger, R., Martin, A., Benton, R. and Delago, A., 2008. Diagnostic performance of 64-multidetector row coronary computed tomographic angiography for evaluation of coronary artery stenosis in individuals without known coronary artery disease. *Journal of the American College of Cardiology*, 52(21), pp.1724-1732.
21. Johnson NP, Kirkeeide RL, Gould KL. Coronary anatomy to predict physiology: fundamental limits. *Circulation: Cardiovascular Imaging*. 2013 Sep;6(5):817-32.
22. Prati F, Di Vito L, Biondi-Zoccai G, Occhipinti M, La Manna A, Tamburino C, Burzotta F, Trani C, Porto I, Ramazzotti V, Imola F. Angiography alone versus angiography plus optical coherence tomography to guide decision-making during percutaneous coronary intervention: the Centro per la Lotta contro l'Infarto-Optimisation of Percutaneous Coronary Intervention (CLI-OPCI) study. *EuroIntervention*. 2012 Nov 22;8(7):823-9.
23. Dehkordi MT, Sadri S, Doosthoseini A. A review of coronary vessel segmentation algorithms. *Journal of medical signals and sensors*. 2011 Jan;1(1):49.
24. Miller JM, Rochitte CE, Dewey M, Arbab-Zadeh A, Niinuma H, Gottlieb I, Paul N, Clouse ME, Shapiro EP, Hoe J, Lardo AC. Diagnostic performance of coronary angiography by 64-row CT. *New England Journal of Medicine*. 2008 Nov 27;359(22):2324-36.
25. Voros S, Rinehart S, Qian Z, Joshi P, Vazquez G, Fischer C, Belur P, Hulten E, Villines TC. Coronary atherosclerosis imaging by coronary CT angiography: current status, correlation with intravascular interrogation and meta-analysis. *JACC: Cardiovascular Imaging*. 2011 May;4(5):537-48.

26. Papadopoulou SL, Girasis C, Dharampal A, Farooq V, Onuma Y, Rossi A, Morel MA, Krestin GP, Serruys PW, de Feyter PJ, Garcia HM. CT-SYNTAX score: a feasibility and reproducibility Study. *JACC: Cardiovascular Imaging*. 2013 Mar;6(3):413-5.
27. Farooq V, Van Klaveren D, Steyerberg EW, Meliga E, Vergouwe Y, Chieffo A, Kappetein AP, Colombo A, Holmes Jr DR, Mack M, Feldman T. Anatomical and clinical characteristics to guide decision making between coronary artery bypass surgery and percutaneous coronary intervention for individual patients: development and validation of SYNTAX score II. *The Lancet*. 2013 Feb 23;381(9867):639-50.
28. Cavalcante R, Sotomi Y, Lee CW, Ahn JM, Farooq V, Tateishi H, Tenekecioglu E, Zeng Y, Suwannasom P, Collet C, Albuquerque FN. Outcomes after percutaneous coronary intervention or bypass surgery in patients with unprotected left main disease. *Journal of the American College of Cardiology*. 2016 Sep 6;68(10):999-1009.
29. Decramer I, Vanhoenacker PK, Sarno G, Van Hoe L, Bladt O, Wijns W. Effects of sublingual nitroglycerin on coronary lumen diameter and number of visualized septal branches on 64-MDCT angiography. *AJR Am J Roentgenol*. 2008 Jan. 190(1):219-25.
30. Schuetz GM, Zacharopoulou NM, Schlattmann P, Dewey M. Meta-analysis: noninvasive coronary angiography using computed tomography versus magnetic resonance imaging. *Ann Intern Med*. 2010 Feb 2. 152(3):167-77.
31. Taylor AJ, Cerqueira M, Hodgson JM, et al. ACCF/SCCT/ACR/AHA/ASE/ASNC/NASCI/SCAI/SCMR 2010 .Appropriate Use Criteria for Cardiac Computed Tomography. A Report of the American College of Cardiology Foundation. 2010 Nov 23. 122(21):e525-55.
32. Abdulla J, Abildstrom SZ, Gotzsche O, Christensen E, Kober L, Torp-Pedersen C. 64-multislice detector computed tomography coronary angiography as potential alternative to conventional coronary angiography: a systematic review and meta-analysis. *European heart journal*. 2007 Dec 1;28(24):3042-50.

33. Mowatt G, Cummins E, Waugh N, Walker S, Cook JA, Jia X, Hillis GS, Fraser CM. Systematic review of the clinical effectiveness and cost-effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of coronary artery disease. *Health Technology Assessment* 2008; 12: 17.
34. Paech DC, Weston AR. A systematic review of the clinical effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of suspected coronary artery disease. *BMC cardiovascular disorders*. 2011 Dec;11(1):1-1.
35. Stein PD, Yaekoub AY, Matta F, Sostman HD. 64-slice CT for diagnosis of coronary artery disease: a systematic review. *The American journal of medicine*. 2008 Aug 1;121(8):715-25.
36. Sajjadih A, Hekmatnia A, Keivani M, Asoodeh A, Pourmoghaddas M, Sanei H. Diagnostic performance of 64-row coronary CT angiography in detecting significant stenosis as compared with conventional invasive coronary angiography. *ARYA atherosclerosis*. 2013 Mar;9(2):157.
37. Wever-Pinzon O, Romero J, Kelesidis I, et al. Coronary computed tomography angiography for the detection of cardiac allograft vasculopathy: a meta-analysis of prospective trials. *J Am Coll Cardiol*. 2014 May 20. 63(19):1992-2004.
38. Pugliese F, Mollet NR, Hunink MM, Cademartiri F, Nieman K, van Domburg RT, Meijboom WB, Van Mieghem C, Weustink AC, Dijkshoorn ML, de Feyter PJ. Diagnostic performance of coronary CT angiography by using different generations of multisection scanners: single-center experience. *Radiology*. 2008 Feb;246(2):384-93.
39. Steigner ML, Otero HJ, Cai T, Mitsouras D, Nallamshetty L, Whitmore AG, Ersoy H, Levit NA, Di Carli MF, Rybicki FJ. Narrowing the phase window width in prospectively ECG-gated single heart beat 320-detector row coronary CT angiography. *The international journal of cardiovascular imaging*. 2009 Jan;25(1):85-90.

40. Mori S, Endo M, Nishizawa K, Murase K, Fujiwara H, Tanada S. Comparison of patient doses in 256-slice CT and 16-slice CT scanners. *The British journal of radiology*. 2006 Jan;79(937):56-61.
41. Neves PO, Andrade J, Monção H. Coronary artery calcium score: current status. *Radiologia brasileira*. 2017 May;50:182-9.
42. Elliott J, Bodinier B, Bond TA, Chadeau-Hyam M, Evangelou E, Moons KG, Dehghan A, Muller DC, Elliott P, Tzoulaki I. Predictive accuracy of a polygenic risk score–enhanced prediction model vs a clinical risk score for coronary artery disease. *Jama*. 2020 Feb 18;323(7):636-45.
43. Lopez EO, Ballard BD, Jan A. Cardiovascular disease. InStatPearls [Internet] 2021 Aug 11. StatPearls Publishing.
44. Chen HY, Li SC, Chen LF, Wang W, Wang Y, Yan XW. The effects of cigarette smoking and smoking cessation on high-density lipoprotein functions: implications for coronary artery disease. *Annals of clinical biochemistry*. 2019 Jan;56(1):100-11.
45. Joshi H, Shah R, Prajapati J, Bhangdiya V, Shah J, Kandre Y, Shah K. Diagnostic accuracy of computed tomography angiography as compared to conventional angiography in patients undergoing noncoronary cardiac surgery. *Heart Views: The Official Journal of the Gulf Heart Association*. 2016 Jul;17(3):88.
46. Mannan N, Basher MA, Mohammad J, Jahan MU, Momenuzzaman NA, Haque MA. Comparison of coronary CT angiography with conventional coronary angiography in the diagnosis of coronary artery disease. *Bangladesh Medical Research Council Bulletin*. 2014 Sep 5;40(1):31-5.
47. Meinel FG, Schoepf UJ, Townsend JC, Flowers BA, Geyer LL, Ebersberger U, Krazinski AW, Kunz WG, Thierfelder KM, Baker DW, Khan AM. Diagnostic yield and accuracy of coronary CT angiography after abnormal nuclear myocardial perfusion imaging. *Scientific Reports*. 2018 Jun 15;8(1):1-0.

48. Nielsen LH, Ortner N, Nørgaard BL, Achenbach S, Leipsic J, Abdulla J. The diagnostic accuracy and outcomes after coronary computed tomography angiography vs. conventional functional testing in patients with stable angina pectoris: a systematic review and meta-analysis. *European Heart Journal–Cardiovascular Imaging*. 2014 Sep 1;15(9):961-71.
49. Wu Z, He Y, Li W, Cheng S. Computed tomography coronary angiography vs. standard diagnostic procedure for the diagnosis of angina due to coronary heart disease: A cross-sectional study. *Experimental and Therapeutic Medicine*. 2019 Apr 1;17(4):2485-94.
50. Abd Ella TF, Elsayed EE, ELkersh AM, Moawad MM. Diagnostic accuracy of computed tomography angiography compared with catheter angiography in coronary artery disease. *Menoufia Medical Journal*. 2021 Jul 1;34(3):1034.
51. Nikolaou K, Knez A, Rist C, Wintersperger BJ, Leber A, Johnson T, Reiser MF, Becker CR. Accuracy of 64-MDCT in the diagnosis of ischemic heart disease. *American Journal of Roentgenology*. 2006 Jul;187(1):111-7.
52. Kerl JM, Schoepf UJ, Zwerner PL, Bauer RW, Abro JA, Thilo C, Vogl TJ, Herzog C. Accuracy of coronary artery stenosis detection with CT versus conventional coronary angiography compared with composite findings from both tests as an enhanced reference standard. *European radiology*. 2011 Sep;21(9):1895-903.
53. Knaapen P. Computed tomography to replace invasive coronary angiography? Close, but not close enough. *Circulation: Cardiovascular Imaging*. 2019 Feb;12(2):e008710.
54. Li F, He Q, LIXUE X, Zhou Y, Sun Y, Xu Y, et al. Diagnostic Accuracy of Subtraction Coronary CT Angiography in Patients with Severe Calcification: Comparison between Readers with Different Levels of Experience. *Frontiers in Cardiovascular Medicine*. 2022;9:184.