



Antibiotic resistances of *Salmonella* spp. in rectal samples from farm animals in Al-Diwaniyah City, Iraq

Hiba Shehab Ahmed

College of Biotechnology/ University of AL-Qadisiyah , Iraq

Corresponding Author Email: hiba.shehab@qu.edu.iq

Abstract

The principles of this work were to isolate the bacterium *Salmonella* and evaluate its antibiotic resistance capability from rectal fecal samples of some types of animals from Al-Diwaniyah City, Iraq. A total of 38 randomly collected samples (13, 16, and 9) from healthy cows, sheep, and goats, respectively, were employed to take samples via the use of commercially-available sterile cotton swabs. The samples were cultivated using different agar media. Then, the suspected *Salmonella* growth from the nutrient agar was subjected to biochemical identification. Later, antibiotic susceptibility assay was performed on the pure *Salmonella* isolates using Mueller-Hinton (M-H) agar and via the use of Kirby-Bauer (K-B) method. The cultivation and biochemical tests resulted in 7 (53%) for cows, 5 (31%) for sheep, and 5 (55.5%) for goats) pure isolates of *Salmonella* spp. Antibiotic resistance was varied among the isolates with MAR index of 0.39 for cows, 0.32 for sheep, and 0.24 for goats. Most isolates were resistant to almost all antibiotics, especially streptomycin, ampicillin, kanamycin, and novobiocin. The current findings reveal that fecal samples from cows, sheep, and goats have *Salmonella* spp with antibiotic resistance traits.

Keywords: Antibiotic resistance, farm animals, *Salmonella*.

Introduction

As a serious public health risk, salmonella illnesses have been linked to gastro-intestinal health problems. Antibiotics are routinely employed to cure and manage food-animal related salmonellosis, leading to the rising frequency of antibiotic resistance rates in *Salmonella* that has attracted international concern (1). Despite the fact that antibiotic resistance has existed for a long time, the "resistome" is a growing concern. Antibiotic usage in animal industries and human health care facilities, lack of sanitation, the migration of wildlife animals, and an inadequate sewage system are major factors in the rise of the global resistome. In healthcare, antibiotics are one of the most often utilized treatments for infections. During the 'golden age' of antibiotics, which spanned the 1930s to 1960s, a large number of antibiotics were

developed. Antibiotic development slowed due to the emergence of drug-resistant organisms, and as a result, this period came to an end. The establishment of antibiotic resistance is predisposed by a lack of new antibiotics being developed or discovered, as well as overuse of existing antibiotics (2, 3, 4). As the problem of antimicrobial resistance grows, so does concerns for human, animal, and environmental well-being. Multidrug-resistant (MDR) bacteria, sometimes known as "superbugs," are to blame for this antimicrobial resistance. Five MDR bacteria reside in the animal, human, and environmental niches. Growing antibiotic usage in people and animals, antibiotics marketed over-the-counter, expanded worldwide travel, inadequate sanitary conditions and manure/feces discharge of



nonmetabolized antibiotics or their residue are all possible factors for "the global resistome" or antimicrobial resistance. These elements put the emergence of MDR pathogens in the population under genetic selection pressure (5,6). Latest information on the utilization of antibiotics in livestock has shown where the greatest concentration of antibiotic usage is taking place throughout the world, which impacts both economic sectors and public health. Cattle, chicken, and pigs are among the most widely antibiotic-given food animals, and antibiotic usage is expected to rise by up to 67% by 2030 in the globe's most inhabited nations (7). The aim of this work was to isolation of *Salmonella* from rectal fecal samples of farm animals and evaluate its antibiotic resistance capability in Al-Diwaniyah City, Iraq.

Materials and methods

Sampling

A total of 38 animals (13 cows, 16 sheep, and 9 goats) were employed to take samples via the use of sterile cotton swabs dipped in sterile saline solution before they were used to take rectal fecal samples and put them in containers, which pre-loaded with 5ml sterile saline solution. Then, these containers were icebox-transported within two hours to the Laboratory of Microbiology, College of Veterinary Medicine, University of Al-Qadisiyah, Al-Diwaniyah City, Iraq, for immediate Lab processing.

Salmonella cultivation and identification

The samples were cultivated using different agar media, such as 10ml-

Results

The cultivation and biochemical tests resulted in 17 (7 (53%) for cows, 5 (31%) for sheep, and 5 (5.5%) for goats) pure isolates of *Salmonella* spp. (Table 1 and 2). Antibiotic

RAPPAPORT-VASSILIADIS-Soya broth for enrichment (incubated for 24hrs at 37°C), MacConkey agar, Brilliant Green agar, Salmonella-Shigella agar, deoxycholate citrate agar, and Xylose Lysine Deoxycholate agar, which were streaked from the enrichment growth and incubated for 24hrs at 37°C. After that, the growth was sub-cultivated onto nutrient agar plates for 24hrs at 37°C. Then, the suspected *Salmonella* growth from the nutrient agar was subjected to biochemical identification using Triple Sugar Iron agar, Synthetic Complete agar, methyl red test, and indole test, which were performed by using methods from Coles (8,9).

Antibiotic susceptibility test

antibiotic susceptibility assay was performed on the pure *Salmonella* isolates using Mueller-Hinton (M-H) agar and via the use of Kirby-Bauer (K-B) method. The M-H plates were incubated for 24hrs at 35-37°C with the utilization of 13 antibacterial agents followed by calculating the multiple antimicrobial resistance (MAR) index (8,10). Determination of MAR index: Determination of MAR index followed the procedure described by (4), in which the number of antibiotics an isolate is resistant is divided by the total number of the antibiotics used in the study.

Statistical analysis:

For statistical analysis, a Chi square test (X^2) was performed to assess the significance, with the IBM Statistical Package for Social Sciences (SPSS), version 27. Values less than or equal to 0.05 were considered statistically significant.

resistance was varied among the isolates with MAR index of 0.39 for cows, 0.32 for sheep, and 0.24 for goats (Table 2 and 3 and Figure 1).

**Table 1: *Salmonella* occurrence in rectal fecal samples.**

Type of animals	Number of samples	Number of <i>Salmonella</i> positive samples	Percentage (%)
Cows	13	7	53
Sheep	16	5	31
Goat	9	5	55.5

Table 2: *Salmonella* spp. based biochemical features

Features	Outcomes
Capsule	-
Catalase	+
Citrate	-
Flagella	+
Gas	-
Gelatin hydrolysis	-
Gram staining	-
Urease	-
H ₂ S	+
Indole	-
Motility	+

Negative, +: Positive.

Table 3: Antimicrobial susceptibility test of *Salmonella* spp isolates from cows, sheep, and goats

No.	Isolate code	Antibiotic Sensitivity												
		Tet 30µg	ST 20 µg	Cho 30 µg	Bac 10 µg	Gen 10 µg	Pol 10 µg	Pen 10 µg	Met 5 µg	Amp 10 µg	Nit 200 µg	Kan 30 µg	Nor 5 µg	Van 30 µg
1.	SC4	RES	RES	RES	SEN	SEN	RES	RES	SEN	RES	RES	RES	RES	RES
2.	SC5	RES	RES	SEN	RES	SEN	SEN	RES	RES	RES	RES	RES	RES	RES
3.	SC8	SEN	RES	SEN	RES	RES	RES	SEN	SEN	RES	SEN	RES	RES	SEN
4.	SC10	SEN	RES	RES	SEN	RES	RES	SEN	RES	RES	SEN	RES	RES	SEN
5.	SC11	SEN	RES	SEN	SEN	RES	RES	SEN	RES	RES	SEN	RES	RES	SEN
6.	SC12	SEN	RES	SEN	RES	RES	RES	RES	RES	RES	RES	RES	RES	SEN
7.	SC13	RES	RES	SEN	RES	I	RES	RES	RES	RES	RES	RES	RES	RES
8.	SS1	SEN	RES	SEN	SEN	RES	RES	RES	SEN	RES	RES	RES	RES	RES
9.	SS3	RES	RES	SEN	RES	RES	RES	RES	SEN	RES	SEN	RES	RES	RES
10.	SS12	RES	RES	I	SEN	RES	RES	RES	RES	RES	SEN	RES	RES	RES
11.	SS14	RES	RES	RES	SEN	RES	RES	RES	SEN	RES	SEN	RES	RES	RES
12.	SS8	SEN	RES	RES	RES	RES	SEN	RES	SEN	RES	SEN	RES	RES	SEN
13.	SG3	SEN	RES	RES	RES	RES	SEN	RES	SEN	RES	SEN	RES	RES	SEN
14.	SG4	RES	RES	RES	RES	RES	RES	SEN	SEN	RES	SEN	RES	RES	RES
15.	SG5	RES	RES	SEN	RES	SEN	RES	SEN	RES	RES	SEN	RES	RES	RES
16.	SG7	RES	RES	SEN	RES	SEN	RES	RES	RES	RES	RES	RES	RES	RES
17.	SG9	SEN	RES	SEN	RES	RES	RES	RES	RES	RES	RES	RES	RES	RES



TE: Tetracycline TE, ST: Streptomycin, Cho: Chloramphenicol, Bac: Bacitracin, Gen: Gentamicin, Pol: Polymyxin-B, Pen: Penicillin-G, Met: Methicillin, Amp: Ampicillin, Nit: Nitrofurantoin, Kan: kanamycin, Nor: Novobiocin, and Van: vancomycin.




Figure 1: Multiple antimicrobial resistance (MAR) Index of antibiotic susceptibility test for *Salmonella* spp. isolates from cows, sheep, and

Discussion

Among foodborne pathogens, *Salmonella* species rank as the third highest cause of mortality from diarrheal infections in the whole globe. Food derived from animals is the principal way through which this disease is transmitted to humans (11). The current findings revealed that antibiotic resistance varied among the current study isolates with MAR index of 0.39 for cows, 0.32 for sheep, and 0.24 for goats, which agree with those by Ramatla *et al* (12), who mentioned that *Salmonella* spp. recovered from different animal and environmental niches showed antibiotic resistant in (92.0%), (89.3%), (77.4%), (72.6%), (67.4%), and (52.2%) of the isolates against sulphonamides, enrofloxacin and erythromycin, oxytetracycline, imipenem, tetracycline, and trimethoprim, respectively. Tetracyclines decrease protein production by adhering to the small ribosomal subunit and inhibiting aminoacyl-tRNA from adhering to the protein manufacturing machinery. Tetracycline resistance may be acquired by the complex synthesis of several systems such as

efflux pumps, enzyme inhibition, and mutations. Tetracyclines are the main frequently utilized or excessively utilized antibiotics in food animal industry. This is attributable to the truth that over-the-counter veterinary drugs are both reasonably priced and extensively accessible (13, 14, 16). Eagar *et al.* (17) reported that the most frequently used antibiotic agents in animals at 16.7% and 12.4% were tetracyclines and sulphonamides, respectively. The analysis by Ramatla *et al* (12) revealed that streptomycin was resisted by about 38% of *Salmonella* isolates. However, high rates of 80%, 95%, 65.5%, and 20.1% of resistance against streptomycin were detected from Saudi Arabia, Italy, and Egypt. Due to the fact that streptomycin is on the top-listed of the most utilized antibiotics in food animal industry, frequent rates of antibiotic resistance by *Salmonella* isolates can be highly recorded (16,18,19,20).



Conclusion

The current findings reveal that fecal samples from cows, sheep, and goats have *Salmonella* spp with antibiotic resistance traits. This problem should be faced with specific

strategies to decrease antibiotic resistance in farm animals via the use of controlled antibiotic utilization and only in critical needs to use these medicines.

References

- 1-Vinueza-Burgos C, Baquero M, Medina J, De Zutter L. Occurrence, genotypes and antimicrobial susceptibility of *Salmonella* collected from the broiler production chain within an integrated poultry company. *Int J Food Microbiol* [Internet]. 2019 Jun 16 [cited 2022 Mar 9];299(6):1–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/30933685/>
- 2-Nathan C, Cars O. Antibiotic resistance--problems, progress, and prospects. *N Engl J Med* [Internet]. 2014 Nov 6 [cited 2022 Mar 9];371(19):1761–3. Available from: <https://pubmed.ncbi.nlm.nih.gov/25271470/>
- 3-Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clin Microbiol Rev* [Internet]. 2011 Oct [cited 2022 Mar 9];24(4):718–33. Available from: <https://pubmed.ncbi.nlm.nih.gov/21976606/>
- 4-Singer AC, Shaw H, Rhodes V, Hart A. Review of Antimicrobial Resistance in the Environment and Its Relevance to Environmental Regulators. *Front Microbiol* [Internet]. 2016 Nov 1 [cited 2022 Mar 9];7(11):1728–1,749. Available from: <https://pubmed.ncbi.nlm.nih.gov/293508501/>
- 5-Davies J, Davies D. Origins and Evolution of Antibiotic Resistance. *Microbiol Mol Biol Rev* [Internet]. 2010 [cited 2022 Mar 9];74(3):417–33. Available from: <https://pubmed.ncbi.nlm.nih.gov/20937522/>
- 6-Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, et al. Global trends in antimicrobial use in food animals. *Proc Natl Acad Sci U S A* [Internet]. 2015 May 5 [cited 2022 Mar 9];112(18):5649–54. Available from: <https://pubmed.ncbi.nlm.nih.gov/26426470/>
- 7-Roca I, Akova M, Baquero F, Carlet J, Cavaleri M, Coenen S, et al. The global threat of antimicrobial resistance: science for intervention. *New Microbes New Infect* [Internet]. 2015 Jul 1 [cited 2022 Mar 9];6(4):22–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/26446399/>
- 8-Coles EH. *Veterinary clinical pathology*. 4th ed. W.B. Saunders; 1986. 490 p.
- 9-ISO (International Organization for Standardization). *Microbiology of food and animal feeding stuff-horizonal method for the detection of Salmonella spp*. Geneva, Switzerland; 2000. 511–525 p.
- 10-Osundiya O, Oladele R, Oduyebo O. Multiple Antibiotic Resistance (MAR) indices of *Pseudomonas* and *Klebsiella* species isolates in Lagos University Teaching Hospital. *African J Clin Exp Microbiol*. 2013 Aug 5;14(3):164–8.
- 11-Ferrari RG, Rosario DKA, Cunha-Neto A, Mano SB, Figueiredo EES, Conte-Juniora CA. Worldwide Epidemiology of *Salmonella* Serovars in Animal-Based Foods: a Meta-analysis. *Appl Environ Microbiol* [Internet]. 2019 Jul 1 [cited 2022 Mar 10];85(14):e00591–619. Available from: <https://pubmed.ncbi.nlm.nih.gov/36606869/>
- 12-Ramatla T, Tawana M, Onyiche TE, Lekota KE, Thekisoe O. Prevalence of antibiotic resistance in salmonella serotypes concurrently isolated from the environment, animals, and humans in South Africa: A systematic review and meta-analysis. *Antibiotics* [Internet]. 2021 Dec 1 [cited 2022 Mar 10];10(12):1435–48. Available from: <https://pubmed.ncbi.nlm.nih.gov/38698067/>
- 13-Hao H, Sander P, Iqbal Z, Wang Y, Cheng G, Yuan Z. The Risk of Some Veterinary Antimicrobial Agents on Public Health Associated with Antimicrobial Resistance and their Molecular Basis. *Front Microbiol* [Internet]. 2016 Oct 18 [cited 2022 Mar 10];7(10):1626–36. Available from: <https://pubmed.ncbi.nlm.nih.gov/275067539/>
- 14-Mokgophi TM, Gcebe N, Fasina F, Adesiyun AA. Antimicrobial Resistance Profiles of *Salmonella* Isolates on Chickens Processed and Retailed at Outlets of the Informal Market in Gauteng Province, South Africa. *Pathogens* [Internet]. 2021 Mar 1 [cited 2022 Mar 10];10(3):1–15. Available from: <https://pubmed.ncbi.nlm.nih.gov/38000370/>
- 15-Theobald S, Etter EMC, Gerber D, Abolnik C. Antimicrobial Resistance Trends in *Escherichia coli* in South African Poultry: 2009-2015. *Foodborne Pathog Dis* [Internet]. 2019 Sep 1 [cited 2022 Mar 10];16(9):652–60. Available from: <https://pubmed.ncbi.nlm.nih.gov/316965260/>



- <https://pubmed.ncbi.nlm.nih.gov/31099598/>
- 16-Vaez H, Ghanbari F, Sahebkar A, Khademi F. Antibiotic resistance profiles of Salmonella serotypes isolated from animals in Iran: a meta-analysis. *Iran J Vet Res* [Internet]. 2020 [cited 2022 Mar 10];21(3):188–97. Available from: [/labs/pmc/articles/PMC7608045/](https://pubmed.ncbi.nlm.nih.gov/31099598/)
- 17-Eagar H, Swan G, Van Vuuren M. A survey of antimicrobial usage in animals in South Africa with specific reference to food animals. *J S Afr Vet Assoc* [Internet]. 2012 [cited 2022 Mar 10];83(1):16–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/23327130/>
- 18-Velasquez CG, MacKlin KS, Kumar S, Bailey M, Ebner PE, Oliver HF, et al. Prevalence and antimicrobial resistance patterns of Salmonella isolated from poultry farms in southeastern United States. *Poult Sci* [Internet]. 2018 Jun 1 [cited 2022 Mar 10];97(6):2144–52. Available from: <https://pubmed.ncbi.nlm.nih.gov/29608757/>
- 19-Elhadi N. Prevalence and antimicrobial resistance of Salmonella spp. in raw retail frozen imported freshwater fish to Eastern Province of Saudi Arabia. *Asian Pac J Trop Biomed* [Internet]. 2014 [cited 2022 Mar 10];4(3):234–8. Available from: [/labs/pmc/articles/PMC3868795/](https://pubmed.ncbi.nlm.nih.gov/29608757/)
- 20-Wang X, Biswas S, Paudyal N, Pan H, Li X, Fang W, et al. Antibiotic Resistance in Salmonella Typhimurium Isolates Recovered From the Food Chain Through National Antimicrobial Resistance Monitoring System Between 1996 and 2016. *Front Microbiol* [Internet]. 2019 [cited 2022 Mar 10];10(5):985–96. Available from: [/labs/pmc/articles/PMC6514237/](https://pubmed.ncbi.nlm.nih.gov/29608757/)