



Research article

Comparison of different methods in the identification of *Staphylococcus aureus* in cattle

Abeer L. mohammed

Ali A. Al-Iedani

Department of Microbiology; College of Veterinary Medicine; University of Basrah, Basrah, Iraq.

Corresponding Author Email:

Co-author Email:

Abstract

This study aimed to compare the efficiency of conventional microbiological , Miniaturize kit GP24 and molecular technique by using PCR of nuc gene in identification of S. aureus. Also to evaluate efficiency of the tube coagulase and DNase activity tests. 104 of suspected S. aureus were isolated from different cattle samples, (68.3%) and (75%) were positive for coagulase and DNase tests, respectively. The sensitivity and specificity of coagulase and DNase were 92.5%, 75.7% and 79.1% , 32.4% respectively. The identification rate of S. aureus by GP24 kit and PCR detection using nuc gene were 68.26% and 64.42% respectively. Out of 104(31.7%) of isolates were identified as coagulase negative staphylococci by using GP24 kit. Antimicrobials susceptibility assays of 67 S. aureus isolates revealed that, the highest resistance rates were against penicillin (97%) followed by tetracycline (25.4%). While all isolates were found to be highly sensitive toward Imipenem, vancomycin and chloramphenicol (100%) for each. In conclusion the coagulase and conventional microbiological techniques have more reliable results in comparison with DNase test for identification of S. aureus. On other hand, the results of GP24 kit and coagulase tube test were similar in sensitivity and specificity.

Keywords: *S. aureus, GP24 kit, coagulase, DNase, nuc gene.*

Introduction

Staphylococcus aureus belongs to the family *Staphylococcaceae* which are spherical gram positive bacteria , approximately of 0.5 – 1.5 μm in diameter, which form irregular clusters resembling bunches of grapes. aerobic, facultative anaerobic, non-spore forming and non-motile (1).*S. aureus* is an

important etiologic agent producing infections in animals (2) and cause a wide range of infections including dermatitis, pneumonia, septicemia, osteomyelitis and meningitis as well as mastitis in cattle (3).

Routine identification and enumeration of S. aureus isolates are usually carried out by conventional



methods based on the use of selective media such as Baird-Parker and mannitol salt agar (MSA), followed by identification of suspicious colonies by biochemical reactions which include catalase positive and oxidase negative (4, 5) and coagulase positive, which is an enzyme secreted by *S. aureus* causing the clotting of plasma in the host (6, 7). Coagulase production is an important phenotypic determinant of *S. aureus* which is associated with virulence as it resists phagocytosis and helps bacteria in virulence (8). Tube coagulase test detects secreted extracellular free coagulase that reacts with a substance in plasma called "Coagulase-Reacting Factor" (CRF) to form a complex, which in turn reacts with fibrinogen to form fibrin (the clot) (9).

Seven species of coagulase-positive staphylococci (CoPS) have been identified including *Staphylococcus aureus*, *S. intermedius*, *S.*

schleiferi subsp. coagulans, *S. hyicus*, *S. lutrae*, *S. delphini*, and *S. pseudintermedius*. In addition to *S. aureus*, the other CoPS species can cause severe infections compared with those caused by coagulase-negative staphylococci (CoNS) (10).

S. aureus strains produce an extracellular thermostable nuclease protein with a molecular mass of 17,000 Da. encoded by *nuc* gene. It is an endonuclease, degrading both DNA and RNA (86R), this gene is a specific marker gene that can be used for distinguishing *S. aureus* from other staphylococcus spp. (11, 12).

This study aimed to compare the efficiency of conventional microbiological, Miniaturize kit GP24 and molecular technique (PCR of *nuc* gene) in identification of *S. aureus*. Also to evaluate the efficiency of the tube coagulase and DNase activity tests.

Materials and Methods

Samples collection: A total of 212 samples from cattle including ; 42 mastitis milk samples detected by using California mastitis test (CMT) according to (13), the study was conducting during the period from January to June in Basrah (2018) as shown in table 1.

Table (1): Number and types of collected samples

Samples	Number
Mastitic milk (detected by CMT)	42
Raw milk from local market	70
Nasal swabs	46
Udder surface swabs	32
Coat Swabs	22
Total	212

Conventional microbiological technique:

All samples were subjected to colony morphology characteristics on selective media (MSA), Gram's staining, conventional biochemical tests (catalase, oxidase, DNase and coagulase) according to (14).

Miniaturize kit GP24:

The suspected isolates were submitted to GP24 (Slovak, slovakia) a ready-to-use test consisting of 24 biochemical tests to which an homogenous bacterial suspension (100µl) at 3McFarland turbidity is added. Urea (URE) and Arginine (ARG) wells (H1 and H2 wells) were covered with 2 – 3 drops of paraffin oil. The plate was incubated at



37°C for 24 h. Result of identification was obtained by identification table and identification online software (DIAGNOSTICS s. r. o.)

Molecular detection:

Genomic DNA of suspected *S. aureus* isolates was extracted by DNA kit (Geneaid, USA) and according to manufacturer's protocol. *S. aureus* isolates were confirmed by PCR detection using *nuc* gene(423bp) primer *nuc*

F: 5'- GCT TGC TAT GAT TGT GGT AGC C 3' and

R: 5'- TCT CTA GCA AGT CCC TTT TCC A 3' according to (15).

Determination of the antimicrobials susceptibility of isolates:

The confirmed *S. aureus* isolates were subjected to antimicrobial susceptibility test. Twelve antimicrobials from (Bioanalyse/ Turkey) were used to detect the isolates susceptibilities using disk diffusion method according to (16) including, Cefoxitin (30 µg), Ceftriaxone (30 µg), Chloramphenicol (30 µg), Ciprofloxacin (5 µg), Erythromycin (15 µg), Gentamycin (10 µg), Imipenem (10µg), Oxacillin (1 µg), Penicillin (10 µg), Tetracycline(30 µg), Trimethoprim(5 µg) and Vancomycin (30 µg). Cefoxitin was used as a surrogate for study of methicillin resistance.

Statistical analyses

The data were statistically using analyzed the software package IBM SPSS

22. Next a 2 × 2 contingency table was used for diagnostic specificity and sensitivity . Diagnostic sensitivities and specificities were calculated as follows (17):

$$\text{Sensitivity} = TP / (TP + FN)$$

$$\text{Specificity} = TN / (TN + FP)$$

*TP :true positive, FP: false positive, TN: true negative, FN: false negative

Results

Out of 212 tested samples 104 (49.1%) isolates were considered as suspected *S. aureus* by using conventional microbiological techniques. Out of 104 suspected isolates, 71(68.3%) and 78(75%) were positive for coagulase and DNase tests, respectively, as shown in table (2) and figure (1). The sensitivity and specificity of DNase were 79.1% and 32.4% respectively. Whereas, for coagulase were 92.5% and 75.7% respectively. However the sensitivity and specificity of coagulase and DNase tests were calculated in comparison with detection of *nuc* gene (table 2).

Table (2) Number of isolates which identified by using different diagnostic methods

Sample tested	No. of Isolates identified by *	Dnase test (%)	Coagulase (%)	<i>nuc</i> gene (%)
212	104	(75%)	(68.3%)	(64.4%)



* Culturing on MSA, Gram stained smear and biochemical tests (Oxidase and catalase tests).

Sensitivity and specificity of coagulase test= 92.5%, 75.7%.

Sensitivity and specificity of Dnase test=79.1%, 32.4%.




Figure (1) A: Tube Coagulase Test, P: coagulase positive (clot), N: coagulase negative (not clot); B:DNase test, P: positive result and N: negative result

All suspected isolates (104) were subjected to identification using GP24 kit figure (2) and PCR detection using *nuc* gene figure (3). The identification rate of *S. aureus* was 68.26% and 64.42% respectively table (3). By using molecular detection, the results showed that, the high percentage of *S. aureus* isolates were from udder surface samples (93.33%), followed by subclinical

mastitis (64.28%), whereas the lowest percentage was observed from the coat swabs (55.55%). The sensitivity and specificity of GP24 kit were 92.5% and 75.7% respectively. The sensitivity and specificity of GP24 kit to identify *S. aureus* was calculated in comparison with detection of *nuc* gene

Table(3): Comparison between conventional microbiological techniques, GP24 kit and molecular technique in identification of *S. aureus* from different samples sources.


Samples	No. of <i>S. aureus</i> using		
	conventional microbiological technique	GP24 kit	<i>nuc</i> gene
Subclinical mastitis	28	(64.28%)	(64.28%)
Raw milk from local markets	37	(72.97%)	(56.75%)
Nasal swabs	15	(66.66%)	(60.00%)



Udder surface swabs	15	(73.33%)	(93.33%)
Coat swabs	9	(55.55%)	(55.55%)
Total	104	(68.26%)	(64.42%)

Sensitivity and specificity of GP24 kit = 92.5%, 75.7%.

*The ratios were calculated by dividing the number of cell on number of conventional microbiological technique cell.



Taxa	Identification (%)	Differentiation (T)
<i>Staphylococcus aureus</i> ssp. <i>aureus</i>	excellent 100.00	excellent 1.00

CAT	URE	MLT	SOR	LAC	FRU	ARA	RAF	bGA		
+	+	+	-	+	+	-	-	+		
OK	OK	OK	OK	OK	OK	OK	OK	OK		
COA	NOV	ARG	MAN	TRE	CEL	MNS	RIB	MLZ	GLR	PHS
+	0	+	+	+	-	+	-	-	-	0
OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	OK
bHEM	NIT	ESL	MLB	SUC	GAL	XYL	NAG	bGL	HIP	
0	+	-	-	+	+	-	+	+	0	
OK	OK	OK	OK	OK	OK	OK	OK	OK	OK	
OXI	VP	PYR	LAP	BAC	YEP	AHE				
-	0	0	0	0	0	0				
OK										

Figure (2) A: The plate of GP24 kit
(The plates designed for 4 isolates)

B: Online identification table using GP24 kit




Fig (3): Electropherogram of *nuc* gene amplification

The product size (423) bp, the mixture was run in 1.5 % agarose gel, stained with ethidium bromide, L: Ladder.

The 104 isolates (68.26%) were identified as *S. aureus* and the rest (31.7%) were identified as the other *Staphylococcus* spp by using miniaturize GP24, (30.3%),

(27.3%), (27.3%), (12.1%) and (3.0%) of isolates were identified as *S. chromogenes*, *S. hominis*, *S. xylosus*, *S. capitis* and *S. vitulinus* respectively (table 4).

Table (4): Number and percentages of other *Staphylococcal* spp. identified by using GP24 kit.

<i>Staphylococcus</i> spp.	mastitis	Raw	Udder	Nasal	Coat	Total
----------------------------	----------	-----	-------	-------	------	-------



	isolates (%)	milk of market isolates (%)	isolates (%)	isolates (%)	isolates (%)	%
<i>S. hominis</i>	3	3	3	0	0	27.3%
<i>S. xylosus</i>	4	2	1	2	0	27.3%
<i>S. chromogenes</i>	4	3	0	2	1	30.3%
<i>S. vitulinus</i>	0	1	0	0	0	3.0%
<i>S. capitis</i>	0	0	0	2	2	12.1%
Total	33.3%	27.3%	12.1%	18.2%	9.1%	33

* The ratios were calculated by dividing the number of cell on total number cell

The results showed that, the highest resistance was against penicillin (97%) followed by tetracycline (25.4%), Ceftriaxone (20.9%) and Ciprofloxacin (18%). While all isolates were found to be highly sensitive toward Imipenem (100%), vancomycin (100%) and chloramphenicol

(100%). Out of 67 isolates 44(65.7%) were resistant to Cefoxitin, whereas, 48(71.64%) were resistant to oxacillin, Cefoxitin were used to detect the phenotype of methicillin resistant *Staphylococcus aureus* (MRSA). Table (5)

Table (5): Antimicrobial susceptibility of *S. aureus* isolates from

Antimicrobial agents	Sensitive	Intermediate	Resistance
Ceftriaxone	(52.2%)	(26.9%)	(20.9%)
Chloramphenicol	(100%)	(0%)	(0%)
Ciprofloxacin	(71.6%)	(10.4%)	(18%)
Erythromycin	(56.8%)	(32.8%)	(10.4%)
Gentamycin	(80.6%)	(11.9%)	(7.5%)
Imipenem	(100%)	(0%)	(0%)
Penicillin	(3%)	-	(97%)
Tetracycline	(40.3%)	(34.3%)	(25.4%)
Trimethoprim	(55.2%)	(28.4%)	(16.4%)
Vancomycin	(100%)	(0%)	(0%)



Cefoxitin	(34.3%)	-	(65.7%)
Oxacillin	(22.4%)	(5.9%)	(71.64%)

Discussion

The initial identification of *S. aureus* depended on colonial appearance on the conventional selective media MSA, their biochemical profile and confirmed by molecular method (PCR) (18). In this study the suspected isolates using conventional microbiological techniques were (49.1%), this result was compatible with (19) who reported that (40.81%) of samples were carried *S. aureus*. Whereas this result was higher than (20), who revealed that *S. aureus* was isolated from (37%) raw milk of cow in Basrah, and lower than (19) who detected 85.7% of suspected *S. aureus* by using conventional microbiological techniques.

Out of 104 isolates (68.3%) were coagulase positive, The current result is compatible with the studies of (21, 22) who reported coagulase production in percentages 78.5% and 77.8% respectively. While, higher percentage 89% was recorded by (18). In contrast, a much lower percentages were reported by the studies of (23, 24) 28.07% and 16.67%, respectively. Sensitivity and specificity of coagulase test were evaluated by comparing with *nuc* gene, the results were 92.5% and 75.7% respectively. Similar results were obtained by (18) who revealed that sensitivity of tube coagulase test was more than 91% and (25) who reporting a sensitivity of 97% but a specificity of 100%. Higher percent of coagulase positive isolates comparing to *nuc* gene result may be due to present other coagulase positive staphylococcus spp.

The DNase and thermostable nuclease activity (TNase) tests are being used for presumptive identification and appear to be a consistent property of *S. aureus* (26). In present investigation

78(75%) showed DNase activity on the DNase agar, this result is in agreement with (18) who reported 75% of cultures were positive for DNase test, however, this result is lower than (21) who found DNase activity in 86.93%, and higher than (23) who reported (65.7%). On the other hand DNase test had a sensitivity of 79.1% and specificity 32.4%. congruous, in result of sensitivity was obtained by (18, 27) who reported 75% and a 96%. Concerning the specificity, the present result is consistent with (28) who reported the sensitivity and specificity of DNase as 53.13% and 41.84% respectively.

Seventy one out of 104 isolates (68.3%) were identified by Miniaturize kit GP24 as *S. aureus* and the rest (31.7%) of isolates were identified as coagulase negative staphylococci (CoNS). Higher occurrence rates of CoNS were observed in both mastitic and raw milk 11(33.3%) and 9(27.3%) respectively, similar result was obtained by (29 and 30) who recorded the isolation rates of CoNS from bovine subclinical mastitis ranged from 7.4% to 53.5%. CoNS are normal flora of the skin of the teat and common in subclinical mastitis (31). This result was higher in compared with the study of (32) who reported the isolation rate of CoNS from subclinical mastitis was 15.55%. The sensitivity of GP24 kit was high 92.5%, and specificity was 75.7%. On other hand this results are similar to result of coagulase test.

Concerning the molecular detection of *S. aureus* by pcr technique, targeting of *nuc* gene encodes for a specific thermostable nuclease (33). This gene were used to detect *S. aureus* in previous studies (34, 35, 36). In present study 67 (64.42%) of isolates were identified as *S. aureus*,



This result is in line with many studies such as (37, 38) In contrast, a much lower occurrence rates of *S. aureus* were found by (39, 40, 41).The highest rates of *S.aureus* were observed in udder surface and nasal swabs (93.33% and 60%, respectively),this result explains the causes of mastitis may be due to smudge the teats with dung and mud, also the direct source of bacteria of milk.While the causes which increase this microorganism in nares may be due to warm, moist and nutrients.

The antimicrobial susceptibility test profile obtained in the present study showed that a significant percentage of *S. aureus* isolates were resistant to most of the commonly used antibiotics such as penicillin, Oxacillin and Cefoxitin. The cefoxitin diffusion method was used as an alternative marker for the detection of methicillin resistance (16), the resistance of *Staphylococcus* spp. to penicillins is a well-known phenomenon worldwide, and 10 to 70 % of *S. aureus* in cows may be resistant to these drugs, depending on geographical location (42). This result compatible with local study (43). However(44) found that resistant to penicillin was (76.92%) followed by gentamicin(51.28%), oxacillin (38.46 %), tetracyclin (28.21%) and erythromycin (23.08%) in *Staphylococcus* species from mastitic milk.

On other hands, all the *S.aureus* isolates were sensitive to vancomycin chloramphenicol and Imipenem (100%), the present study are in line with studies (45, 46) .Many factors may have contributed to the above level of resistance toward the tested antibacterial drugs. Three major factors determine this crisis, the increasing frequency of antimicrobial resistance phenotypes among microbes is an evolutionary response to the widespread use of antimicrobials, the resistance of bacteria may be resulted from arbitrary uses of antimicrobial and the extensive and often unnecessary use of antimicrobials by humanity provides the strong selective

pressure that is driving the evolutionary response in the microbial world (47).

In conclusion, the coagulase plus conventional microbiological techniques have more reliable results in comparison with DNase test and can be used for initial identification of *S. aureus*. However, the DNase test plus conventional microbiological techniques may use as screening test for primary isolation of *S. aureus*. Concerning GP24 kit, the results were correlated with coagulase test because this kit depend on adding the result of coagulase to identify the gram positive bacteria, moreover, the sensitivity and specificity of GP24 similar to that of coagulase test . The number of *S. aureus* isolates which identified using GP24 kit, coagulase or DNase were more than that identified by using *nuc* gene, these results were due to the presence of other coagulase positive staphylococci which have phenotypes similar to *S. aureus*.

References

- 1- Markey, B K, Leonard, F C, Archambault, M., Cullinane, A. and Maguire, D. Clinical veterinary microbiology. (2013) 2nd . *Staphylococcus* spp. 105- 120
- 2- Eslampour MA, Hovareshti P, Feizabadi MM. Molecular characterization of *Staphylococcus aureus* isolated from bovine mastitis in Iran: Vet Microbiol. 2009; 39: 207-8.
- 3- Hasman, H., Moodley, A. and Guardabassi, L, Spa type distribution in *Staphylococcus aureus* originating from pigs, cattle and poultry: *Vet. Microbiol.* .(2010).141:326–331.
- 4- Amagliani, G. Glammarini, C. Omicciolo, E. Brandi, G .Magnami, Food Control: (2007) 18 :1137–1142.
- 5- Sommer, L. Y. Kashi, J. Food Prot. .(2003). 66 : 1658–1665.
- 6- Ryan KJ, Ray CG. Sherris Medical microbiology: An Introduction to Infectious Diseases. (2004) 4th edn. McGraw Hill Publishers.



- 7- Windria, S., Widianingrum, D.C. and Salasia, S.I.O. Identification of *Staphylococcus aureus* and coagulase-negative staphylococci isolates from mastitis milk of Etawa crossbred goat. *Res. J. Microbiol.*, (2016) 11: 11-19
- 8- Zhang K, Sparling J, Chow BL, Elsayed S, Hussain Z, Church DL. New quadruplex PCR assay for detection of methicillin and mupirocin resistance and simultaneous discrimination of *Staphylococcus aureus* from coagulase-negative staphylococci. *J Clin Microb.* (2004). 42(11): 4947-55.
- 9- Aldhar, C.N., Chilgunde, S.N., Gaikwad, R.V., Smad, A. and Roy, S. Current trends in diagnosis of mastitis in Veterinary practice. *The J. Remt. Vet. Corp.* (2005) 44(3): 133-140
- 10- Quinn, P J, Markey, B K, Leonard, F C, Fitzpatrick, E S, Fanning, S and Hartigan, P J. Veterinary microbiology and microbial disease: *Staphylococcus* species. (2011) 2nd Wiley-Blackwell 179 -187
- 11- Tucker, P. W., E. E. Hazen, and F. A. Cotton. Staphylococcal nuclease reviewed: a prototypic study in contemporary enzymology. I. Isolation, physical and enzymatic properties. *Mol. Cell. Biochem.* 1978. 22:67-77.
- 12- Zhang K, Sparling J, Chow BL, Elsayed S, Hussain Z, Church DL. New quadruplex PCR assay for detection of methicillin and mupirocin resistance and simultaneous discrimination of *Staphylococcus aureus* from coagulase-negative staphylococci. *J Clin Microb.* (2004). 42(11): 4947-55.
- 13- Aldhar, C.N., Chilgunde, S.N., Gaikwad, R.V., Smad, A. and Roy, S. Current trends in diagnosis of mastitis in Veterinary practice. *The J. Remt. Vet. Corp.* (2005) 44(3): 133-140
- 14- Aclafaddin, J. F. Biochemical tests for identification of medical bacteria. (2000). 3rd Ed. Lippincott Williams and Wilkins USA.
- 15- Ongboot W., Chomvarin C., Engchanil C. and Chaimanee P. Multiplex PCR for detection of superantigenic toxin genes in methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* isolated from patients and carriers of a hospital in northeast Thailand. *Southeast Asian J. Trop. Med. Public Health* (2013). 44: 660-671.
- 16- National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Disk Susceptibility Tests—Eighth Edition; Approved Standard M2-A8. (2013). NCCLS, Wayne, PA, USA
- 17- Atam Nahavandi K. Calculating Sensitivity, Specificity and Predictive Values for Q Medical Diagnostic Tests, *Gene Cell Tissue.* 2018 ; 5(2):e80270. doi: 10.5812/gct.80270.
- 18- Ateete1, D.P, Kimani, C.N, Katabazi, F.A., Okeng, A, Okee, M. S, Ann Nanteza, A, Joloba, M. L and Najjuka, F.C. Identification of *Staphylococcus aureus*: DNase and Mannitol salt agar improve the efficiency of the tube coagulase test. *Annals of Clinical Microbiology and Antimicrobials*(2010). 9:23.
- 19- Boud, W.A., Molecular Detection of Vancomycin and Methicillin-Resistant Genes of *Staphylococcus aureus* Isolated from Animal and Human Specimens. (2019). M.Sc. Thesis, College of Veterinary Medicine, University of Basrah
- 20- Ahdi, A, SH .Molecular Detection of Superantigens in *Staphylococcus aureus* Isolated from Human ,Cattle and Buffalo Milk Samples. (2019). M.Sc. Thesis, College of Veterinary Medicine, University of Basrah



- 21- Singh, R.S.; Kumar, R. and Yadav, B.R. Distribution of pathogenic factors in *Staphylococcus aureus* strains isolated from intra-mammary infections in cattle and buffaloes. Indian J. Biotechnol. (2011). 10: 410-416.
- 22- urkyilmaz, S. and Kaya, O. Determination of some Virulence factors in *Staphylococcus* spp. isolated from various clinical samples. Turk. J. Vet. Anim. Sci. (2005). 30: 127-132.
- 23- Avid, F., Taku, A., Altaf Bhat, M., Badroo, G. A., Mudasir, M. and Ahmad Sofi, T. Molecular typing of *Staphylococcus aureus* based on coagulase gene. Veterinary World, EISSN: (2018). 2231-0916
- 24- Al-Iedani, A.A. and Jumaa, M. A. Detection of different *Staphylococcus* spp. on outer surfaces of farm animals, their owners and study their genotypic and phenotypic abilities to resist methicillin. Int. J. Agric. Sc & Vet. Med. (2017). 5 (2): 120-131
- 25- Davis, T. E., Fuller, D. D. and Aeschleman, E. C. Rapid, direct identification of *Staphylococcus aureus* and *Streptococcus pneumoniae* from blood cultures using commercial immunological kits and modified conventional tests. Diagn. Microbiol. Infect. Dis. (1992). 15:295-300.
- 26- Yadav, R., Sharma, S. K., Yadav, J., Nathawar, P and Kataria, A.K. Phenotypic and Genotypic Characterization of *Staphylococcus aureus* of Mastitic Milk Origin from Cattle and Buffalo for some Virulence Properties. J PURE APPL
- 27- Lagace-Wiens PRS, Alfa MJ, Manickam K, Karlowsky JA. Thermostable DNase Is Superior to Tube Coagulase for Direct Detection of *Staphylococcus aureus* in Positive Blood Cultures. J Clin Microbiol, (2007) 45(10):3478-3479.
- 28- Pumipuntu, N, Kulpeanpravit, S, Santajit, S, Tunyong, W, Thida Kongngoen, T., Hinthong, W and Indrawattana, N. Screening method for *Staphylococcus aureus* identification in subclinical bovine mastitis from dairy farms. Veterinary World, EISSN: (2017). (10) P: 2231-0916
- 29- Giannechini R, Conch, C., Rivero, R., Delucci, I. and Moreno Lopez, J. Occurrence of Clinical and Subclinical Mastitis in Dairy Herds in the West Littoral Region in Uruguay. Acta Vet. Scand., (2002) 43:221-230
- 30- Myllys, V., Asplund, K., Brofeldt, E., Hirvela-Koski V, Honkanen-Buzalasko T, Junttila, J., Kulkas, L., Myllykangas, O., Nikanen, M., Saloniemi, H., Sandholm, M., and Saranpää T. Bovine mastitis in Finland in 1988 and 1995-changes in prevalence and antimicrobial resistance. Acta Vet. Scand. (1998) 39:119-126
- 31- Pyorala S and Taponen, S. Coagulase-negative staphylococci emerging mastitis pathogens. Vet Microbiol., (2009) 134:3-8.
- 32- AL-Edany, A, Khudor, M and AL-Mousawi, Kh. Comparison of three indirect tests for the diagnosis of bovine subclinical mastitis caused by coagulase negative staphylococci with their susceptibility to seven antibiotics. Bas. J. Vet. Res. (2012). Vol.11, No.1
- 33- Foster TJ. Immune evasion by staphylococci. Nat Rev. Microbiol (2005) 3: 948-958.
- 34- Palomares, C., Torres, M.J., Torres, A., Aznar, J. and Palomares, J.C. Rapid detection and identification of *Staphylococcus aureus* from blood culture specimens using real-time fluorescence PCR, Diagnostic Microbiology and Infectious Disease, (2003) 45, 183-189.
- 35- Costa, A.M., Kay, I. and Palladino, S. Rapid detection of *mecA* and *nuc* genes in staphylococci by real-time multiplex



polymerase chain reaction, Diagnostic Microbiology and Infectious Disease, (2005)51, 13–17.

36- Asfour, H.A.E. and Darwish, S.F. Phenotypic and genotypic detection of both *mecA* and *blaZ* genes mediated β -lactam resistance in *Staphylococcus* strains isolated from bovine mastitis, Global Veterinaria, (2011) 6, 39–50.

37- Hammad, A.M. Microbiological studies on raw milk and some dairy products (2004). M. V. Sc. Thesis, Minufiya Univ., Egypt.

38- Rall, V. L. M.; Vieira, F. P.; Rall, R.; Vieitis, R. L.; Fernandes, A.; Candeias, J. M. G.; Cardoso, K. F. G. and Araujo, J. P. PCR detection of staphylococcal enterotoxin genes in *Staphylococcus aureus* strains isolated from raw and pasteurized milk. Vet. Microbiol. (2008) 132(3):408-413

39- Jahan, M. ; Rahman, M. ; Parvej, M. ; Chowdhury, S. ; Haque, M. ; Talukder, M . and Ahmed, S. Isolation and characterization of *S.aureus* from raw cow milk in Bangladesh. *Journal of Advanced Veterinary and Animal Research*, (2015) 2(1), 49-59.

40- Liu, H ; Li, S ; Meng, L ; Dong, L ; Zhao, S ; Lan, X and Zheng, N. Prevalence, antimicrobial susceptibility, and molecular characterization of *S.aureus* isolated from dairy herds in northern China. *Journal of Dairy Science*, (2017) 40 (5),1–8.

41- Momtaz, H. ; Tajbakhsh, E. ; Rahimi, E . and Momeni, M. Coagulase gene polymorphism of *S.aureus* isolated from clinical and sub-clinical bovine mastitis in Isfahan and Chaharmahal va Bakhtiari provinces of Iran. *Comparative Clinical Pathology*, (2011) 20(5), 519–522.

42- Vintov, J., Aarestrup, F.M., Zinn, C.E. and Olsen, J.E., Association between phage types and antimicrobial resistance among bovine *Staphylococcus aureus* from 10 countries, Veterinary Microbiology, (2003) 95, 133–147.

43- Idbeis, H.I. Molecular Analysis of *Staphylococcus aureus* Isolated From Mastitis and Study Their Biofilm Activity against Cinnamon Oil. (2019) PhD thesis. College of Veterinary Medicine, University of Basrah

44- Alian, F., Rahimi, E., Shakerian, A., Momtaz, H., Riahi, M. and Momeni, M. Antimicrobial resistance of *Staphylococcus aureus* isolated from bovine, sheep and goat raw milk, Global Veterinaria, (2012) 8, 111–114.

45- Al-Iedani, A. A. Phenotypic study on the capacity of biofilm production in *S.aureus* isolated from bovine subclinical mastitis and their impact on resistance to antimicrobials. Basrah Journal of Veterinary Research, . (2016) 15(2): 111–127.

46- Aliyu, Y. ; Reuben, C. ; Sani, A . and Salawu, E. Occurrence and antibiogram of *S.aureus* isolated from locally-pasteurised cow milk (kindirmo) sold in parts of Nasarawa Town, Nasarawa State, Nigeria. *Microbiology Research Journal International*, (2018) 23(4), 1–11.

47- Michael, C.A, Domine-Howes, D and Labbate, M. The antimicrobial resistance crisis: causes, consequences, and management. (2014) Vol. (2) www.frontiersin.org.