

Reasons for Preoperative Anxiety and Severity in Psychological Stress of Women Before Caesarean Section

Hadeel Fadel Farhoud¹, Hayder H. AL-Hadrawi².

Department of Mental Health Nursing, College of Nursing, University of Kufa, Najaf, Iraq. Email: hadeelfadel03@gmail.com
Department of Mental Health Nursing Department, College of Nursing University of Kufa, Najaf, Iraq. Email: hayderh.alhadrawi@uokufa.edu.iq

Article information	Abstract
<p>Article history: Received on April 20, 2023 Accepted on July 03, 2023 Available online January 12, 2024</p>	<p>Background: Preoperative fear is a common and neutral response of human beings as expecting surgery even if the surgery is significantly needed. There are several factors that cause preoperative stress or anxiety, which affect the physical and psychological health of people.</p> <p>Study aim: This study aims to measure the possible reasons for preoperative anxiety and the severity of psychological stress of women prior to caesarean section.</p> <p>Methods: The descriptive and comparative study design was used in a convenience sample of 120 pregnant women before caesarean section. The women were selected from the Al-Zahra Teaching Hospital in the province of Al-Najaf. The psychological stress level was measured using a brief measure of the emotional preoperative stress scale.</p> <p>Results: The study findings show that the level of psychological stress of women before caesarean delivery ranged from mild (56%) to moderate (43%). The study highlighted the importance of considering the increase in psychological stress levels for women prior to caesarean section and the need for more research into interventions that can reduce psychological stress.</p> <p>Conclusions and recommendations: Psychological stress during pregnancy negatively affects the health of women and their fetus. Many antianxiety medications are contraindicated for pregnant women; therefore, it is recommended that people use a stress management intervention before any type of surgery to reduce surgery-related psychological stress.</p>
<p>Keywords: preoperative fear, psychological stress, cesarean section</p>	
<p>Correspondence: Hadeel Fadel Farhoud hadeelfadel03@gmail.com</p>	

DOI: [10.33899/mjn.2024.182198](https://doi.org/10.33899/mjn.2024.182198), Authors, 2024, College of Nursing, University of Mosul.
This is an open-access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

INTRODUCTION

Cesarean section is a life-saving procedure that should be available to all women who require it for certain complications during pregnancy and childbirth and includes additional risks for hysterectomy, abnormal placentation, uterine rupture, stillbirth, and preterm birth in the next pregnancy. With more caesarean sections, there is a greater incidence of bleeding, the need for blood transfusion, pelvic adhesions,

intraoperative surgical damage, and hysterectomy (Sandall et al., 2018). Studies have shown that caesarean section has a psychological effect on the mother after caesarean section, leading to an increased risk of depression, increased significant psychological trauma, and posttraumatic stress related to birth (Vanahor et al., 2018)². Psychological stress is a common side effect of

expecting surgery even if surgery is significantly needed. However, the response to stress is influenced by changes in the metabolic, neurohormonal, and immunological systems, as well as anxiety. Thus, these alterations have been observed as a result of surgery and stress has been associated with elevated cortisol levels in the blood (Leardi et al., 2007). Preoperative fear increases the risk during surgical procedures, as well as postoperative risks by adding a negative emotional response that causes depression or anxiety that affects the mother's recovery or delays the healing of the mother after the operation (Bakr et al. 2014). Postoperative physiological and psychological outcomes are influenced by preoperative psychological stress; therefore, the impact of psychological stress on women's health outcomes has been assessed and shown to impact the process of wound healing and postoperative problems in the first month after surgery (Levett & Grimmett, 2019). Numerous variables, including gender, marital status, fear of complications, apprehensions, previous surgical experience, socioeconomic status, postoperative pain, type of anaesthesia, lack of preoperative information, smoking and drinking, education and occupation have been related to the severity of preoperative stress. As a result, the need for more information on anaesthesia and surgical procedures is essential to help reduce stress prior to surgery (Kiriakopoulos et al., 2019; Jiwanmall et al., 2020). Pregnancy-specific stress has a stronger influence on birth outcomes than general stress and that stress is magnified in high-risk pregnancies. In particular, there is a significant increase in the risk of anxiety and stress in pregnant hospitalized patients who are given bed rest (Hobel et al., 2008; Teckenberg-Jansson et al., 2019; Kareem & Ali, 2021).

METHODS AND MATERIALS

Study design: The descriptive-comparative study design is used to measure the effects of possible reasons for preoperative anxiety and the severity of psychological stress of women prior to caesarean section.

Ethical Considerations: Before starting the sample collection, the study was approved by the Research Ethics Committee of the University of Kufa, Faculty of Medicine, and that study was free of any possible harm to the participants (physical or psychological). The women were informed that their participation in

the study was voluntary, and their information would be kept confidential.

Study Setting:

The study is carried out at Al-Najaf Al-Ashraf Government, Al-Zahraa Teaching Hospital for Delivery and Children. Al-Zahraa Hospital was chosen because of its preoperative admission policy, which requires admission for one day before cesarean delivery.

Sample and sampling plan:

A nonprobability "convenience" sampling technique was used to gather information from 120 pregnant women prior to the cesarean section. Cohen's methods were used to determine the appropriate sample size.

Inclusion and Exclusion Criteria: The inclusion criteria are women of all ages who are admitted for elective cesarean section. Women who are admitted at least one day before the cesarean section. Women are excluded if they are admitted for emergency cesarean delivery and those with a history of mental health problems.

Study Instrument:

The brief measure of the emotional preoperative stress scale was used to measure the psychological stress of women. This scale was originally developed by (Caumo et al., 2016) and it contains 15 items. The higher score indicates a severe level of preoperative psychological stress.

Data Collection:

Data were collected using a structured questionnaire. The translated version of the brief measure of the emotional preoperative stress scale was used to collect data from pregnant women. It was a self-administered process, and the estimated time to complete the questionnaire ranged between 20 and 25 minutes.

Statistical analysis: Microsoft Excel (2010) and SPSS software version 26 were used to analyze the collected data. Descriptive statistics were used to describe demographic data, health-related variables and psychological stress levels of pregnant women.

Results

The sociodemographic variables of 120 pregnant women are described in Table 1. The age ranges between (22–25) years (31.7%). Women with

primary education (35.0%); those not working (86.7%); Women living in urban areas constitute (74.2%). Figure 1 describes the various reasons for preoperative anxiety. Fear of death, concern for family, fear of postoperative pain, fear of anaesthesia, fear of medical errors, complications, and fear of apparent cause are the most common reasons women as causes for their preoperative anxiety. The

psychological stress levels are explained in Figure 2. Most pregnant women experience psychological stress ranging from mild (43%) to moderate levels (56%). Table 2 shows that of 17 reasons for preoperative fear, only 10 reasons contribute significantly to the psychological stress before caesarean delivery ($P = < 0.05$).

Table 1: Descriptive statistics of sociodemographic variables of women

Age Groups	f	%	Residence	f	%
17 - 21	20	16.7	Rural Area	31	25.8
22 - 25	38	31.7	Urban Area	89	74.2
26 - 30	32	26.7	Total	120	100.0
31 - 34	19	15.8	Marital Status	f	%
35 - 40	11	9.2	Married	118	98.3
Total	120	100.0	Separated	2	1.7
			Total	120	100.0
Educational Level	f	%	Working Status	f	%
Do not Read and Write	24	20.0	Not Working	104	86.7
Primary Education	42	35.0	Free Job	3	2.5
Intermediate School	28	23.3	Governmental Employee	13	10.8
High School	8	6.7	Total	120	100.0
Diploma or Bachelor Degree	18	15.0			
Total	120	100.0			





Table 2: Differences in psychological stress of women based on reasons of preoperative anxiety.

Reason for Preoperative Anxiety		ANOVA Test		
		Mean Square	F	Sig.
Reason 1	Fear of fluids or intravenous treatment	4.876	.222	.639
Reason 2	Fear of medical equipment	373.373	19.781	.0001
Reason 3	Fear of the unexpected results of the operation	103.754	4.903	.029
Reason 4	Post-operative pain	45.511	2.102	.150
Reason 5	Fear of anesthesia	183.755	8.971	.003
Reason 6	Fear of death	19.501	.892	.347
Reason 7	Worry about the material costs of the operation	18.802	.859	.356
Reason 8	Fear of physical disability	142.330	6.832	.010
Reason 9	Fear of complications due to the operation	100.060	4.722	.032
Reason 10	Fear of medical errors	76.782	3.590	.061
Reason 11	Absence from work due to the operation	127.317	6.074	.015
Reason 12	Concern about family	102.866	4.860	.029
Reason 13	Need for blood transfusion	77.037	3.602	.060
Reason 14	Effect of the operation on the appearance of the body	26.130	1.198	.276
Reason 15	Delayed healing of the wound	133.741	6.397	.013
Reason 16	Fear of delirium due to anaesthesia	93.868	4.419	.038
Reason 17	Fear without apparent cause	104.070	4.919	.028



DISCUSSION

The findings of this study indicate that the psychological stress of women before caesarean section mainly ranges between mild to moderate levels. Stress is the normal response for several reasons for preoperative fear such as fear of anaesthesia, fear of death, fear of medical errors, and concern about family, and so on, a reason for preoperative fear that can be increased by psychological stress level (Abarghooe et al.,2022). A study conducted in Pakistan found that preoperative fear of death causes a high level of pre-operative anxiety that can lead to increased postoperative analgesic requirement, prolonged

hospital stay, significant contribution to adverse perioperative outcomes and poor patient satisfaction (Jawaid et al., 2007). A similar study conducted in Turkey showed that perioperative anxiety is associated with increased autonomic fluctuations, a greater need for anaesthesia, a greater possibility of experiencing nausea and vomiting, and increased pain in the immediate postoperative period of surgery (Celik & Edipoglu, 2018). Furthermore, a study conducted in Ethiopia showed that increases the possibility of high psychological stress before the operation resulting from the fear of anaesthesia. In addition, the stress response is influenced by changes in the metabolic, neurohormonal, and

immunological systems as well as anxiety. Therefore, these alterations have been observed because of surgery and stress has been associated with cardiovascular changes and elevated cortisol levels in the blood (Bedaso & Ayalew,2019). In addition, other studies in Turkey have shown that difficulty waking up from anaesthesia, changes in lifestyle, degradation of body image, postoperative pain, disability, and fear of death are the main reasons for their anxiety before surgery (Akutay & Ceyhan,2023). Fear of delayed wound healing can be a reason for preoperative psychological stress that causes certain physiological alterations through manipulation of the immune and nervous systems. However, observational studies have shown that stress and fear before surgery can slow healthy wound healing (Basu et al.,2022).

Conclusions and recommendations:

Psychological stress during pregnancy is caused by several factors and negatively impacts the health of women and their fetuses. Many antianxiety medications are contraindicated for pregnant women; therefore, it is recommended that the stress management intervention be the safest way of reducing surgery-related psychological stress in pregnant women prior to caesarean section.

DECLARATION SECTION

Acknowledgements

We would like to thank and express our deep gratitude to all pregnant women who participated willingly in our study and generously spent time to complete the questionnaire and apply the management therapy. Their active participation was essential for the study to measure the effectiveness of the therapeutic intervention.

Ethical Considerations

Before starting the sample collection, the study was approved by the Research Ethics Committee of the University of Kufa, Faculty of Medicine, that the study was free of any possible harm to the participants (physical or psychological). The women were informed that their participation in the study was voluntary and their information would be kept confidential.

DECLARATION SECTION

Acknowledgments

We thank the anonymous referees for their useful suggestions.

Ethical Considerations

Before starting the sample collection, the study was approved by the Research Ethics Committee of the University of Kufa, Faculty of Medicine, and that study was free of any possible harm to the participants (physical or psychological). The women were informed that their participation in the study was voluntary, and their information would be kept confidential

Conflict of interest

The authors report that there is no conflict of interest.

Funding:

None to be declared.

Data availability

Data are available by contacting the corresponding author by email.

Authors contribution

All authors contributed equally to the conception and design of the study, data collection, and analysis, and drafted the initial manuscript. All authors critically reviewed and edited the manuscript. All authors approved the final version of the manuscript for submission.

Reference

- Abarghooe, S. N., Mardani, A., Baha, R., Aghdam, N. F., Khajeh, M., Eskandari, F., & Vaismoradi, M. (2022). Effects of the Benson relaxation technique and music therapy on anxiety of primiparous women prior to cesarean section: A Randomized Controlled Trial. *Anesthesiology research and practice*, 2022.
- Akutay, S., & Ceyhan, Ö. (2023). The relationship between fear of surgery and factors in surgical patients. *Perioperative Medicine*, 12(1), 1-8. <https://doi.org/10.1186/s13741-023-00316-0>
- Bakr, S. A., Ali, A. P. D. S. K., & Khudhr, A. P. D. S. A. (2014). Effect of age and Previous Surgery Experience on preoperative anxiety. *Kufa Journal for Nursing Science* Vol. 4 (1) <https://doi.org/10.36321/kjns.vi20141.2457>
- Basu, S., Goswami, A. G., David, L. E. & Mudge, E. (2022). Psychological stress on wound healing: A silent player in a complex background. *The International Journal of Lower Extremity Wounds*, [15347346221077571](https://doi.org/10.1177/15347346221077571). <https://doi.org/10.1177/15347346221077571>
- Bedaso, A. & Ayalew, M. (2019). Preoperative anxiety among adult patients undergoing elective surgery: A prospective survey at a

- general hospital in Ethiopia. Patient safety in surgery, 13(1).
<https://doi.org/10.1186/s13037-019-0198-0>
- Celik, F., & Edipoglu, I.S. (2018). Evaluation of preoperative anxiety and fear of anaesthesia using the APAIS score. *European Journal of Medical Research*, 23(1), 1-10.
<https://doi.org/10.1186/s40001-018-0339-4>
- Hobel, C. J., Goldstein, A. M. Y., & Barrett, E. S. (2008). Psychosocial stress and pregnancy outcome. *Clinical Obstetrics and Gynecology*, 51(2), 333-348.
<https://doi.org/10.1097/GRF.0b013e31816f2709>
- Jawaid, M., Mushtaq, A., Mukhtar, S., & Khan, Z. (2007). Preoperative anxiety before elective surgery. *Neurosciences Journal*, 12(2), 145-148.
<https://doi.org/10.1155/2022/9986587>
- Jiwanmall, M., Jiwanmall, SA, Williams, A., Kamakshi, S., Sugirtharaj, L., Poornima, K., & Jacob, K.S. (2020). Preoperative anxiety in adult patients undergoing day care surgery: Prevalence and associated factors. *Indian journal of psychological medicine*, 42(1), 87-92.
https://doi.org/10.4103/IJPSYM.IJPSYM_180_19
- Kaream, A., & Ali, R. (2021). Evaluation of pregnant women's knowledge and health behaviours regarding coronavirus disease precaution measures in primary health care centres in Baghdad City-Iraq. *Kufa Journal for Nursing Sciences*, 11(2), 146-153.
<https://doi.org/10.36321/kjns.vi20212.2058>
- Kiriakopoulos, N., Grigoriadis, S., Maziotis, E., Philippou, A., Rapani, A., Giannelou, P., ... & Simopoulou, M. (2019). Investigating the stress response during vaginal delivery and elective cesarean section through evaluation of cortisol, interleukin 6 (IL-6), growth hormone (GH) and insulin-like growth factor 1 (IGF-1). *Journal of Clinical Medicine*, 8(8), 1112.
<https://doi.org/10.3390/jcm8081112>
- Leardi, S., Pietroletti, R., Angeloni, G., Necozone, S., Ranalletta, G., & Del Gusto, B. (2007). A randomized clinical trial that examines the effect of music therapy on the stress response to day surgery. *Journal of British Surgery*, 94(8), 943-947.
<https://doi.org/10.1002/bjs.5914>
- Levett, D. Z. H., & Grimmett, C. (2019). Psychological factors, pre-habilitation and surgical outcomes: evidence and future directions. *Anesthesia*, 74, 36-42.
<https://doi.org/10.1111/anae.14507>
- Sandall, J., Tribe, R. M., Avery, L., Mola, G., Visser, G. H., Homer, C. S., ... & Temmerman, M. (2018). Short- and long-term effects of caesarean section on the health of women and children. *The Lancet*, 392(10155), 1349-1357.
[https://doi.org/10.1016/S0140-6736\(18\)31930-5](https://doi.org/10.1016/S0140-6736(18)31930-5)
- Teckenberg-Jansson, P., Turunen, S., Pölkki, T., Lauri-Haikala, M. J., Lipsanen, J., Henelius, A., ... & Huotilainen, M. (2019). Effects of live music therapy on heart rate variability and self-reported stress and anxiety among hospitalized pregnant women: A randomized controlled trial. *Nordic Journal of Music Therapy*, 28(1), 7-26.
<https://doi.org/10.1080/08098131.2018.1546223>
- Vanahor, R. P. (2018). Psychological Effects of Caesarean Section on Mothers Delivered at Kenyatta National Hospital. Doctoral dissertation, University of Nairobi, 3972