

## Prevalence of hyperuricemia in diabetes mellitus

*\*Mohammed Abdul Hessein*

### **Abstract**

#### *Objective :*

To explore the prevalence of hyperuricemia in diabetes mellitus patients.

#### **Design :**

Eighty diabetic patients attended diabetic clinic center at teaching hospital in Najaf city between February and June 2006 were evaluated to explore the prevalence of hyperuricemia, thirty of them (group A) had joint problems and fifty of them (group B) did not.

#### **Result:**

Ten patients (33%) from group A had hyperuricemia versus one (2%) from group B.

#### **Conclusion:**

Data from observation study suggest that there is correlation between joint problems and high serum uric acid in diabetes mellitus.

### **Introduction:-**

Uric acid is produced from purines in the liver, which are mostly pass through the kidneys and is excreted in the urine, normally these processes keep the concentration of uric acid in the blood plasma at a healthy level, which is below 6.8 mg/dl<sup>1</sup>.

Under certain circumstances, however, the body produces too much uric acid or excretes too little. In either cases, concentration of uric acid increase in the blood. This condition is known as hyperuricemia<sup>1'2'11</sup>.

Hyperuricemia is related to the individual components of the diabetes mellitus in complex ways. According to one theory, hyperuricemia enhances the effect of insulin resistance on renal urate absorption leading to glucose intolerance<sup>3'4'5'12'15</sup>.

Hyperuricemia is associated with gout , but a minority of people develop it, when the serum is saturated with urate so that other disorders that produce gout-like symptom or causes hyperuricemia should be ruled out eg. pseudogout and septic arthritis 2,6. Hyperuricemia has long been suspected to be a cardiovascular risk factor especially in hypertension 4'7'13 that itself can induce hyperuricemia by decreasing renal flow , increasing urate reabsorption; or by the use of diuretic to control blood pressure8'9'14.

### **Patients and methods :-**

The study was conducted on in-patients in Teaching Hospital in Najaf city during the period from February to June 2006 .

Thirty diabetic patients (group A) ranging in age between( 26-80) years mean age (53) years with joint problems were intended into this study (8 of them male and 22 were female) versus fifty diabetic patients (group B) ranging in age between (18-80) years mean age(49) years . Thirty of them male and 20 were female without joint problems.

Patients were considered to have joint problems if they had joint pain , stiffness , limitation of movement and swelling.

A detailed history and thorough physical examination were performed for each diabetic patient especially looking for duration of diabetes , types of treatment , drugs history related to hyperuricemia , cardiovascular and renal complication .

Laboratory test which were done include the following:

- 1-fasting blood sugar (mmol/l).
- 2- blood urea and serum creatinine (mmol/l).
- 3- serum uric acid (mg/100ml).

Uremic patients and those with history of drug-related hyperuricemia were excluded from the study .

Data obtained were analysed by *chi* square and *p*- value <0.05 was considered significant.

### **Result:-**

In group(A) , ten patients (33%) were found to have hyperuricemia ,where as in group(B) only one patient (2%) was

found to have hyperuricemia , this was statistically significant (P. value <0.05) . Table I, figure(A).

Nineteen patients (63%) in group (A) had a cardiovascular disease in a form of hypertension or ischemic heart disease compared with thirteen patients (26%) in group(B) had a cardiovascular disease which was statistically significant. Table II , figure (B).

Six of the ten hyperuricemic patients in group (A) (60%) had a cardiovascular disease and the only one hyperuricemic patient in group (B) had a cardiovascular disease, this was statistically not significant (P. value>0.05) . Table III.

There was an increase in the prevalence of hyperuricemia in diabetes mellitus patients with increasing age . Table IV , figure (C)

There was a correlation between the age and the involvement of the joints so that in the middle age groups ,lower limbs involvement is more than the upper limbs involvement and mostly one joint involvement while in the old age groups upper limbs involvement is more ,with more than one joint involvement . Table V.

Table I ( prevalence of hyper uricemia in DM with joint problems versus DM without joint problems) .

group	Number of patient	Number of hyperuricemia	%
DM with joint problems	30	10	33
DM without joint problems	50	1	2
total	80	11	13

**P. value <0.05 (statistically significant)**

**DM = diabetes mellitus.**

Table II. (prevalence of cardiovascular disease in diabetes mellitus)

group	Number of patient	Number of cardiovascular disease	%
<b>DM with joint problems</b>	<b>30</b>	<b>19</b>	<b>63</b>
<b>DM without joint problems</b>	<b>50</b>	<b>13</b>	<b>26</b>
<b>Total</b>	<b>80</b>	<b>32</b>	<b>40</b>

Table III. (prevalence of hyperuricemia in cardiovascular disease)

group	Hyperuricemic patient	Hyperuricemia with cardiovascular disease	%
<b>DM with joint problems</b>	<b>10</b>	<b>6</b>	<b>60</b>
<b>DM without joint problems</b>	<b>1</b>	<b>1</b>	<b>1</b>
<b>Total</b>	<b>11</b>	<b>7</b>	<b>63</b>

**P.value < 0.05**

**P. value > 0.05 (statistically not significant)**

Table IV. (prevalence of hyperuricemia with age)

Age/years	Number of patient	Hyperuricemia with joint problems	%
<b>&lt; 24</b>	<b>0</b>	<b>0</b>	<b>0</b>
<b>25-34</b>	<b>3</b>	<b>0</b>	<b>0</b>
<b>35-44</b>	<b>3</b>	<b>0</b>	<b>0</b>
<b>45-54</b>	<b>3</b>	<b>1</b>	<b>33</b>
<b>55-64</b>	<b>13</b>	<b>3</b>	<b>23</b>
<b>65-74</b>	<b>5</b>	<b>3</b>	<b>60</b>
<b>75-84</b>	<b>3</b>	<b>3</b>	<b>1</b>
<b>Total</b>	<b>30</b>	<b>10</b>	<b>33</b>

Table V.( correlation between age and involvement of joint)

Age/years	Upper limb joint involvement	Lower limb joint involment
40-59 (middle)	0	3
<b>60-80( old)</b>	<b>4</b>	<b>3</b>




Figure A (relation ship between serum uric acid & prevalence of hyperuricemia)





Figure B relation ship between serum uric acid & prevalence of cardio vascular disease.



## Discussion and Conclusion:-

We have taken thirty diabetic patients with joint problems, (33%) of them were found to have hyperuricemia and these result were similar to study done by Andrew J . Luk 5'11 who studied hyperuricemia and diabetic in sixteen patients with (18%) had joint problems versus eight without joint problems .

He found that hyperuricemia induce impaired glucose tolerance by oxidative stress (glucose auto-oxidation and advanced glycosylation end product )4"8"9'12.

The prevalence of hyperuricemia with a cardiovascular disease was correlated with diabetic patients with joint problems these result were consistent with study done by Framingham heart study found no independent association between hyperuricemia and increased risk of cardiovascular diseases (in both studies, statistically not significant. 5"7"13 )

The prevalence of hyperuricemia in diabetic patients with joint problems was increased with advancing age 1"2.

There was a correlation between age and limbs involvement in the middle age groups; lower limb joints involvement being more than the upper limb joints; while in old age groups ,upper limb joints involvement is more than that in the lower and this is correlated with study done by Niskaneu Lk , who explained that middle age adult association in groups with obesity ,high blood pressure, unhealthy cholesterol level and heavy alcohol use, while elderly is associated with kidney problems and the use of diuretic 5"10"14.

As seen from this study there is correlation between joint problems and high serum uric acid in diabetes mellitus.

References:-

1. Harvey simon ,MD, gout an arthritic condition association with chronic hyperuricemia (long lasting high concentration of uric acid in the blood).(800-283-7800),review data:5/13/2004
2. Wallac SL, Robinson H, Masi AT . classification of acute arthritis of primary gout .arthritis Rheum 1977 ; 20 : 895-900.
3. Hall Ap , Barry PE , Dawber TR . epidemiology of gout and hyperuricemia .A long – term population study. Am J Med.1967 ; 42 : 27-37.
4. Boyko EJ , De Courten M . feature of metabolic syndrome predict high risk diabetes and impaired glucose tolerance : a prospective study in Mauritius. diabetes care .2000 ; 23 : 1242 - 1248.
5. Andrew J , Luk , MD , Mph . recent advance in epidemiology of gout and hyperuricemia . AM J Manage care . 2005 ; 11 : S435 – S442.
6. Campion EW, Glynn RJ , Delabry LO . asymptomatic hyperuricemia . risks and consequences in the normative aging study . AM J Med 1987 ; 82 :421-6.
7. Baker JF, Krishnan E , Chenl . serum uric acid and a cardiovascular disease :recent development , and where do they leave us ? Am J Med 2005;118: 816-26.
8. Culleton BF, Larson MG , Kannel WB .serum uric acid and risk for cardiovascular disease and death: the Framingham heart study . Ann intern Med.1999 ; 131 : 7-13.
9. Melvin R Haydn. Uric acid new book at an old risk mark for a cardiovascular disease , metabolic syndrome , type 2 diabetes mellitus ( urate redox shuttle).Mrh29@usmo.com ;Revised August 23,2004.
10. Niskanen LK , Laaksonen DE . uric acid level a risk factor for a cardiovascular and all cause mortality in the middle age men : a prospective cohort study .Arch intern Med . 2004 ; 164:1546- 1551 .



11. Lin KC , Lin HY , Chou P . community based epidemiological study on hyperuricemia and gout in kin-Hu , Kinmen. *J rheumatol.* 2000 ; 27 : 1045 – 1050.
12. Meigs JB , Wilson PW, Nathan Dm . prevalence and characteristics of the metabolic syndrome in the san Antonio heart and Framingham of spring studies diabetes . 2003 ; 52 : 2160-2167 .
13. Johnson RJ , Feig DI . resurrection of uric acid as a causal risk factor in essential hypertension . 2005 ; 45 : 18-20.
14. Choi HK , Atkieson K. obesity , weight change , hypertension , diuretic use , and risk of gout in men : the health professional follow up study . *Arch intern Med* . 2005 ; 165 : 742 – 748 .
15. Arromdee E , Michet CJ . epidemiology of gout : is the incidence rising ? *J Rheumatol.* 2002 ; 29 : 2403 – 2406 .