

**A PROTOCOL FOR SHOOT REGENERATION FROM LEAVES
PETIOLES TISSUE CULTURE OF NEEM TREES (*Melia azedarach*)**

Mozahim K. AL- Mallah

Shifa M. Salih

College of Education , Dept. of Biology,
University of Mosul , Mosul – Iraq**ABSTRACT**

Shoots were regenerated from petioles – derived callus on solidified Murashige and Skoog (MS) medium supplemented with 1.5 mg / l BA only and MS medium containing 4.0 mg / l IBA and 1.0 mg / l BA . The percentage of shoot formation was 30.0 % . The regeneration frequency was reasonable to obtained sufficient shoots by this method . This simple protocol is the first report in tissue culture of this woody plant which encourage other researchers to worke other studies in tissue culture of neem . Moreover , callus culture will be useful to obtain a wide range of industrial plant products .

INTRODUCTION

Sababnah (*Melia azedarach* L.) is a fast growing , deciduous tree, belonging to the family Meliaceae which contains other species of neem trees, *Azadirachta indica* (Allan *et al.* ,1999) . This woody tree is widely exploited for its effects on insects (Kearney *et al.* , 1994) . Current interests in its use as a natural insecticide is concentrated on compounds that exhibit antifeedant and insect growth regulatory effects (Mordue & Blackwell , 1993) . Most of the active principles from neem fruits as well as from seeds , stem and root bark are tetranortriterpenoids , although biologically active diterpenoids , triterpenoids constituents are also present . More than three hundred compounds have been isolated from various parts of the tree (Kumar *et al.*, 1996) . and most attention has been directed towards the major principle , Azadiractin . Several studies have centered on *A. indica* callus induction and organogenesis (Sanyal *et al.*, 1981 ; Schultz , 1984 ; Ramesh & Padhya , 1988) Ramesh and Padhya (1990) studied the production of adventitious shoot buds from leaf discs on Wood and Braun medium supplemented with Kinetin , Benzyl adenine and adenine sulphate . They reported that although there was no growth on basal medium , addition of kinetin or benzyl adenine alone resulted in callus production . While addition of both at concentration of 4mM each resulted in shoot formation . Similar results were found by culturing nodal segments and leaf discs of *A. indica* on MS (Murashige & Skoog ,1962) medium supplemented with 1.0 mg/L benzyl adenine and 0.5 mg/L kinetin (Subramani *et al.*, 1993) . In later studies , Eeswara (1997) failed to obtain shoot formation in any of the above combinations . The possibility of callus initiation and shoots regeneration of *Melia azedarach* was investigated in this study and according to our knowledge there are no published reports on tissue culture of *Melia azedarach* in our country, The paper represent the

first report on tissue culture of *Melia azedarach*.

MATERIALS AND METHODS

Collection of plant materials : Trees of *Melia azedarach*, grown in Mosul University's park (Fig.1,A) were used as a source of plant material which include young leaves and its petioles .

Surface sterilization of specimens : Young leaves were picked from Sabahbah trees (15 years old) , they were thoroughly washed with tap water before surface sterilization . The petioles were cut into segments of 1 cm length while the leaves cut into pieces of 0.5 cm .These explants were surface sterilized according to Eeswara *et. al.*, (1997) with some modification , by immersion into 0.1 % HgCl₂ solution for 5 min. and washed with sterile water . These treated explants were immersed in 3% commercially available sodium hypochlorite solution for 30 min.,then washed carefully with sterile water

Callus initiation : Sterilized leaves and petioles of *Melia azedarach* were cultured on the surface of 15 ml of agar –solidified MS (Murashige & Skoog ,1962) medium supplemented with different combinations of Naphthalen acetic acid (NAA) with Benzyladenine (BA) or 2,4-dichlorophenoxy acetic acid (2,4-D) with Kinetin . Also Indole butyric acid (IBA) was used . Many initiation media were tested for callus formation from leaf and petiole segments as indicated below :

NAA and BA combinations

- MSO (Hormone - free) control
- MS + 1.5 mg/l BA
- MS + 0.9 mg/l NAA
- MS + 0.2 mg/l NAA + 0.1 mg/l BA
- MS + 2.0 mg/l NAA + 2.0 mg / l BA
- MS +1.0 mg/l NAA + 2.0 mg l BA

2,4-D and Kinetin combinations

- MS +0.1 mg /l 2,4 - D + 0.2 mg /l Kin.
- MS + 0.5 mg /l 2,4 - D + 1.0 mg /l kin.
- MS + 1.0 mg /l 2,4 - D + 1.0 mg /l kin.
- MS + 1.0 mg /l 2.4 - D + 2.0 mg /l Kin.
- MS + 2.0 mg /l 2,4 - D + 4.0 mg /l Kin

IBA and BA combinations

- MS + 0.1 mg /l IBA
- MS + 2.0 mg /l IBA + 1.0 mg /L BA
- MS + 4.0 mg /l IBA + 1.0 mg /L BA

Plant specimens were kept in culture room at $25 \pm 1^{\circ}\text{C}$, 1500 Lux white fluorescent tube for 16 hrs . light 8 hrs. dark .Other group of samples were incubated at same condition in the dark .

RESULTS AND DISCUSSION

This paper indicates the response of *Melia azedarach* tree to plant tissue culture technique throughout the good response of leaves and petioles explants to callus formation on MS medium supplemented with various combination of different auxins (NAA , IBA and 2,4-D)and cytokinines(BA and kinetin) in the dark .

Various concentrations of BA and NAA were helpful in inducing callus initiation from leaves and petioles explants (Table 1) .

This good response of explants for callus formation may be due to the efficient selection of cells in an explants (Al-Mallah , & Hassan ,1994) coupled with compatibility between the endogenous hormonal levels and the addition of growth regulator to the induction medium .

Table 1 : Callus formation from leaves and petioles explants of *Melia azedarach* on agar-solidified MS medium supplemented with NAA and BA

Medium (mg /L)	Cult. Segment / Init. Segment		Callus formation (%)	
	Leaves	Petioles	Leaves	Petioles
MS + 1.5 mg/L BA	40 [*] /20 ^{**}	40/33	50	82.5
MS + 0.2 mg/LNAA + 0.1 mg/L BA	30/0	30/0	0.0	0.0
MS + 0.9 mg/LNAA	30/0	30/0	0.0	0.0
MS + 1.0 mg/LNAA + 2.0 mg/L BA	40/32	40/21	80.0	52.5
MS + 2.0 mg/LNAA + 2.0 mg/L BA	30/3	30/2	10	6.6
MSO (control)	30/0	30/0	0	0

* Cult . :cultured segments , **Init .:Initiated segments

The obtained results indicate clearly that MS medium supplemented only with 1.5 mg /l BA sustained callus induction . The petiole explants started callus formation after two weeks of culture and the percent of callus initiation was 82.5 .It was obvious that increasing NAA level inhibited callus formation from 82.5% to 6.4 % (Table 1) .These results confirm that this plant have enough levels of endogenous hormones and does not require the addition of high levels of exogenous growth regulators (Wala & Jasrai , 2003).Subramani *et al.* (1993) noticed that agar solidified MS medium supplemented with 3.5 mg /l BA was proper for callus initiation in neem. The results indicated that replacing NAA with IBA seems to be very useful to callus initiation as the induction of callus increased (Table 2) .

The petiole explants began callus induction within twelve days (Fig.1 ,B). While leaves explants started callus formation after twelve days on agar solidified MS medium containing IBA 4.0 mg /L and BA 1.0 mg / L (Fig.1 ,C) .The results showed that this medium was the best one for callus growth, so it was used for the subculturing and maintenance of callus every 4 weeks interval . Callus was friable and yellow light brown in colour (Fig.1,D). Previous study (Allan *et at* .,1994) showed that agar solidified MS medium supplemented with

IBA 4.0 mg / L and BA 1.0 mg / L was suitable for callus initiation from leaves explants of *Azadirachta indica* .

Table 2 : Initiation of callus from leaves and petioles of *Melia azedarach* on agar solidified MS medium supplemented with IBA and BA

Medium	Cult. Segment / Init. Segment		Callus formation %	
	Leaves	Petioles	Leaves	Petioles
MS + 0.1 mg/l IBA	30/0	30/0	0.0	0.0
MS + 2.0 mg/l IBA + 1.0 mg/L BA	30/1	30/2	3.3	6.6
MS + 4.0 mg/l IBA + 1.0 mg /l BA	50/21	50/42	42	84
MSO(control)	20/0	20/0	0.0	0.0




Fig .1: Shoot regeneration from leaves and petioles tissue culture of *Melia azedarach* L.

(A) : *Melia azedarach* tree used as a source of plant material ,(B) : Petioles explants of *Melia azedarach* after twelve days of culture on MS medium supplemented with IBA 4.0 mg / L and BA 1.0 mg /L ,(C) : Leaves explants after twelve days of culture on MS medium supplemented with IBA 4.0 mg / L and BA 1.0 mg /L. ,(D) : Callus of *Melia azedarach* after 30 days on(MS + IBA 4.0 mg / L + BA 1.0 mg /L) ., (E) : Shoots formation from Petioles arrowed derived callus on MS medium supplemented with 1.5 mg / L BA. ,(F) : Shoots formation arrowed from Petioles explants on (MS + 4.0 mg / L IBA + 1.0 mg /L BA) .

Moreover , 2,4-D and Kinetin can be used in callus initiation of *Melia Azedarach* since callus was continuously induced at ratio of 60.0% (Table 3) .

Table 3 : Callus formation from leaves and petioles of *Melia Azedarach* on agar solidified MS medium supplemented with different concentration of 2,4-D and kinetin

Medium (mg /l)	Cult. Segment / Init. Segment		Callus formation %	
	Leaves	Petioles	Leaves	Petioles
MS + 0.1 mg/l 2,4 - D + 0.2 mg /l kin	30/0	30/0	0.0	0.0
MS + 0.5 mg /l 2,4 - D + 1.0 mg /l kin	40/2	40/3	5.0	7.5
MS + 1.0 mg /L 2,4 - D + 1.0 mg /L ki	30/0	30/0	0.0	0.0
MS + 1.0 mg /l 2,4 - D + 2.0 mg /l kin	30/1	30/3	3.3	10.0
MS + 2.0 mg /l 2,4 - D + 4.0 mg /l kin	30/8	30/18	26.6	60.0
MSO(control)	30/0	30/0	0.0	0.0

This result express the capability of this tree to invitro culture . As previously published with *Azadirachta indica* Nirmalakumari *et al.* (1996) indicated that MS medium containing 2.0 mg / L 2,4-D and 0.5 mg /l kin. promoted callus formation from cotyledones and leaves explants .

All the above results of callus initiation occurred in the dark .Conditions of 8 hrs. dark 16 hrs. Light was not suitable for all tested materials and we do not notice any sign for callus induction .Wewetzer (1998) indicated that the dark was preferable for callus initiation of *Azaderacta indica* .

Plant regeneration:The results showed that among all the different agar solidified MS media used , those supplemented with 1.5 mg /l BA only and MS medium containing 4.0 mg / l IBA and 1.0 mg / l BA stimulated shoots formation from petioles derived callus of *Melia Azedarach* .(Fig.1,E)and from petiole explants (Fig.1,F) and shoot formation reached 30% , and 6.0 % respectively (Table 4) .

Table 4 : Shoot regeneration from petioles-derived callus of *Melia azedarach* on agar solidified MS medium supplemented with IBA and BA

Regeneration medium	No. of cultured explants	No. of shoots	shoot regeneration (%)
MS + 1.5mg/l BA	40	12	30.0
MS + 4.0 mg/l IBA +1.0 mg /l BA	50	3	6.0

It was observed that there are different factors influencing tissue growth and differentiation , these factors include explant source and the addition of growth regulators (Mihaljevi *et al.* , 2002)

In conclusion this finding represents a simple protocol for callus induction and shoot formation from leaves and petioles of *Melia azedarach* . This will enhance tissue cultures works on this woody plant by other researchers and promote other workers to start extracting Azadirachtin and other secondary metabolites from callus cultures of this important tree ,or regenerating transformed *Melia azedarach* plants as happened with medicinal plant *Solanum nigrum* (AL-Mallah &Salih ,

Mesopotamia J. of Agric

(ISSN 1815-316 X)

Vol. (34) No. (1)

2006

2005). And for future work electrostimulation can be applied to increase probably growth of callus , since electrostimulation was found to enhance growth of callus of tobacco (Al-Mallah ,1993),and black night shad *S .nigrum* (Al-Mallah &Salih ,2003) .

طريقة بسيطة للحصول على الافرع الخضرية من المزارع النسيجية لاعناق اوراق اشجار النيم (*Melia azedarach*)

مزارح قاسم الملاح
شفاء مهدي صالح
قسم علوم الحياة/ كلية التربية /جامعة الموصل / الموصل / العراق

الخلاصة

تمكنت الدراسة الحالية من الحصول على الافرع الخضرية من كالس أعناق الاوراق لاشجار السبج *Melia azedarach* في وسط موراشيج وسكوك (MS) الصلب المدعم باضافة ١.٥ ملغم / لتر بنزائل أدنين (BA) ووسط (MS) الصلب المدعم باضافة ٤.٠ ملغم / لتر (IBA) و ١.٠ ملغم / لتر (BA) وأظهرت النتائج تمايز مزارع الكالس وبلغت نسبة تكون الافرع الخضرية ٣٠% MS في ٦.٠ و ٦.٠ على التوالي . ومن المحتمل ان تكون هذه الدراسة الاولى في مجال الزراعة النسيجية لاشجار السبج في العراق .ان هذه الدراسة تشجع العديد من الدراسات في مجال الزراعة النسيجية لهذه الاشجار او تلك المتعلقة بالحصول على النواتج الايضية او بالتحول الوراثي لهذه الاشجار المنتشرة في البيئة العراقية .

REFERENCES

- Al- Mallah, M. K. (1993) . Electroporation enhanced plant regeneration from callus of *Nicotiana tabacum* . 2nd Arab conf. Modern Biotech. Amman . Jordan, 331 –335.
- Al- Mallah, M. K. and S.M. ,Salih (2003) . Electroporation increased growth of callus , regeneration capability and protein content of *Solanum nigrum* L. Plants . Raf . J . Sci . 14: 35 –42 .
- Al- Mallah, M. K. and S.M. , Salih (2005) . An efficient technique for obtaining transformed *Solanum nigrum* L.plants via *Agrobacterium rhizogenes* R1601 . Raf . J . Sci . (accepted) .
- Al- Mallah, M. K.and H. A. , Hassan (1994).Callus formation from cultured leaf and stem explants of Harmala (*Peganum harmala*).J.Educ. Sci.20:96 – 101.
- Allan , E. ; P , Janakie . and J , Easwara . (1994) . The production of azadirachtin by *in vitro* , tissue cultures of neem , *Azadirachta indica* . Pestic . Sci . 42 : 147 –152 .
- Allan, E.J ; T , Stuchbury .and M , Luntz .(1999) . *Azadirachta indica* A .Juss. (Neem Tree): In vitro culture , Micropropagation ,and the production of Azadirachtin and Other secondary metabolites . Biotechnology in Agriculture and Forestry .43:11-41.
- Eeswara , J. (1996) . The production of the insect antifeedant azadirachtin and related metabolites by plant cell and tissue culture of neem (*Azadirachta indica* A. Juss) . Ph.D thesis , University of Aberdeen . UK.

- Eeswara , J. ; T , Stuchbury . ; E , Allan . and A , Mordue . (1997) . A standard procedure for the micropropagation of neem (*Azadirachta indica* A. Juss) . Plant Cell Repts . 17: 215- 219 .
- Kearnay ,M ; E.J , Allan . ; J , Hooker. and A , Mordue . (1994) . Antifeedant effects of in vitro culture extracts of neem tree . *Azadirachta indica* against the desert locust *Schistocerca gregaria* (Forskal) . Plant Cell Tiss. Org .Cult . 37 : 67_71.
- Kumar , C. ; M , Srinivas.and S , Yakkundi. (1996) .Limonoids from the seeds of *Azadirachta indica* .Phytochemistry 43 : 451 -455 .
- Mihaljevi , S. ; I , Bjedov . And M , Kova . (2002) . Effect of explant source and growth regulators on in vitro callus growth of *Taxus baccata* L. Washingtonii . Food Technol . Biotechnol . 4 :299 –303.
- Mordue , A .and A , Blackwell .(1993) .Azadirachtin . An update .J Insect Physiol . 39 : 903-924.
- Murashige , T. and F , Skoog . (1962) . A revised medium for rapid growth and bioassays with tobacco culture . Physiol Plant , 15 : 473 – 497 .
- Nirmalakumari , A. ; N , Ramaswamy . and R , Sree.. (1996) . Tissue Culture Studies in Neem (*Azadirachta indica* A. Juss) .In : Singh R P , Chari , MS. , Raheja AK , Kraus W (eds) Neem And Environment , Vol 2 Science publishers , Lebanon , New Hampshire , PP : 982 –992 .

- Ramesh , K. and M , Padhya. (1988) . Isolation of nimbin from *Azadirachta indica* leaves and its callus cultures . Indian Drug . 12 :526-527 .
- Ramesh , K. and M , Padhya . (1990) .In vitro propagation of neem *Azadirachta indica* (A. Juss)from leaf discs. Indian J. Exp. Biol.28 : 932-935 .
- Sanyal , M. ; A , Dass. ; M , Banerjee. and P , Datta. (1981) . In vitro hormone induced chemical and histological differentiation in stem callus of neem *Azadirachta indica* A. Juss . Indian J. Exp. Biol . 19: 1067 - 1068 .
- Schulz ,F. (1984) . Tissue culture of *Azadirachta indica* . In : Schmutterer H, , AscherKRS (eds) . Natural pesticides from the neem tree and other tropical plants .proc. 2nd . Int Neem Conf. GTZ , Eschborn , pp :539 –542 .
- Subramani , J. ;C. Nair;M. Gopinathan and,S. Pillai. (1993) . Large Scale Production of neem Through in Vitro Methods . In : Kleeberg , H. (ed) Practice Oriented Results on Use And Production of Neem Ingredients And Pheromones . Proc. 2nd workshop Druck and Graphic , Giessen . pp : 137- 143 .
- Wala , B. and,Y. Jasrai (2003) . Micropropagation of an endangered medicinal plant : *Curculigo orchoides* Gaertn . Plant Tissue Cult . 1 :13 –19 .
- Wewetzer , A. (1998) . Callus cultures of *Azadirachta indica* and their potential for the production of Azadirachtin . Phytoparasitica 26: 47 – 52 .