



Available online at: [www.basra-science\\_journal.org](http://www.basra-science_journal.org)

ISSN -1817 -2695



## ***Podaxis pistillaris* (Gasteromycetes) from the desert of southern Iraq, an addition to the known mycota of Iraq**

Tawfik M Muhsin, Abass F Abass and Ekhlas K Al-Habeeb

Department of Biology, College of Education, University of Basrah, Iraq

Email: [tmuhsin2001@yahoo.com](mailto:tmuhsin2001@yahoo.com)

Received 20-2-2012 , Accepted 12-6-2012

### **Abstract**

The desert gastroid mushroom *Podaxis pistillaris* (L. ex Pers.) Fr. collected from sandy desert habitat in southern Iraq was identified for the first time from this region. A detailed description and photographs are provided in addition to some information on its world distribution and traditional medicinal significances are discussed based on the available literature.

**Key words:** Desert mushroom, folk medicine, *Podaxis*, Iraq

### **1. Introduction**

*Podaxis* is a gastroid mushroom within the family Agaricaceae (Basidiomycota) which is morphologically identical to stalked-puffballs and widely distributed in desert soils of arid and semi-arid over the world [1,2]. In spring after rainfall, the fruit bodies of this mushroom appear singly or in groups. Taxonomically, about 50 species have been described within this genus, nonetheless, many of them may represent morphotypes of *Podaxis pistillaris* [1]. It has been speculated that *Podaxis* was related to the genus *Coprinus* [3], which recently has been confirmed through the use of DNA sequencing techniques [4]. *Podaxis* has been used in traditional folk medicine in many countries such as in Yemen for the treatment of skin diseases [5], in South Africa against sunburn [6], and in Mali for

wound-healing [7]. In addition, it has been used as face paint and to darken white hair by aborigines in Australia [8] and used as food in countries such as India, Afganistan [9] and Saudi Arabia [10].

The southern desert of Iraq is a vast arid area dominated by sandy coarse alkaline soils. Sodium chloride contributes to the salinity level reaching 14.8 mmohs/cm [11]. Plants communities cover the desert mainly xerophytic, xerotolerant and xerohalophytic species [11]. Despite *P. pistillaris* is quite common in the southern desert of Iraq there is a lack of information, however, regarding this mushroom and there is only a single report on some other mushrooms in southern Iraq [12]. In the present report a description of *P. pistillaris* as a first record in Iraq is provided.

## 2. Materials and method

Fruiting bodies of *P. pistillaris* were collected from sandy soils in Basrah province, southern Iraq (latitude 30 N and longitude 47 E) during February of 2009 and 2010 after recent rain fall. Samples were placed in paper bags and brought to

the laboratory and immediately examined, described and identified according to the available literature. Specimens from our collections were also confirmed by Dr. Michael A. Castellano, United State Department of Agriculture, Corvallis. USA.

## 3. Results and discussion

### 3.1. Taxonomy

*Podaxis pistillaris* (L. ex Pers.) Fr.

Figures (1-2)

BASIDIOMATA (Gasterocarps) up to 15 cm in height, 0.8-1.6 cm in diam., whitish at early stages becoming yellowish to rusty-brown in color at maturity, covered

with scales when young. PILEUS cylindrical to ellipsoidal, 5-9 cm in length, 1-2 cm in diam., white becoming yellowish-brown in color.



Figure 1. A. Field habit of fruit body of *P. pistillaris* growing in sandy desert, B. Different stages of fruit bodies, C. Spores at immature stage (40 x magnification), D. Mature spores with brown color and germ pores (40 x magnification).




Figure 2. A. Mature fruit body of *P. pistillaris* with cracked peridium covers a mass of dark brown spores, B. Longitudinal section of young fruit body, C. Section shows a young gleba with pinkish colored spores, D. Transverse section of the stipe with a reddish mycelia tissue and a hollow.

PERIDIUM non-dehiscent and remaining fused with the stipe (secotioid), cracking or splitting when dried to release spores. STIPE 3-14 cm long, 0.8-1.3 cm diam., white to yellow-brown, straight, bulbous at the base, possessing a fibrous to woody texture, hollow in the centre when mature. GLEBA pink in color when young becoming reddish to brown at maturity. BASIDIOSPORES 10-14 x 9-13  $\mu$ m, globose to subglobose, smooth, thick-

walled, purple when young becoming reddish to dark brown at maturity, with a germ pore.

**Specimens examined:** Collected from Safwan and Um-kasir desert sandy coarse soil, at February 2009, February 2010. Specimen No 110, 111 TMM, AFA. Slide No 120, 121 Deposited at the Mycology Lab, Biology Department, Basrah University, Iraq.

### 3.2. Remarks

The morphological characteristics of *P. pistillaris* inhabiting the desert region of southern Iraq is similar to the other specimens described elsewhere. [1,2,13,14,15] with minor variations such as spore shape in our collections being mostly globose compared with those previously described.

In comparison with reports from nearby desert areas, the specimen from southern Iraq it looks morphologically identical with those reported from Kuwait [16] and Qatar [17], however, no microscopic details were given by those authors.

This species is not uncommon as a saprophyte in sandy ecosystems (Dr. Michael Castellano, pers. comm.). Although *P. pistillaris* is considered by many authors to be a stalked puffball but it is more closely allied with the shaggy mane (*Coprinus comatus* (Mull.) Pers. than with puffballs [4].

Physiological studies on *P. pistillaris* using heavy metals indicated that the growth of this mushroom is affected by amending cadmium and lead in culture

medium [10]. Temperature effects on the growth of this fungus have been also conducted [18]. Cultivation of fruiting bodies of this desert mushroom has also been reported [19]. Medicinally, this species has been used in many traditional folk treatments (Table 1) and recently it has been demonstrated that *P. pistillaris* exhibits a bioactive compounds against bacteria [5,20]. The edibility of this mushroom in some places over the world has also been reported [5,10,21]. Moreover, chemically, the fruiting bodies of *P. pistillaris* contain 76% moisture, 5% total nitrogen, 22-37% total crude protein, 18.5% carbohydrates, 2.3% total lipids and 2.4% ash [22]. It is worth mentioning that this desert mushroom is eaten by local inhabitants living in the southern Iraq. Seemingly, this mushroom possesses a unique life strategy to ensure its survival under extreme desert environments. Therefore, more research is needed to investigate the tolerance of this fungus to such a harsh habitat and whether it lives as a saprophyte or behaves as an ectomycorrhizal partner in association with some desert inhabiting plants.

**Table 1. Distribution and significances of the desert mushroom *P.pistillaris* over the world**

Geographical locations	Local names	Significances	References
Afghanistan		Edible	Jiskani (2001) [9]
Africa		[Folk medicine	Dring (1964) [2]
Argentina			Martinez (1971) [23]
Australia	Desert shaggy mane	Darken the white hair in the whiskers and face paint	Cleland & Johnston (1933) [8]
Brazil			Basiea & Galvao (2002) [15]
China		Inflammation treatment	Mao (2000) [24]
India	Khumbahi	Edible	Jiskani (2001) [9]
Iran			Watling and Gregory (1977) [25]
Pakistan			Sultana et al. (2007) [26]
Mexico			Martin et al. (2005) [27]
South Africa		Folk medicine against sunburn	Bottomley (1948) [6]
South Africa (Sahara)		Poisonous	Walley & Rammeloo (1994) [28]
Saudi Arabia	Al-Arjoon	Edible	Hashem & Al-Rahmah, (1993) [10]
Kuwait	Al-Arjoon		Moustafa (1975) [16]
Qatar	Al-Arjoon		Al-Thani (2010) [17]
Yemen		Treatment of skin disease	Al-Fatimi et al. (2006) [5]
USA	Desert shaggy mane		Brasfield (1937) [13]
Iraq	Kama?	Edible	Present study

#### 4. Acknowledgment

We would like to thank Dr. Leonard Hutchinson, Department of Plant Pathology, College of Agriculture, University of LakeHead, Thunderbay, Ontario, Canada, for revising this manuscript and for his

valuable suggestions. Thanks are due to Dr. Michael A. Castellano, United State Department of Agriculture, Corvallis. USA. for the confirmation of the identification of the fungus specimen.

## 5. References

1. Morse EE. 1933. A study of the genus *Podaxis*. Mycologia 25: 1-33.
2. Dring DM. 1973. Gasteromycetes. In: The Fungi. Vol. IVB. GC Ainsworth, FK Sparrow, AS Sussman (Eds.), Academic Press, New York.
3. Smith AH 1973. Agaricales and related Secotoid Gasteromycetes. In: The Fungi. Vol. IVB. GC Ainsworth, FK Sparrow, AS Sussman (Eds.), Academic Press New York.
4. Hopple JS Jr, Vilgalys R. 1994. Phylogenetic relationships among coprinoid taxa and allies based on data from restriction site mapping of nuclear rDNA. Mycologia 86: 96-107.
5. Al-Fatimi MA, Julich WD, Jansen R, Lindequist U. 2006. Bioactive components of the traditional used mushroom *Podaxis pistillaris*. eCAM . 3: 87-92.
6. Bottomley EA 1984. Gasteromycetes of South Africa. Bothalia 4:473-810.
7. Diallo D, Sogn C, Samake FB, Paulsen BS, Michaelsen TE, Keita A. 2002. Wound healing plants in Mali, the Bamako region. An ethnobotanical survey and complement fixation of water extracts from selected plants. Pharmac Biol 40(2): 117-128
8. Cleland JB, Johnston TH. 1933. The ecology of the aborigines of Central Australia; botanical notes. Trans Proc Royal Soc South Austr 57: 113-124.
9. Jiskani MM. 2001. Growing mushrooms: Step ahead to boost up the economy of Pakistan. Pakis J Econom Manag . p 15-17.
10. Hashem AR and Al-Rahmah AN. 1993. Growth of *Podaxis pistillaris* collected from Saudi Arabia at different concentrations of Cadmium and lead. J King Saud Univ 5:127-135.
11. Thalen DCP. 1979. Ecology and utilization of desert shrub rangelands in Iraq. W. Junk Publish. The Hague, Netherland PP.448.
12. Ewaze JO, Muhsin TM. 1988. New records of Agaricales from Iraq. Basra J Agric Sci 2: 105-111..
13. Brasfield TW. 1937. The morphology of *Podaxis*. Univ Iowa Study Nat Hist 17: 100-121.
14. Bessey EA. 1964. Morphology and Taxonomy of fungi. Hafner Pub. Co. New York.
15. Baseia IG, Galvao TC. 2002. Some interesting Gasteromycetes (Basidiomycota) in dry areas from Northeastern Brazil. Acta Bot Bras 16:1-9
16. Moustafa AF. 1975. A preliminary annotated list of fungi from Kuwait. J Univ Kuwait 2:67-87.
17. Al-Thani RF. 2010. Survey of macrofungi (including truffles) in Qatar. KBM J Biology 1:26-39.
18. Khan S M, Khan DA. 1979. Temperature studied on *Podaxis pistillaris*. Mycologia 71: 861-867.
19. Phutela RP, Kaur H, Sohdi HS. 1998. Physiology of edible gasteromycete *Podaxis pistillaris* (L. ex Pers.) Fr. Indian J Mycol Plant Pathol 28:31-37.
20. Khalaf KT. (2008). Extraction, purification and identification of bioactive substances from basidiomycetes. PhD Thesis, University of Basrah, Iraq. Pp 126.
21. Gupta S & Singh SP. 1991. Nutritive value of mushroom *Podaxis pistillaris*. Indian J Mycol Plant Pathol 21:273-276.
22. Khaliel AS, Abou-Heilah AN, Kassim MY. 1991. The main constituents and nutritive value of *Podaxis pistillaris*. Acta Bot Hung 36:173-179.
23. Martinez A. 1971. Notes sobre el genero *Podaxis* (Gasteromycetes)



- en Argentina. *Biol Soc Argen Bot* 14:73-87.
24. Mao XL. 2000. The macrofungi in China. Zhengzou, Henan Science and Technology Press, China.
25. Watling R, Gregory NM. 1977. Larger fungi from Turkey, Iran and neighboring countries. *Karstenia* 17: 70.
26. Sultana K, Gul M, Firdous SS, Asghar R. 2007. Hymenomycetes from Multan District. *Pakistan J Bot* 39: 651-657.
27. Martin E, Coronades M, Sanchez A, Perez-Silva E, Herrera T. 2005. Macromycetes of Pinacate and Great Altar desert biosphere, Sonoron, Mexico. *Mycotaxon* 95: 31-90.
28. Walley R, Rammeloo J. 1994. The poisonous and useful fungi of Africa south of the Sahara: a literature survey. *Scripta Bot Belgica* 10: 1-56.