

A STUDY ON THE PATHOLOGICAL AND DIAGNOSIS OF EIMERIA SPECIES INFECTION IN JAPANESE QUAIL

N.H. Mohammad

Department of microbiology, College of Veterinary Medicine, University of Mosul,

Mosul, Iraq

(Received 20 March 2012 ,Accepted 24 June 2012)

Keywords: *Eimeria* spp, Japanese quails

ABSTRACT

This study was conducted to detect the intestinal coccidial types in 87 Japanese quails (*coturnix coturnix japonica*) the birds were purchased from different parts of Nenevah governorate. The current study reveals that the total percentage of infection with *Eimeria* spp. oocysts 49.4%. Three species were diagnosed, according to their percentage they are: *E. tsunodai* 44.8%, *E. uzura* 34.5%, *E. bateri* 24.1%. The higher rates were in young birds. There was no significant difference with respect to sex. The results showed the higher infection rate and intensity was recorded in caecum infection with three species of *Eimeria* was most frequent in Japanese quails with percentage 46.5%. Macroscopical lesions of infected birds revealed that softening of feces, thickening of mucosa and hemorrhage in caecum. Microscopically lesions characterized by severe hyperplasia of epithelial cell with construction of intestinal gland cavities in small intestine, caecum and presence of edema between muscle fibers with different development epithelial cells and infiltration with inflammatory cells. This study is regarded the first study in detection of species and pathological effect of *Eimeria* in Japanese quails in Nenevah governorate.

INTRODUCTION

Coccidiosis is one of the most important and common protozoal disease in various avian species (1). This parasitic infection occurs in the epithelial cells of the intestine, it results in a greateconomic loss all over the world (2). Quails which consider a branch of the modern poultry industry. These birds are raised primarily for production of eggs, meat and used as laboratory animals similar to rats and mice (3, 4). Quails are most susceptible to various diseases such as coccidiosis which recognized as a serious parasitic disease problem limiting quail industry(5). Clinically coccidiosis is characterized by diarrhea loss of appetites and a general appearance of malaise, weight loss, watery diarrhea that may be greenish or bloody and retarded growth (6). In some cases a number of quails died at 5-6 day occurred in experimental study(7). Various species of *Eimeria* I been isolated from the different species of quails such as *E.tsunodai*, *E.uzura*, *E.bateri* from Japanese quails (6) and *E.lophortygis*, *E.okanaganensis* described from California quails while *E.crusti*, *E.oreortygis* are described from mountain quail(8) and *E.conturnicis*, *E.bateri* are described from grey quail(4) and *E.colini*, *E.lettyae* from bob white quail(9) also *E.tahamensis* from Arabian quail(10). The present study was designed to provide preliminary information on the prevalence rates, type of *Eimeria* with describing both gross and microscopic changes caused by these parasites in Japanese quails in Nenevah governorate, Iraq.

MATERIALS AND METHODS

This study was conducted on 87 Japanese quails (*coturnix coturnix japonica*) and were classified as young quails (less than 10 weeks) and adult (more than 10 weeks) from both sexes (45 females, 42 male) after slaughtered the Japanese quails fecal samples were collected directly from the duodenum, small intestine and caecum. direct

smear method(11) and sheath flotation technique were used for diagnosis of oocysts of *Eimeria* spp(12). Identified according to the site of infection, size of oocysts and sporocysts, presence or absence of micropyle(13, 14). according to (15) for detection the degree of intensity of infection with *Eimeria* spp. wet smears of mucosa were prepared from duodenum and small intestine and caecum scraping stained with Giemsa for microscopic examination of *Eimeria* spp. stage(16), the gross characteristics of the lesion were described and recorded. Selectively tissue specimens were collected, preserved in 10% formalin solution and processed by histopathological techniques and stained with haematoxylin and eosin and examined histologically for the presence of *Eimeria* spp. stage and pathological changes(17). The data were analyzed statistically by using chi-square(18).

RESULTS

Out of the 87 Japanese quails examined during the study period, 43 cases (49.4%) were positive for *Eimeria* spp. Three *Eimeria* spp. were identified in naturally infected birds belonged to *Eimeria tsunodai*, *Eimeria uzura*, *Eimeria bateri* (figure 1,2,3). Differentiation of *Eimeria* sp. was based on morphological specific feature and by microscopic measurements by using ocular micrometer (tables 1,2). The prevalence of *Eimeria* spp. and site of infection recovered from Japanese quails are given in (table 3). A higher percentage was reported in *Eimeria tsunodai* 44.8%. Statistical analysis showed significant differences between *Eimeria* sp. from (table 4) it is evident that there were three degree of infection according to the site of infection and intensity of infection with *Eimeria* sp. The highest rate of infection was found in young quails and lowest percentage in adult. Statistical analysis showed significant difference between the ages. While no significant

difference between male and female (table 5). The results reveal that high percentage of infection with three species of *Eimeria*(mixed infection) was 46.5% while single infection the lowest was 20.9%,with significant difference was observed between them(table 6) and presence of development stage of coccidia in scrupling stained with Giemsa stain(figure 4) this study including present of macroscopical pathological lesions represented by softening of feces in duodenum ,small intestine and more clearance in caecum,with thickening of mucosa and light hemorrhage in caecum while the histopathological lesion characterized by sever hyperplasia of epithelial cells with construction of intestinal gland cavities in small intestine and caecum (figure 5),with presence of developmental stage of parasite in epithelial layer lining of intestinal glands(figure6,7),infiltration with inflammatory cellsrepresented by esinophils and presence of edema between the musclefibers in small intestine and caecum (figure 8).

Table(1): The dimensions of the oocysts and morphological of *Eimeriaspp*determined in Japanese quail.

| Species | Oocyst size(μ) | | Morphology | Wall | Micropyle |
|-------------------|------------------------------|--------------------|------------------|--------|-----------|
| | mean \pm StE. | Range | | | |
| <i>E.tsunodai</i> | 19-14.6 0.88 \pm 0.76 | (16-20)(18-14.5) | Ovoid | Double | - |
| <i>E.uzura</i> | 23.4-17.7 2.02 \pm 0.61 | (19-28.5)(21.5-16) | Broaid ellipsoid | Double | + |
| <i>E.bateri</i> | 22-16.4 1.05 \pm 0.89 | (16-30) (21-14.5) | subspherical | Double | - |

*(10-15)of the oocysts measured from each species

Table(2):The dimensions of the *Eimeria* sporocyst detected in the Japanese quails.

| Species | Sporocyst size(μ) | |
|--------------------|------------------------------|---------------------|
| | mean \pm StE. | Range |
| <i>E. tsunodai</i> | (5.1-10.5) 0.4 \pm 0.83 | (10.7-9.1)(5.3-4.7) |
| <i>E. uzura</i> | (6-11.5) 1.2 \pm 0.8 | (12-10.8)(6-5) |
| <i>E. bateri</i> | (6.8-10.2) 0.5 \pm 0.9 | (11.2-8.5)(7-6) |

*(8-10)of the sporocyst measured from each species

Table (3):Prevalenceand the site of infection with *Eimeria* spp in Japanese quails

| <i>Eimeria</i> species | No.of positive | Rate of infection | Site of infection | | |
|------------------------|----------------|-------------------|-------------------|-----------------|--------|
| | | | duodenum | Small intestine | Caecum |
| <i>E.tsunodai</i> | 39 | 44.8 ^a | - | - | - |
| <i>E.uzura</i> | 30 | 34.5 ^b | + | + | + |
| <i>E.bateri</i> | 21 | 24.1 ^c | + | + | + |

*different letters have significant differences at $P \leq 0.05$

Table(4): Distribution and intensity of infection with *Eimeria* spp. according to site of infection

| Organs | No.of positive sample | Rate of infection | Intensity of infection |
|-----------------|-----------------------|---------------------|------------------------------|
| Duodenum | 6 | 6.9% ^a | Lowdegree |
| Small intestine | 9 | 10.3% ^a | Moderatedegree low degree |
| Caecum | 28 | 32.9 % ^b | High degree low degree |

*rate with different letters have significant difference at $p \leq 0.05$

Lowdegree: 5 oocyt/hpf.,Moderate degree: 10-15 oocyst/hpf.,High degree:50 oocyst/hpf.

Table(5):Relationship of *Eimeria* infection with sex and age in Japanese quails.

| Sex | Female | | | Male | | |
|--------------|----------------------|--------------|--------------------|----------------------|--------------|--------------------|
| | No.of birds examined | No.+ve birds | Rate of infection | No.of birds examined | No.+ve birds | Rate of infection |
| Age of birds | | | | | | |
| Young birds | 27 | 16 | 59.3% ^a | 26 | 16 | 61.5% ^a |
| Adult birds | 18 | 6 | 33.3% ^b | 16 | 5 | 31.3% ^b |
| Total | 45 | 22 | 48.9% ^a | 42 | 21 | 50% ^b |

Vertical different small letters have significant difference at $P \leq 0.05$.

Horizontal different large letters no significant difference at $P \leq 0.05$.

Table(6): Type of infection with *Eimeria* spp.in Japanese quails.

| Type of infection | No.+ve birds | rate of infection |
|-------------------|--------------|-------------------|
| Single infection | 9 | 20.9% a |
| Double infection | 14 | 32.6% b |
| Mixed infection | 20 | 46.5% c |
| Total | 43 | 100 |

Different letters have significant differences at $P \leq 0.05$.




Figure 1: *Eimeria tsunodai* oocyst by using direct smear method x400




Figure 2:*Eimeria uzura* oocyst by using flotation technique x400.




Figure 3:*Eimeria bateri* oocyst by using direct smear method.x400.




Figure 4: Caecum of quails showing presence of development stage of *Eimeria* spp in scrapling stained with Gimsas stain, 1000X.




Figure 5: Histological section of small intestine showing hyperplasia of epithelial cells (a), with construction of intestinal gland cavities (b). H&E X370.




Figure 6: Histological section of small intestine showing presence of development stage of *Eimeria* spp .in the epithelial cells (a).H&E X370.




Figure 7: Histological section of small intestine showing presence of *Eimeria* spp.in various stages of development in epithelial cells (a), H&EX370.




Figure 8: Histological section of caecum showing edema between the muscle fibers (a). H&E X370.

DISCUSSION

Quails considered a branch of the modern poultry industry in Iraq. Several *Eimeria* spp are highly pathogenic to their host causing great economic losses in quail breeding and limiting development of this industry (19). The present study described the incidence, morphological and pathological characters of infection with *Eimeria* spin slaughtered Japanese quails examined fecal samples collected from intestinal tract revealed that 43(49.4%) were infected with *Eimeria* spp. This result is apparently similar to that obtained by (20) who detected *Eimeria* infection in Azerbaijan 52% of examined Japanese quails. But lower than study in the Okanagan valley of British Columbia (21) and higher than study in Saudi Arabia (13) the difference in the percentage of infection in many studies may be related to different factors such as, environmental conditions, seasonal fluctuations, type of anticoccidial drugs may

have contributed to this difference(22). In this study three *Eimeria* were diagnosed naturally infected Japanesequails(*Eimeria tsunodai*,*Eimeria bateri*,*Eimeria uzura*). These results are in agreement with reports of 6,13. The measurement oocysts and sporocysts which observed in this study are in agreement with those described by (4,6,13,and 23). There was significant with different in prevalence between species of *Eimeria*. This might be due to immunity status of the host, stress factor, environmental conditions and misuse of coccidiostats or the development of local strain of *Eimeria* spp to variable compounds(24), the infection of *Eimeria* spp in intestinal tract showed that no significant differences *between* the infection of duodenum ,small intestine. While significant was noticed between the infection of caecum and the duodenum ,small intestine the cause of this difference might be due to high prevalence rate of *E.tsunodai* appeared in caecum(7) refers to the pathogenic of this species of *Eimeria* is more effect than other species and expressed the pathogenic and clinical symptoms of *Eimeria tsunodai* resemble to those of chickens infected with *E .tenella*, the intensity of infection with *Eimeria* of Japanese quails revealed that high degree of infection appeared in caecum the result was in agreement with (15,25). According to the age, sex the infection rate of *Eimeria* spp was low in adult while it was high in young quails. These results not agreement with studies carried out on quails in California(21) noted that the prevalence of coccidiosis increased with the age of quails. While agreement with (26) reported that young birds are more susceptible to coccidiosis effects and it is suggested that older birds evidentially develop some kind of immunity to infection. On other hand no significant occurred in the incidence of infection between the male and female this result agreement with experiences of (21). The mixed infection with three species of *Eimeria* recorded the highest rates this result is in agreement with experiences of (22). While not agreement with (27) there was

asignificant differences was noticed between mixed infections, double and single infection .this might be due to many different factors such as farm management practices, oocyst contamination of food and drinking water, differences in management of the anticoccidial programs (28). The histopathological changes observed in duodenum, small intestine, caecum were more or less similar to those reported in quails by (5,6,7,29). The histopathological changes in intestinal tract pointed to the serious effect of *Eimeria* sp. in quails. This observation has an important value since the pathogenicity of *Eimeria* spp in quails had not been discussed.

ACKNOWLEDGEMENT

This study was supported by the College of Veterinary Medicine, University of Mosul.

دراسة تشخيصيه وإمراضية للاصابة بانواع طفيلي *EIMERIA* في طائر السمان الياباني

نادية حامد محمد

فرع الاحياء المجهرية، كلية الطب البيطري، جامعة الموصل، الموصل، العراق

الخلاصة

تضمنت الدراسة الكشف عن انواع الاكريات المعوية لـ 87 من طائر السمان الياباني تم الحصول عليها من مختلف مناطق محافظة نينوى. اشارت الدراسة الحالية الى ان نسبة الخمج الكليه بداء الاكريات 49.4% وتم تشخيص ثلاثة انواع من *Eimeria* وهي *E. tsunoda* 44.8%، *E. uzura* 34.5% و *E. bateri* 24.1%. وبينت الدراسة ان اعلى نسبة خمج كانت في الفئة العمرية الصغيرة ولم يلاحظ وجود فرق معنوي في نسبه الخمج بين الجنسين. ولوحظ ان اعلى نسبة خمج وشدته سجلت في الاعور. ان خمج طائر السمان الياباني بثلاثة انواع من الايميريا هو النمط الاكثر شيوعا وبنسبة 46.5%. اظهرت نتائج الدراسة الحالية وجود تغيرات مرضية عيانية تمثلت بوجود تلين البراز في كل من الاثني عشر والامعاء الدقيقة وكاناكثر وضوحا في الاعور مع وجود نتخن في بطانة الامعاء فضلا عن النزف الخفيف في الاعور. واطهر الفحص النسجي وجود تغيرات مرضية

تمثلت بفرط التنسج في ظهارة الامعاء والاعور مع وجود الودمة بين الالياف العضلية مع المراحل التطورية للطفيلي وارتشاح للخلايا الالتهابية. وتعتبر الدراسة الاولى من نوعها في الكشف عن انواع الاكربيات وتأثيرها المرضي في طائر السمان الياباني في محافظة نينوى.

REFERENCES

- 1-Al-natour M Q, Suleiman M. Flock-level prevalence of *Eimeria* spp. among broiler chicks in northern Jordan. *Pre Vet Med*,2002;53:305-310.
- 2-Radi Z A. An epizootic of combined *Clostridium perfringens*, *Eimeria* spp. and *Capillaria* spp. Enteritis and *Histomonas* spp. hepatitis with *Eischerichia coli* septicemia in bob white quails (*colinus virginianus*). *Int J Poult Sci*,2004;3(7):438-441.
- 3-Sreeranjini A R, Lyyangar MP, Pramod kumar D. Histological study on the fibrous architecture of kidney and ureter of Japanese quail (*coturnix coturnix japonica*). *Tam J Vet Ani Sci*,2010;6(2):107-110.
- 4-Tsutsumi Y. *Eimeria tsunodai* Spp. nov. (Protozoa: Eimeriidae) acecal coccidia of Japanese quails (*coturnix coturnix japonica*). *Jap J Vet Sci*,1972;34:1-9.
- 5-Seok S, Park J, Cho S, Baek M, Lee H, Kim D. coccidian (*Eimeria* spp) in small intestine of Japanese quail (*coturnix coturnix japonica*). *Kor J Lab Ani Sci*,2003;19(2):90-91.
- 6- Teixeira M, Teixeira fwl, Lopes CWG. Coccidiosis in Japanese quails (*coturnix japonica*) characterization of a naturally occurring infection in a commercial rearing farm. *Rev Bras Cienc Avic*,2004;6:129-134.
- 7-Tsutsumi Y. Pathogenicity of *Eimeria tsunodai* for Japanese quails (*coturnix coturnix japonica*) and susceptibility of the coccidium to some drugs. *Jap J Vet Sci*, 1972;34:115 – 121.

- 8-Duszynski DW, Gutierrez RJ. The coccidia of quail in the United States. *J Wild Dis*, 1981;17:371 – 379.
- 9-Ruff MD. Life cycle and biology of *Eimeria lettyae* sp.(Protozoai: *Eimeriidae*) from the northern bob white, *colinus virginianus*. *J Wild Dis*, 1985;21:361 – 370.
- 10-Amoudi MA. *Eimeria tahamensis*. n.spp (Apicomplexa *Eimeriidea*) from the Arabian quail (*coturnix delegorguei arabica*). *J Protozool*, 1987;34:455-456.
- 11-Urquhart G M, Armour J, Duncan JL, Dunn AM, Jennings FW. *Veterinary parasitology*. 2nd ed. black well science Ltd. Oxford, 2003. 276p.
- 12-Foreyt WJ. *Veterinary parasitology. Reference manual*. 5th ed. Iowa State University Press USA. Zool. pp: 5 – 9.
- 13-Bashtar AR, Abdel-Ghaffar F, Al-Rasheid KAS, Mehlhorn H, Al Nasr I. Light microscopic study on *Eimeria* species infecting Japanese quails reared in Saudi Arabian farms. *Para Res*, 2010;107:409-416.
- 14-Tsunodai K, Muraki Y. New coccidium of the Japanese quail *Eimeria uzura* sp.nov. *Jap Vet Sci*, 1971;33:227-235.
- 15-AL-Alousi TI, Daoud MS, AL.Bayati MM. A study of endoparasites of turkeys in Mosul –Iraq. *Iraqi J Vet Sci*, 1994;7(3):123-129.
- 16-Soulsby E JL. *Helminths, Arthropods and protozoa of domesticated animals*. 7th ed. London: Philadelphia Bailliere, Tindal, 1982. pp: 777.
- 17-Luna LG. *Manual of histological staining methods of the Armed forces institute of pathology*, 3rd edition, the blakiston division, McGraw-Hill book company, Newyork, 1968 pp: 34, 127.
- 18-Petrie A, Watson p. *Statistics for veterinary and animal science*. Black Well Science. London, 2003.

- 19-Abdel-Ghaffar f A,Bashtar AR,Mohamed A. Incidence and prevalence of *Eimeria* infections among turkeys in Egypt. J Egy Ger Soe Zool,1990;1:125-134.
- 20-Musaev M,Gaibova G, Ismailova G,Alieva F,Iskenderova N. The coccidia of the gallinaceous birds in Azerbaijan. Tr J Vet Anim Sci,1998;22:409-413.
- 21-Liburd E M. Incidence of coccidia in California quail(*lophortyx californicus*)from the Okanagan valley,British Columbia.Can JZool.1969;47(4):645-648.
- 22-Nematollahi A, Moghaddam Gh, Niyazpour F. Prevalence of *Eimeria sp.*among broiler chicks in Tabriz (northwest of Iran). Res JPoult Sci,2008;2(3):72-74.
- 23-Norton CC, Peirce MA. The life cycle of *Eimeria bateri*(protozoa:Eimeriidae)in the Japanese quail *coturnix coturnix japonica*. JProtozool, 1971;18:57-62.
- 24-Kaya G. Prevalence of *Eimeria* species in lambs in Antakyaprovince.Turk J Vet Anim Sci, 2004;28:687-692.
- 25-Al-Tae A F M. Incidence of broiler caecal coccidiosis in Ninevah governorate 1999-2004.Iraq J Vet Sci, 2007;21(1):45-63.
- 26-Mcdougald LR, Fuller L,Mattiello R.Asurvey of coccidia on 43 poultry farms in Argentina. Avian Dis,1997;41:923-929.
- 27-Al-Neema MS.Diagnostic study on *Eimeria*species in chickens in Al-Hamdania region. [Master's thesis]. College of VeterinaryMedicine, University of Mosul;2011.
- 28-Sulaiman EG,Talib Q,Daham E,Arsalan S H. Study of some eggs and oocysts of internal parasites in sheep in Mosul. Iraqi J Vet Sci, 2005;19(1):21-32
- 29-Ruff MD,Fagan JM,Dick JW. Pathogenicity of coccidians in the Japanese quail (*coturnix coturnix japonica*).Poult Sci,1984;63:55-60.