

Fuzzy Linear Transformations **التحويلات الخطية الضبابية**

Dr. Ali H. Battor and Areej Tawfeeq Hameed AL–Budarub
Assistant Professor , Department of Mathematics,
College of Education for Girls, University of Kufa

ABSTRACT

In this paper , the concept of fuzzy linear transformation have been investigated , this lead us to study and give some properties concerning with it .

Moreover , we give some types of fuzzy as a fuzzy Kernel and its relationships with fuzzy linear transformation and a characterizations of fuzzy linear transformation is presented .

المستخلص

في هذا البحث قدمنا التحويلات الخطية الضبابية كتعميم للتحويلات الخطية الاعتيادية , والذي قادنا الى دراسة واعطاء العديد من الخواص المتعلقة بهذا المفهوم .
كذلك برهنا النتائج الاساسية المناظرة للمفهوم الاعتيادي , بالاضافة الى ذلك اعطينا بعض المفاهيم الضبابية الاخرى مثل النواة وغيرها لبيان مدى علاقتها بالتحويلات الخطية الضبابية .

INTRODUCTION

The present paper introduces and studies fuzzy linear transformation . In fact , some basic definitions and results which will be needed later are recalled.

In section one , we applies the concept of fuzzy set on a vector space and we give some of properties , a binary operations addition and scalar multiplication . Finally , we studied and debated some properties that are necessary in this work .

In section two , we studied and discusses the concept of fuzzy linear transformation on vector space , a binary operations addition and scalar multiplication and some theorems we studied and discusses the concept of a fuzzy kernel on a vector space .

In section three , we shall give definition and some properties of fuzzy coset and we shall give definition and some properties of quotient fuzzy ring . Moreover , we studied the concept of a fuzzy isomorphism .

Throughout this paper $(R,+, \cdot)$ be a commutative ring with identity .

PRELIMINARY CONCEPTS

In this section some basic definitions and results which we will be used in the next section are considered .

Let X be a nonempty set , **A fuzzy subset of X** is a function from X into $[0,1]$, ([7] , [2]) .

Let A and B be fuzzy subset of X . If for all $x \in X$, $A(x) \leq B(x)$ then we write $A \subseteq B$. If $A \subseteq B$ and there exists $x \in X$ such that $A(x) < B(x)$, then we write $A \subset B$ and we say that A is a proper fuzzy subset of B , [7] . Note that $A = B$ if and only if $A(x) = B(x)$, for all $x \in X$, ([7] , [8]) .

Let λ_X denote the characteristic function of X defined by $\lambda_X(x) = 1$ if $x \in X$ and $\lambda_X(x) = 0$ if $x \notin X$, ([8] , [1]) .

Let $(R,+, \cdot)$ be a commutative ring with identity , for each $t \in [0,1]$, the set $A_t = \{ x \in R \mid A(x) \geq t \}$ is called **a level subset of R** and $A = B$ if and only if $A_t = B_t$ the set $A_* = \{ x \in R \mid A(x) > 0 \}$ is called **the support of R** , ([2] , [1]) .

Let $x \in X$ and $t \in [0,1]$, let x_t denote the fuzzy subset of X defined by $x_t(y) = 0$ if $x \neq y$ and $x_t(y) = t$ if $x = y$ for all $y \in R$. x_t is called a **fuzzy singleton**, ([1],[8]). If x_t and y_s are fuzzy singletons, then $x_t + y_s = (x + y)_\lambda$ and $x_t \circ y_s = (x \cdot y)_\lambda$, where $\lambda = \min \{ t, s \}$, ([7],[4],[8]).

Let $I^R = \{A_i \mid i \in \Lambda\}$ be a collection of fuzzy subset of R . Define the fuzzy subset of R (**intersection**) by $(\bigcap_{i \in \Lambda} A_i)(x) = \inf \{ A_i(x) \mid i \in \Lambda \}$ for all $x \in R$, ([2],[4]). Define the fuzzy

subset of R (**union**) by $(\bigcup_{i \in \Lambda} A_i)(x) = \sup \{ A_i(x) \mid i \in \Lambda \}$ for all $x \in R$, ([3],[4]).

The empty fuzzy subset of R denote by ϕ is definition by : $\phi(x) = 0$ for all $x \in R$, ([7],[4]).

Let A and B be fuzzy subsets of R , **the product $A \circ B$** define by : $A \circ B(x) = \sup \{ A(y), B(z) \mid x = y \cdot z \}$ $y, z \in R$, for all $x \in R$, ([2],[3]). And **the addition $A+B$** define by : $A+B(x) = \sup \{ A(y), B(z) \mid x = y + z \}$ $y, z \in R$, for all $x \in R$, ([4],[2]).

The complement of A denoted by $E = A^c$ and define by $E(x) = A^c(x) = 1 - A(x)$, for all $x \in R$, [7].

When we say fuzzy subset we mean a non empty fuzzy subset. We let $\text{Im}(A)$ denotes **the image of A** . We say that A is a **finite -valued** if $\text{Im}(A)$ is finite and $|\text{Im}(A)|$ denotes the cardinality of $\text{Im}(A)$, ([2],[8]).

Let $f : X \rightarrow Y$, A and B are two nonempty fuzzy subsets of nonempty sets X and Y respectively, the fuzzy subset $f(A)$ of Y defined by : $f(A)(y) = \sup A(x)$ if $x \in f^{-1}(y) \neq \emptyset$, $y \in Y$ and $f(A)(y) = 0$, otherwise, where $f^{-1}(y) = \{x : f(x) = y\}$. It is called **the image of A under f** and denoted by $f(A)$. The fuzzy subset $f^{-1}(B)$ of R defined by : $f^{-1}(B)(x) = B(f(x))$, for $x \in X$. (i.e. $f^{-1}(B) = (B \circ f)$). Is called **the inverse image of B** and denoted by $f^{-1}(B)$, [2]. A fuzzy subset A of X is called **f-invariant** if $f(x) = f(y)$ implies $A(x) = A(y)$, where $x, y \in X$, [8]. A is called **the sup property**, if every set of $\text{Im}(A)$, the image of A has a maximal element, ([8],[4]).

Let X be a nonempty set and a fuzzy set A in X can be represented by the set of pairs : $A = \{(x, A(x)) : x \in X\}$. **The family of all fuzzy sets in X** is denoted by I^X ([7],[2]).

Let A be a non empty fuzzy subset of a group G , A is called a **fuzzy subgroup of G** if for all $x, y \in G$, $A(x + y) \geq \min \{A(x), A(y)\}$ and $A(x) = A(-x)$, ([8],[8],[4]).

A is a non empty fuzzy subset of R , A is called a **fuzzy ring of R** if and only if for all $x, y \in R$, then $A(x - y) \geq \min \{A(x), A(y)\}$ and $A(x \cdot y) \geq \min \{A(x), A(y)\}$, ([8],[2]).

A non empty fuzzy subset A of R is called a **fuzzy ideal of R** if and only if for all $x, y \in R$, then $A(x - y) \geq \min \{A(x), A(y)\}$ and $A(x \cdot y) \geq \max \{A(x), A(y)\}$, ([4],[2]). It is clear that every fuzzy ideal of R is a fuzzy ring of R , but the converse is not true.

SECTION ONE

Fuzzy Vector Space

In this section, we applies the concept of fuzzy set on vector space and we give some of properties, a binary operations addition and scalar multiplication. Finally, we studied and debated some properties that are necessary in this work.

DEFINITION 1.1 [3]:

A vector space over a field F is a set X , whose elements are called vectors which two operations, addition $(+ : X \times X \rightarrow X)$ and scalar multiplication $(\cdot : F \times X \rightarrow X)$ with conditions is satisfies :-

1. $x + y \in X$, for all $x, y \in X$;
2. $x + y = y + x$, for all $x, y \in X$;
3. $x + (y + z) = (x + y) + z$, for all $x, y, z \in X$;
4. There exists $0 \in X$ such that $0 + x = x$, for all $x \in X$ and 0 is the zero vector or the origin;
5. For all $x \in X$, there is a unique element $(-x) \in X$ such that $x + (-x) = 0$;
6. $\lambda x \in X$, for $\lambda \in F$ and for all $x \in X$;
7. $\lambda(x + y) = \lambda x + \lambda y$, for $\lambda \in F$ and $x, y \in X$;

8. $(\lambda + \alpha)x = \lambda x + \alpha x$, for $\lambda, \alpha \in F$ and $x \in X$;
9. $(\lambda \alpha)x = \lambda(\alpha x)$, for $\lambda, \alpha \in F$ and $x \in X$;
10. $I.x = x .I = x$, for all $x \in X$ and I is the unity element of the field F .

DEFINITION 1.2 [2]:

If X be a vector space over F and $A, B \subseteq X, G \subseteq F$, the following notations will be used :
 $A + B = \{ x = a + b : a \in A, b \in B \}$ and $GA = \{ x = \lambda a : a \in A, \lambda \in G \}$.

DEFINITION 1.3 [15]:

If A, B are fuzzy sets in vector space X over F and let $\lambda \in X, G \subseteq F$. We define $A + B$ and λA by :-

1. $A + B = f(A \times B)$, where $A \times B(x, y) = \min \{ A(x), B(y) \}$ and $f: X \times X \rightarrow X$ is a function defined by $f(x, y) = (x + y)$, for all $x, y \in X$.
2. $\lambda A = g(A)$ where $g: X \rightarrow X$ is a function defined by $g(x) = \lambda x$, for all $x \in X$.

DEFINITION 1.4 [11]:

A fuzzy subset A of a field F is a **fuzzy field of F** if

1. $A(1) = 1$.
2. $A(x-y) \geq \min \{ A(x), A(y) \}$, for each $x, y \in F$.
3. $A(xy^{-1}) \geq \min \{ A(x), A(y) \}$, for each $x, y \in F, y \neq 0$.

Let A be a fuzzy field of F . If $x \in F, x \neq 0$, then $A(0) = A(1) \geq A(x) = A(-x) = A(x^{-1})$.

• **DEFINITION 1.5 [15]:**

Let X be a vector space over F . A fuzzy set A in X is called a **fuzzy subspace over F** if :

1. $A + A \subseteq A$;
2. $\lambda A \subseteq A$, for all $\lambda \in F$.

• **DEFINITION 1.6 [11]:**

A is a fuzzy set of a vector space V over a field F . A is a **fuzzy subspace of V over a fuzzy subfield K of F** if :

1. $A(0) > 0$;
2. $A(x-y) \geq \min \{ A(x), A(y) \}$, for all $x, y \in V$;
3. $A(cx) \geq \min \{ K(c), A(x) \}$, for all $c \in F$, for all $x \in V$.

THEROEM 1.7 [8]:

Let A be a fuzzy set in a vector space X over F , then the following statements are equivalent :

1. A is a fuzzy subspace of X .
2. For all $\alpha, \beta \in F$, we have $\alpha A + \beta A \subseteq A$.
3. For all $\alpha, \beta \in F$ and for all $x, y \in X$, we have $A(\alpha x + \beta y) \geq \min \{ A(x), A(y) \}$.

PROPOSITION 1.8 [16]:

1. If A is a fuzzy subspace of vector space X over F . Then $A(0) > A(x)$, for all $x \in X$.
2. If A is a fuzzy set in a vector space X over F . Then A is a fuzzy subspace of X if and only if A_t is a subspace of X , for all $0 \leq t \leq A(0)$.

PROPOSITION 1.9 [15]:

1. If A, B are fuzzy subspaces of vector space X over F and $\lambda \in F$. Then $\lambda A, A + B, A \cap B$ are fuzzy subspaces in X .

2. Let $x, y \in X$ and A be a fuzzy set of a vector space X over F such that $A(x) > A(y)$, then $A(x + y) = A(y)$.

3. If A is a fuzzy subspace of vector space X over F and $x, y \in X$ with $A(x) \neq A(y)$, then $A(x + y) = \min \{ A(x), A(y) \}$.

PROPOSITION 1.9 [15]:

Let X, Y be two vector spaces over F and let $f: X \rightarrow Y$, be a linear function . Then

1. If A is a fuzzy subspace in X , then $f(A)$ is a fuzzy subspace in Y .
2. If B is a fuzzy subspace in Y , then $f^{-1}(B)$ is a fuzzy subspace in X .

SECTION TWO

Fuzzy Linear Transformations

In this section , we studied and discusses the concept of fuzzy linear transformation on vector space , a binary operations addition and scalar multiplication and citation some theorems . Finally , we studied and debated some properties that are necessary in this work .

DEFINITION 2.1 [3]:

Let X , Y be two vector spaces over F and let $f : X \rightarrow Y$ is called a linear transformation on a vector space if :-

1. $f(x + y) = f(x) + f(y)$, for all $x , y \in X$;
2. $f(\lambda x) = \lambda f(x)$, for all $x \in X$ and $\lambda \in F$.

Or $f(\lambda x + \alpha y) = \lambda f(x) + \alpha f(y)$, for all $x , y \in X$ and $\lambda , \alpha \in F$.

The linear transformation $f : X \rightarrow Y$ is called a linear functional on X .

DEFINITION 2.2 :

Let A , B be two fuzzy subspaces of vector spaces X , Y over F respectively . $K : A \rightarrow B$ is called a fuzzy linear transformation on a fuzzy subspace if :-

$$K(\lambda x_t + \alpha y_h) \geq \min \{ K(x_t) , K(y_h) \} , \text{ for all } x_t , y_h \in A , \lambda , \alpha \in F \text{ and } t , h \in [0,1] .$$

The fuzzy linear transformation $K : A \rightarrow B$ is called a fuzzy linear functional on A .

EXAMPLES 2.3 :

1. Let A be a fuzzy subset of R^3 such that $A(a , b , c) = 1$ for all $(a , b , c) \in R^3$ and B be a fuzzy subset of R^2 such that $B(a , b) = 1/2$ for all $(a , b) \in R^2$. $K : A \rightarrow B$ such that $K(a , b , c) = (a , b)$, for all $(a , b , c) \in R^3$. Is K a fuzzy linear transformation on a fuzzy subspace A .

Solution :

To prove A and B are two fuzzy subspaces of vector spaces R^3 and R^2 respectively .

Let $x , y \in R^3$, $\alpha , \beta \in F$ (F is a field) , then $A(\alpha x + \beta y) \geq \min \{ A(x) , A(y) \}$, where $x = (a_1 , b_1 , c_1)$, $y = (a_2 , b_2 , c_2)$.

$$\begin{aligned} A(\alpha x + \beta y) &= \sup \{ \min \{ A(u) , A(w) \} \mid u + w = \alpha x + \beta y , u , w \in R^3 \} . \\ &= \sup \{ \min \{ A(\alpha x) , A(\beta y) \} \} , \\ &= \sup \{ \min \{ \sup \{ \min \{ A(\alpha) , A(x) \} \} , \sup \{ \min \{ A(\beta) , A(y) \} \} \} \} , \\ &= \sup \{ \min \{ A(\alpha) , A(x) , A(\beta) , A(y) \} \} , [4] , \\ &\geq \sup \{ \min \{ A(x) , A(y) \} \} , \\ &\geq \min \{ A(x) , A(y) \} . \end{aligned}$$

A and B are two fuzzy subspaces of vector spaces R^3 and R^2 respectively .

$K : A \rightarrow B$ such that $K(a , b , c) = (a , b)$, for all $(a , b , c) \in R^3$.

$$\begin{aligned} K(\alpha x + \beta y) &= \sup \{ \min \{ K(\alpha x) , K(\beta y) \} \} , \text{ for all } x , y \in R^3 . \\ &= \sup \{ \min \{ K(s) , K(d) \} \mid s = \alpha x , d = \beta y \} , \text{ for all } s , d \in R^3 . \\ &= \sup \{ \min \{ K(s) , K(d) \} \mid s = (\alpha a_1 , \alpha b_1 , \alpha c_1) , d = (\beta a_2 , \beta b_2 , \beta c_2) \} , \\ &\geq \sup \{ \min \{ K(x) , K(y) \} \} , \\ &\geq \min \{ K(a) , K(b) \} , \end{aligned}$$

Hence K is a fuzzy linear transformation on a fuzzy subspace A .

2. Let A be a fuzzy subset of R^2 such that $A(a , b) = 1/3$ for all $(a , b) \in R^2$ and B be a fuzzy subset of R such that $B(a) = 1/4$ for all $a \in R$. $K : A \rightarrow B$ such that $K(a , b) = a$, for all $(a , b) \in R^2$. Is K a fuzzy linear transformation on a fuzzy subspace A .

Solution :

To prove A and B are two fuzzy subspaces of vector spaces R^2 and R respectively .

Let $x , y \in R^2$, $\alpha , \beta \in F$ (F is a field) , then $A(\alpha x + \beta y) \geq \min \{ A(x) , A(y) \}$, where $x = (a_1 , b_1)$, $y = (a_2 , b_2)$.

$$\begin{aligned} A(\alpha x + \beta y) &= \sup \{ \min \{ A(u) , A(w) \} \mid u + w = \alpha x + \beta y , u , w \in R^2 \} . \\ &= \sup \{ \min \{ A(\alpha x) , A(\beta y) \} \} , \\ &= \sup \{ \min \{ \sup \{ \min \{ A(\alpha) , A(x) \} \} , \sup \{ \min \{ A(\beta) , A(y) \} \} \} \} , \end{aligned}$$

$$\begin{aligned}
 &= \sup \{ \min \{ A(\alpha) , A(x) , A(\beta) , A(y) \} \} , [4] , \\
 &\geq \sup \{ \min \{ A(x) , A(y) \} \} , \\
 &\geq \min \{ A(x) , A(y) \} .
 \end{aligned}$$

A and B are two fuzzy subspaces of vector spaces R^2 and R respectively .

$K : A \rightarrow B$ such that $K(a, b) = a$, for all $(a, b) \in R^2$.

$$\begin{aligned}
 K(\alpha x + \beta y) &= \sup \{ \min \{ K(\alpha x) , K(\beta y) \} \} , \text{ for all } x, y \in R^2 . \\
 &= \sup \{ \min \{ K(s) , K(d) \} \mid s = \alpha x , d = \beta y \} , \text{ for all } s, d \in R^2 . \\
 &= \sup \{ \min \{ K(s) , K(d) \} \mid s = (\alpha a_1 , \alpha b_1) , d = (\beta a_2 , \beta b_2) \} , \\
 &\geq \sup \{ \min \{ K(x) , K(y) \} \} , \\
 &\geq \min \{ K(a) , K(b) \} ,
 \end{aligned}$$

Hence K is a fuzzy linear transformation on a fuzzy subspace A .

REMARK 2.4 :

1. Let A , B be fuzzy subspaces of vector spaces X , Y over F respectively . $K : A \rightarrow B$ is called a **fuzzy Zero transformation** on a vector space if $K(x_t) = 0_t$, for all $x_t \in A$ and $t \in [0,1]$.

2. Let A , B be fuzzy subspaces of vector spaces X , Y over F respectively . $K : A \rightarrow B$ is called a **fuzzy Identity transformation** on a vector space if $K(x_t) = x_t$, for all $x_t \in A$ and $t \in [0,1]$.

THEOREM 2.5 :

Let A , B be fuzzy subspaces of vector spaces X , Y over F respectively . $K : A \rightarrow B$ is a fuzzy linear transformation . Then , for all $t \in [0,1]$,

1. $K(0_t) = 0_t$;
2. $K(-x_t) = -K(x_t)$, for all $x_t \in A$;
3. $K(x_t - y_h) = K(x_t) - K(y_h)$, for all $x_t, y_h \in A$ and $t, h \in [0,1]$;
4. $K(\sum_{i=1}^n \lambda_i x_{ti}) \geq \sum_{i=1}^n \lambda_i K(x_{ti})$, for all $x_{ti} \in A$, $\lambda_i \in F$ and $t_i \in [0,1]$, $i = 1, 2, \dots, n$.

PROOF:

1. Since $0_t \circ 0_t = 0_t$, then $K(0_t) = K(0_t \circ 0_t) = K(0_t) \circ K(0_t) = 0_t$.
2. $K(-x_t) = K[(-1)(x_t)] = -1 K(x_t) = -K(x_t)$, for all $x_t \in A$.
3. $K(x_t - y_h) = K[(x_t) - (y_h)] = K(x_t) + K(-y_h) = K(x_t) - K(y_h)$, for all $x_t, y_h \in A$.
4. Since $K(\lambda_1 x_{t1}) = \lambda_1 K(x_{t1}) = \lambda_1 x_{t1}$, let $K(\sum_{i=1}^k \lambda_i x_{ti}) \geq \sum_{i=1}^k \lambda_i K(x_{ti})$, for all $x_{ti} \in A$, $\lambda_i \in F$ and $t_i \in [0,1]$, $i = 1, 2, \dots, k$.

To prove $K(\sum_{i=1}^{k+1} \lambda_i x_{ti}) \geq \sum_{i=1}^{k+1} \lambda_i K(x_{ti})$, for all $x_{ti} \in A$, $\lambda_i \in F$ and $t_i \in [0,1]$, $i = 1, 2, \dots, k+1$.

$$\begin{aligned}
 K(\sum_{i=1}^{k+1} \lambda_i x_{ti}) &= K(\sum_{i=1}^k \lambda_i x_{ti} + \lambda_{k+1} x_{t_{k+1}}) \\
 &\geq K(\sum_{i=1}^k \lambda_i x_{ti}) + K(\lambda_{k+1} x_{t_{k+1}}) \\
 &\geq \sum_{i=1}^k \lambda_i K(x_{ti}) + \lambda_{k+1} K(x_{t_{k+1}}) \\
 &\geq \sum_{i=1}^{k+1} \lambda_i K(x_{ti})
 \end{aligned}$$

Hence $K(\sum_{i=1}^n \lambda_i x_{ti}) \geq \sum_{i=1}^n \lambda_i K(x_{ti})$, for all $x_{ti} \in A$, $\lambda_i \in F$ and $t_i \in [0,1]$, $i = 1, 2, \dots, n$.

PROPOSITION 2.6 :

Let A be a fuzzy subspace of a vector space X over F and B be a fuzzy subspace of a vector space Y over F . Let $K : A \rightarrow B$, be an epimorphism fuzzy linear transformation , then :

1. $K(A)$ is a fuzzy subspace of B .
2. $K^{-1}(B)$ is a fuzzy subspace of A .

PROOF:

1. Let $x_{t1}, y_{t2} \in B$ such that $K(a_{t3}) = x_{t1}, K(b_{t4}) = y_{t2}$, where $a_{t3}, b_{t4} \in A$, since $t_1, t_2, t_3, t_4 \in [0,1]$ and $a_{t3} = K^{-1}(x_{t1}), b_{t4} = K^{-1}(y_{t2})$, where $a_{t3}, b_{t4} \in A$

$$K(A)(\lambda x_{t1} + \alpha y_{t2}) = \sup \{ \min \{ K(A)(a_{t3}), K(A)(b_{t4}) \} \mid a_{t3} = K^{-1}(\lambda x_{t1}), b_{t4} = K^{-1}(\alpha y_{t2}); \lambda x_{t1} + \alpha y_{t2} = a_{t3} + b_{t4} \},$$

$$\begin{aligned} &\geq \sup \{ \min \{ K(A)(a_{t3}), K(A)(b_{t4}) \} \mid a_{t3} = K^{-1}(x_{t1}), b_{t4} = K^{-1}(y_{t2}) \}, \\ &= \sup \{ \min \{ K(A)(a_{t3}), K(A)(b_{t4}) \} \mid K(a_{t3}) = x_{t1}, K(b_{t4}) = y_{t2} \}, \\ &\geq \min \{ K(A)(x_{t1}), K(A)(y_{t2}) \}; K(a_{t3}) = x_{t1}, K(b_{t4}) = y_{t2} \quad [4]. \end{aligned}$$

$K(A)(\lambda x_{t1} + \alpha y_{t2}) \geq \min \{ K(A)(x_{t1}), K(A)(y_{t2}) \}$. Then $K(A)$ is a fuzzy subspace of B .

2. Let $a_{t3}, b_{t4} \in A$ such that $a_{t3} = K^{-1}(x_{t1}), b_{t4} = K^{-1}(y_{t2})$, where $x_{t1}, y_{t2} \in B$, since $t_1, t_2, t_3, t_4 \in [0,1]$ and $K(a_{t3}) = x_{t1}, K(b_{t4}) = y_{t2}$, where $a_{t3}, b_{t4} \in A$

$$K^{-1}(B)(\lambda a_{t3} + \alpha b_{t4}) = \sup \{ \min \{ B(\lambda x_{t1}), B(\alpha y_{t2}) \} \mid x_{t1} = K(a_{t3}), y_{t2} = K(b_{t4}) \},$$

$$\begin{aligned} &\geq \sup \{ \min \{ B(x_{t1}), B(y_{t2}) \} \mid x_{t1} = K(a_{t3}), y_{t2} = K(b_{t4}) \} \\ &\geq \sup \{ \min \{ K^{-1}(B)(a_{t3}), K^{-1}(B)(b_{t4}) \} \mid a_{t3} = K^{-1}(x_{t1}), b_{t4} = K^{-1}(y_{t2}) \}, \\ &\geq \min \{ K^{-1}(B)(a_{t3}), K^{-1}(B)(b_{t4}) \}; a_{t3} = K^{-1}(x_{t1}), b_{t4} = K^{-1}(y_{t2}), [4]. \end{aligned}$$

$K^{-1}(B)(\lambda a_{t3} + \alpha b_{t4}) \geq \min \{ K^{-1}(B)(a_{t3}), K^{-1}(B)(b_{t4}) \}$. Then $K^{-1}(B)$ is a fuzzy subspace of A .

PROPOSITION 2.7 :

Let A be a fuzzy subspace of a vector space X over F and B be a fuzzy subspace of a vector space Y over F . Let $K : A \rightarrow B$ be a fuzzy linear transformation if and only if $f : X \rightarrow Y$ is a linear transformation on vector space .

PROOF:

(\Rightarrow) Since $K : A \rightarrow B$ is fuzzy linear transformation , that mean :

$$K(\lambda x_{t1} + \alpha y_{t2}) \geq \min \{ K(x_{t1}), K(y_{t2}) \}, \text{ for all } x_{t1}, y_{t2} \in A \text{ and } \lambda, \alpha \in F, t_1, t_2 \in [0,1].$$

To prove $f : X \rightarrow Y$ is a linear transformation on vector space ,(i.e.) $f(\lambda x + \alpha y) = \lambda f(x) + \alpha f(y)$, for all $x, y \in X$ and $\lambda, \alpha \in F$.

Since $x_{t1}, y_{t2} \in A, t_1, t_2 \in [0,1]$, then there exists $x, y \in X$ such that $K(x_{t1}) = f(x), K(y_{t2}) = f(y)$ implies that $x = f^{-1}(K(x_{t1}))$ and $y = f^{-1}(K(y_{t2}))$.

$$\begin{aligned} f(\lambda x + \alpha y) &= f(\lambda x) + f(\alpha y) \\ &= f(\lambda f^{-1}(K(x_{t1}))) + f(\alpha f^{-1}(K(y_{t2}))) \\ &= \lambda f(f^{-1}(K(x_{t1}))) + \alpha f(f^{-1}(K(y_{t2}))) \\ &= \lambda K(x_{t1}) + \alpha K(y_{t2}) \\ &= \lambda f(x) + \alpha f(y), \text{ for all } x, y \in X. \end{aligned}$$

Then $f(\lambda x + \alpha y) = \lambda f(x) + \alpha f(y)$, for all $x, y \in X$ and $\lambda, \alpha \in F$.

Hence $f : X \rightarrow Y$ is a linear transformation on vector space .

(\Leftarrow) Since $f : X \rightarrow Y$ is a linear transformation on vector space , that mean :

$$f(\lambda x + \alpha y) = \lambda f(x) + \alpha f(y), \text{ for all } x, y \in X.$$

To prove $K : A \rightarrow B$ is fuzzy linear transformation , (i.e.)

$$K(\lambda x_{t1} + \alpha y_{t2}) \geq \min \{ K(x_{t1}), K(y_{t2}) \}, \text{ for all } x_{t1}, y_{t2} \in A \text{ and } \lambda, \alpha \in F, t_1, t_2 \in [0,1].$$

Since $\lambda, \alpha \in F$ and $x, y \in X$, then there exists $x_{t1}, y_{t2} \in A, t_1, t_2 \in [0,1]$ such that $K(x_{t1}) = f(x), K(y_{t2}) = f(y)$ implies that $x_{t1} = K^{-1}(f(x))$ and $y_{t2} = K^{-1}(f(y))$.

$$\begin{aligned} K(\lambda x_{t1} + \alpha y_{t2}) &= K(\lambda x_{t1}) + K(\alpha y_{t2}) \\ &= K(\lambda K^{-1}(f(x))) + K(\alpha K^{-1}(f(y))) \\ &= \lambda K(K^{-1}(f(x))) + \alpha K(K^{-1}(f(y))) \\ &= \lambda f(x) + \alpha f(y) \end{aligned}$$

$$= \lambda K(x_{t1}) + \alpha K(y_{t2}), \text{ for all } x, y \in X, \lambda, \alpha \in F.$$

$$\geq K(x_{t1}) + K(y_{t2}), \text{ for all } x, y \in X, \lambda, \alpha \in F.$$

$$K(\lambda x_{t1} + \alpha y_{t2}) \geq K(x_{t1}) \text{ and } K(\lambda x_{t1} + \alpha y_{t2}) \geq K(y_{t2}), [4].$$

$$K(\lambda x_{t1} + \alpha y_{t2}) \geq \min \{ K(x_{t1}), K(y_{t2}) \}, \text{ for all } x_{t1}, y_{t2} \in A \text{ and } \lambda, \alpha \in F, t_1, t_2 \in [0,1].$$

Hence $K : A \rightarrow B$ is a fuzzy linear transformation .

PROPOSITION 2.8 :

Let A be a fuzzy subspace of a finite vector space $X = \{ x_{t1}, x_{t2}, \dots, x_{tn} \}$ over F and B be a fuzzy subspace of a finite vector space $Y = \{ y_{t1}, y_{t2}, \dots, y_{tn} \}$ over F . Then $K : A \rightarrow B$ such that $K(x_{ti}) = y_{ti}$, for all $i = 1, 2, \dots, n, t \in [0,1]$ is a fuzzy linear transformation .

PROOF:

Since a finite vector space $X = \{ x_{t1}, x_{t2}, \dots, x_{tn} \}$ over F and a finite vector space $Y = \{ y_{t1}, y_{t2}, \dots, y_{tn} \}$ over F , then $f : X \rightarrow Y$ such that $f(x_i) = y_i$, for all $i = 1, 2, \dots, n$, by [3].

Then A_t is a subspace of X , for all $0 \leq t \leq A(0)$, and proposition (1.8 (2)), A is a fuzzy subspace of X .

Hence $K \approx f$ by proposition (2.7), and $K(x_{ti}) = y_{ti}$, for all $i = 1, 2, \dots, n, t \in [0,1]$.

To prove K is a fuzzy linear transformation .Let $x_{t1}, x_{t2} \in A$ such that :

$$x_{t1} = \sum_{i=1}^n \lambda_i y_{ii}, x_{t2} = \sum_{i=1}^n \alpha_i y_{ii}, \text{ then } (\lambda x_{t1} + \alpha x_{t2}) = (\sum_{i=1}^n (\beta \lambda_i + \mu \alpha_i) x_{ii}), \beta, \mu \in F.$$

$$\begin{aligned} K(\lambda x_{t1} + \alpha x_{t2}) &= K(\sum_{i=1}^n (\beta \lambda_i + \mu \alpha_i) y_{ii}) \\ &= \beta K(\sum_{i=1}^n \lambda_i y_{ii}) + \mu K(\sum_{i=1}^n \alpha_i y_{ii}) \\ &= \beta K(x_{t1}) + \mu K(x_{t2}) \end{aligned}$$

$$K(\lambda x_{t1} + \alpha x_{t2}) \geq \beta K(x_{t1}) \text{ and } K(\lambda x_{t1} + \alpha x_{t2}) \geq \mu K(x_{t2}).$$

$$K(\lambda x_{t1} + \alpha x_{t2}) \geq K(x_{t1}) \text{ and } K(\lambda x_{t1} + \alpha x_{t2}) \geq K(x_{t2}).$$

$$K(\lambda x_{t1} + \alpha x_{t2}) \geq \min \{ K(x_{t1}), K(x_{t2}) \}.$$

Hence K is a fuzzy linear transformation .

DEFINITION 2.9 [7] :

A linear transformation f from a ring $(R, +, \cdot)$ to a ring $(R', +', \cdot')$ is called **ring homomorphism** if it satisfies the following properties : for all $a, b \in R$,

1. $f(a + b) = f(a) +' f(b)$

2. $f(a \cdot b) = f(a) \cdot' f(b)$.

PROPOSITION 2.10 [7] :

If $f : R \rightarrow R'$ and $g : R' \rightarrow R''$ are homomorphism between the fuzzy subsets A, B and C , then $f \circ g$ is a homomorphism between A and C .

REMARK 2.11 [7] :

1. If f and g are isomorphism, then $g \circ f$ is an isomorphism since f and g are one – to- one and onto implies that $g \circ f$ is one – to - one and onto.

2. If f and g are homomorphism , one – to - one and onto, then $g \circ f$ is an isomorphism .

THEROEM 2.12 :

Let A, B, C be fuzzy subspaces of vector space X, Y, Z over F respectively and let $K : A \rightarrow B, G : B \rightarrow C$ be fuzzy linear transformations . Then $G \circ K : A \rightarrow C$ be a fuzzy linear transformation .

PROOF:

Since K and G are fuzzy linear transformations , then :

$$K(\lambda x_{t1} + \alpha y_{t2}) = \sup \{ \inf \{ \lambda, K(x_{t1}), \alpha, K(y_{t2}) \}, \text{ for all } x_{t1}, y_{t2} \in A \text{ and } \lambda, \alpha \in F, t_1, t_2 \in [0,1].$$

$$G(\lambda z_{t3} + \alpha u_{t4}) = \sup \{ \inf \{ \lambda, G(z_{t3}), \alpha, G(u_{t4}) \}, \text{ for all } z_{t3}, u_{t4} \in B \text{ and } \lambda, \alpha \in F, t_3, t_4 \in [0,1]$$

$$\text{and } K(x_{t1}) = z_{t3}, K(y_{t2}) = u_{t4}, K(z_{t3}) = a_{t5}, K(u_{t4}) = b_{t6}, a_{t5}, b_{t6} \in A, t_5, t_6 \in [0,1].$$

To prove $G \circ K : A \rightarrow C$ be a fuzzy linear transformation .

Let $a_{t5}, b_{t6} \in A$ (that mean $a, b \in X$ and $t_5, t_6 \in [0,1]$), for all $\lambda, \alpha \in F$, then :

$$\begin{aligned} G \circ K (\lambda a_{t5} + \alpha b_{t6}) &= G(K (\lambda a_{t5} + \alpha b_{t6})) \\ &\geq G(\min \{ K(a_{t5}), K(b_{t6}) \}), \\ &= G(\min \{ z_{t3}, u_{t4} \}), \\ &= \min \{ G(z_{t3}), G(u_{t4}) \}, [4] . \\ &= \min \{ G(K(x_{t1})), G(K(b_{t6})) \}, \\ &= \min \{ G \circ K (a_{t5}), G \circ K (b_{t6}) \}. \end{aligned}$$

$$G \circ K (\lambda a_{t5} + \alpha b_{t6}) \geq \min \{ G \circ K (a_{t5}), G \circ K (b_{t6}) \}.$$

Then $G \circ K : A \rightarrow C$ be a fuzzy linear transformation .

DEFINITION 2.13 ([8], [4]):

Let $X : R \rightarrow [0,1], Y : R' \rightarrow [0,1]$ are fuzzy sets . $f : R \rightarrow R'$ be homomorphism between them.

We define **the fuzzy kernel of f**, $\ker f_{zz} : R \rightarrow [0,1]$ by :

$$\ker f_{zz} f(x) = \begin{cases} X(0) & x \in \ker f \\ 0 & x \notin \ker f \end{cases}$$

DEFINITION 2.14 :

Let A be a fuzzy subspace of a vector space X over R and B be a fuzzy subspace of a vector space Y over R' . Let $K : A \rightarrow B$ be a fuzzy linear transformation and $f : X \rightarrow Y$ be homomorphism between them . We define **the fuzzy kernel of K**, $\ker f_{zz} K : A \rightarrow [0,1]$ by :

$$\ker f_{zz} K(x) = \begin{cases} A(0) & x \in \ker f \\ 0 & x \notin \ker f \end{cases} .$$

PROPOSITION 2.15 :

$\ker f_{zz} K : A \rightarrow [0,1]$ is a fuzzy subspace of X .

PROOF:

Let $a, b \in X$, for all $\lambda, \alpha \in F$, since $\ker f_{zz} K(0) = X(0)$, if $x \in \ker f$, then :

$$\ker f_{zz} K(\lambda a + \alpha b) = \begin{cases} A(0) & \lambda a + \alpha b \in \ker f \\ 0 & \lambda a + \alpha b \notin \ker f \end{cases} .$$

$$\ker f_{zz} K(a) = \begin{cases} A(0) & a \in \ker f \\ 0 & a \notin \ker f \end{cases} .$$

$$\ker f_{zz} K(b) = \begin{cases} A(0) & b \in \ker f \\ 0 & b \notin \ker f \end{cases} .$$

Then $\ker f_{zz} K (\lambda a + \alpha b) = \sup \{ \inf \{ \lambda, \ker f_{zz} K (a), \alpha, \ker f_{zz} K (b) \} \}$.

Hence $\ker f_{zz} K$ is a fuzzy subspace of X .

PROPOSITION 2.16 :

Let A be a fuzzy subspace of a vector space X over F and B be a fuzzy subspace of a vector space Y over F and $K : A \rightarrow B$ be a fuzzy linear transformation . Then $\ker f_{zz} K = \phi$ if and only if K is one - to - one.

PROOF:

Since $\ker f_{zz} K = \phi$, then $\ker f = \{0\}$, by theorem (2.1.10) in [3], K is a one - to - one.

SECTION THREE

Fuzzy Coset and Quotient Fuzzy Rings

In this section, two definitions about fuzzy coset and quotient fuzzy ring are given , some properties concerning with this definitions are given and we studied the concept of a fuzzy isomorphism.

DEFINITION 3.1 [3]:

Let A and B be fuzzy subsets of vector space X over F such that $B \subseteq A$ and $x_t \subseteq A$, $t \in [0, A(0)]$. Then $x_t + B$ ($B + x_t$) is called a **fuzzy left (right) coset of B in A with representative x_t** .

REMARK 3.2 [3]:

Let A and B be fuzzy subsets of vector space X over F such that $B \subseteq A$ and $x_t \subseteq A$, $t \in [0, A(0)]$. For all $z \in X$, $(x_t + B)(z) = \inf \{ t , B(z-x) \}$ and $(A/B) = \{ x_t + B : x_t \subseteq A, x \in B \}$ is commutative group under + .

PROPOSITION 3.3 [3]:

Let A and B be fuzzy subsets of vector space X over F such that $B \subseteq A$ and $x_t , y_s \subseteq A$, $t, s \in [0, A(0)]$. Then :

1. For all $z \in G$, $(x_t + B)(z) = \inf \{ t , B(z-x) \}$ and $(B + x_t)(z) = \inf \{ t , B(x + (-z)) \}$.
2. (a) $x_t + B = y_s + B$ iff $\inf \{ t , B(e) \} = \inf \{ s , B((-y)+x) \}$ and $\inf \{ s , B(e) \} = \inf \{ t , B(x + (-y)) \}$.
 (b) $x_t + B = y_s + B$ iff $\inf \{ t , B(e) \} = \inf \{ s , B(x + (-y)) \}$ and $\inf \{ s , B(e) \} = \inf \{ t , B(y + (-x)) \}$.
3. If $B((-y) + x) = B(e)$, then $x_t + B = y_t + B$.

DEFINITION 3.4 [18]:

Let A and B be fuzzy subsets of vector space X over F such that $B \subseteq A$ and $x_t \subseteq A$, $t \in [0, A(0)]$. $B(e) = A(e)$ and B is a fuzzy normal in A . Then $(A/B)_t = \{ x_t + B : x_t \subseteq A, x \in G \}$, for all $t \in [0, 1]$ is a group under “+” . $(A/B)_t$ is called a **quotient group of fuzzy subgroup** .

$(A/B) = \{ x_t + B : x_t \subseteq A, x \in G, t \in [0, 1] \}$. Then $((A/B), +)$ is a semigroup with identity and (A/B) is completely regular ((A/B) is a union of disjoint groups) i.e. , $(A/B) = \bigcup_{t \in [0, A(0)]} (A/B)_{(t)}$.

PROPOSITION 3.5 [3]:

Let A and B be fuzzy subsets of vector space X over F such that $B \subseteq A$ and $x_t \subseteq A$, $t \in [0, A(0)]$. Then $(A/B)_t = A_t / B_t$.

PROPOSITION 3.6 :

Let A and B be two fuzzy subspaces of a vector space X over F such that $B \subseteq A$ and $x_t , y_t \subseteq A$, $t \in [0, A(0)]$. Then (A/B) is a fuzzy subspace over F on (+ and .) such that :

1. $(x_t + B) + (y_t + B) = (x_t + y_t) + B$.
2. $\lambda(x_t + B) = (\lambda x_t) + B$, for all $\lambda \in F$.

PROOF:

Let $x_t , y_t \subseteq A$, $t \in [0, A(0)]$ and $B \subseteq A$, then $(x_t + y_t) \subseteq A$ and $\lambda x_t \subseteq A$. Thus $(x_t + y_t) + A \subseteq (A/B)_t$, then $(A/B, +)$ and $(A/B, .)$ are closure on (+ and .) .

Let $z_t , u_t \subseteq A$, $t \in [0, A(0)]$ and $B \subseteq A$, then $(x_t - z_t) \subseteq A$ and $(y_t - u_t) \subseteq A$, since A is a vector subspace , $(x_t - z_t) + (y_t - u_t) \subseteq A$ implies that $(x_t + y_t) - (z_t + u_t) \subseteq A$ implies that $(x_t + y_t) + A = (z_t + u_t) + A \subseteq A$ implies that $(A/B)_t$ is a well defined of (+) .

And $(x_t - z_t) \subseteq A$ and $(\lambda(x_t - z_t)) \subseteq A$ implies that $(\lambda x_t - \lambda z_t) \subseteq A$ implies that $(\lambda x_t) + A = (\lambda y_t) + A \subseteq A$ implies that $(A/B)_t$ is a well defined of (.) .

Since $A(0) > 0$, $A(x - y) \geq \min\{ A(x) , A(y) \}$, for all $x , y \in X$ and $A(cx) \geq \min \{ F(c) , A(x) \}$, for all $x \in X$ and $c \in F$, then (A/B) is a fuzzy subspace over F on $(+ \text{ and } .)$

THEOREM 3.7 :

Let A and B be fuzzy subspaces of vector space X over F such that $B \subseteq A$ and $x_t , y_t \subseteq A$, $t \in [0, A(0)]$. Then $K : X \rightarrow X / A$ define by: $f(x) = x_t + A$. Then K is an epimorphism fuzzy linear transformation and $\ker f_{zz} K = A$.

PROOF:

Let $x_t , y_t \subseteq X$, $t \in [0, A(0)]$ and $\alpha , \beta \in F$, then :

$$\begin{aligned} 1. \quad K(\alpha x_t + \beta y_t) &= (\alpha x_t + \beta y_t) + A \\ &= \alpha(x_t + A) + \beta(y_t + A) \\ &\geq \min \{ (x_t + A) , (y_t + A) \} \\ &= \min \{ K(x_t) , K(y_t) \} . \end{aligned}$$

Then K is a fuzzy linear transformation .

2. Let $z_t \subseteq X / A$, then there exists $x \in X$ such that $z_t = (x_t + A) = K(x_t)$, then K is a onto .
3. Since the fuzzy kernel of K is $\ker f_{zz} K : A \rightarrow [0,1]$ by :

$$\ker f_{zz} K(x) = \begin{cases} A(0) & x \in \ker f \\ 0 & x \notin \ker f \end{cases} . \text{ Then } \ker f_{zz} K = A .$$

REMARK 3.8 :

The function K is called **fuzzy Canonical function** . In general , K is one – to- one since $x_t , y_t \subseteq A$, $t \in [0, A(0)]$. Then $(x_t - y_t) \subseteq A$ implies that $(x_t + A) = (y_t + A)$, then $K(x) = K(y)$.

DEFINITION 3.9 :

Let X and Y be fuzzy subsets over F . Then we define **fuzzy linear isomorphism** , if there exists $K : X \rightarrow Y$ is a fuzzy linear transformation , one – to – one and onto . We denoted by $X \approx Y$.

THEOREM 3.10 :

Let A , B , C be fuzzy subspaces of vector spaces X , Y and Z over F respectively such that $A = B \oplus C$. Then $B \approx A / C$ or $C \approx A / B$.

PROOF:

Define $K : B \rightarrow A / C$ such that $K(x) = x_t + C$, $x_t \subseteq B$.

Let $x_t , y_t \subseteq B$, $t \in [0, 1]$ and $\alpha , \beta \in F$, then :

$$\begin{aligned} 1. \quad K(\alpha x_t + \beta y_t) &= (\alpha x_t + \beta y_t) + C \\ &= \alpha(x_t + C) + \beta(y_t + C) \\ &\geq \min \{ (x_t + C) , (y_t + C) \} \\ &= \min \{ K(x_t) , K(y_t) \} . \end{aligned}$$

Then K is a fuzzy linear transformation .

2. Let $z_t \subseteq A / C$, then there exists $x_t \in B$ such that $z_t = (x_t + C) = K(x_t)$, but $A = B \oplus C$, then $x_t = (u_t + w_t)$, $u_t \subseteq B$ and $w_t \subseteq C$) implies that $u_t = (x_t - w_t)$, thus $x_t + C = w_t + C$, then $z_t = K(u_t)$. Hence K is a onto .
3. Since $x_t , y_t \subseteq B$ such that $K(x_t) = K(y_t)$, then $x_t + C = y_t + C$, $t \in [0, 1]$ and $(x_t - w_t) \subseteq C$, K is a one – to – one .

Then $B \approx A / C$, by this style $C \approx A / B$.

(First Fuzzy Isomorphism Theorem For Fuzzy Subspaces)

THEOREM 3.11 :

Let X and Y are fuzzy subspaces of a vector subspace over F and K be onto homomorphism between them. Then $X / \ker f_{zz} K \approx K(X)$.

PROOF:

Define $G : X / \ker K \rightarrow K(X)$ such that: $G(a_t + \ker K) = K(a_t)$, for each $a_t + \ker K \in X / \ker K$.

By definition, G is a non empty function of $X / \ker K$ since $g(0_t + \ker K) = K(0_t)$

Let $a_t + \ker K, b_t + \ker K \in X / \ker K$, $a_t + \ker K = b_t + \ker K$ implies that $a_t - b_t \in \ker K$, therefore $K(a_t - b_t) = 0_t$ and K is homomorphism, then $K(a_t) - K(b_t) = 0_t$ implies $K(a_t) = K(b_t)$. Thus $G(a_t + \ker K) = G(b_t + \ker K)$. Hence G is well - define.

Now, we must prove G is an isomorphism

First, if $G(a_t + \ker K) = G(b_t + \ker K)$, then $K(a_t) = K(b_t)$ and $K(a_t) - K(b_t) = 0_t$ implies that $K(a_t - b_t) = 0_t$. Thus $a_t - b_t \in \ker K$ therefore $a_t + \ker K = b_t + \ker K$, G is one - to - one.

Second, for any $b_t \in K(X)$ there exists $a_t \subseteq X$ such that: $K(a_t) = b_t$ since K is onto then $K(a_t) = G(a_t + \ker K) = b_t$, G is onto .

Finally, Let $a_t + \ker K, b_t + \ker K \in X / \ker K$ and $\alpha, \beta \in F$, then :

$$\begin{aligned} G[\alpha(a_t + \ker K) \oplus \beta(b_t + \ker K)] &= G[(\alpha a_t + \beta b_t) + \ker K] \\ &= K(\alpha a_t + \beta b_t) \\ &= \alpha K(a_t) + \beta K(b_t) \\ &= G(\alpha(a_t + \ker K)) + G(\beta(b_t + \ker K)) \\ &\geq \min\{G((a_t + \ker K)), G((b_t + \ker K))\} \end{aligned}$$

Then G is a fuzzy linear transformation .

Hence $X / \ker f_{zz}K \approx K(X)$.

(Second Fuzzy Isomorphism Theorem For Fuzzy Subspaces)

THEOREM 3.12 :

Let A and B be fuzzy subspaces of a fuzzy subspace X over F , with $A \subseteq B$ such that. $B(x) = B(0)$, whenever $A(x) = A(0)$. Then $(X / A) / (B / A) \approx (X / B)$.

PROOF:

Define $G : (X / A) / (B / A) \rightarrow (X / B)$ such that : $G((x_t + A) + (B / A)) = x_t + B$ is an isomorphism by [7] .

By definition, G is a non empty function of $(X / A) / (B / A)$ since $g(0_t + (B / A)) = K(0_t + A)$

Let $(a_t + A + (B / A)), (b_t + A + (B / A)) \in (X / A) / (B / A)$, $(a_t + A + (B / A)) = (b_t + A + (B / A))$ implies that $((a_t - b_t) + A) \in (B / A)$, therefore $K((a_t - b_t) + A) = 0_t + A$ and K is homomorphism, then $K(a_t + A) - K(b_t + A) = 0_t + A$ implies $K(a_t + A) = K(b_t + A)$. Thus $G(a_t + A + (B / A)) = G(b_t + A + (B / A))$. Hence G is well - define.

Now, we must prove G is an isomorphism

First, if $G(a_t + A + (B / A)) = G(b_t + A + (B / A))$, then $K(a_t + A) = K(b_t + A)$ and $K(a_t + A) - K(b_t + A) = 0_t + A$ implies that $K((a_t - b_t) + A) = 0_t + A$. Thus $(a_t - b_t) + A \in (B / A)$ therefore $(a_t + A) + (B / A) = (b_t + A + (B / A))$, G is one - to - one.

Second, for any $(b_t + A) \in K((X / A))$ there exists $a_t + A \subseteq (X / A)$ such that: $K(a_t + A) = b_t + a$, since K is onto then $K(a_t + A) = G(a_t + A + (B / A)) = (b_t + A)$, G is onto .

Finally, Let $(a_t + A + (B / A)), (b_t + A + (B / A)) \in (X / A) / (B / A)$ and $\alpha, \beta \in F$, then :

$$\begin{aligned} G[\alpha(a_t + A + (B / A)) \oplus \beta(b_t + A + (B / A))] &= G[(\alpha(a_t + A) + \beta(b_t + A)) + (B / A)] \\ &= K(\alpha(a_t + A) + \beta(b_t + A)) \\ &= \alpha K(a_t + A) + \beta K(b_t + A) \\ &= G(\alpha(a_t + A + (B / A))) + G(\beta(b_t + A + (B / A))) \\ &\geq \min\{G(a_t + A + (B / A)), G(b_t + A + (B / A))\}. \end{aligned}$$

Then G is a fuzzy linear transformation .

Hence $(X / A) / (B / A) \approx K(X / A)$.

REFERENCES

- 1) AL- Khamees Y. and Mordeson J.N., 1998, Fuzzy Principal Ideals and Simple Field Extensions , Fuzzy Sets and Systems, vol.96, pp.247 – 253.
- 2) AL-Khfaji S.M. , 2010 , On Fuzzy Topological Vector Spaces , M.Sc.Thesis, University of Al-Qadisiyah , College of Computer Sciences and Mathematics .
- 3) AL-Mayahi N.F. and Battor A. H. , 2005 , Introduction to Functional Analysis , AL-Nebras Company .
- 4) Bhambert S.K. , Kumar R. and Kumar P. , 1995 , Fuzzy Prime Submodules and Radical of a Fuzzy Submodules , Bull. Col. Math. Soc. ,vol.87 , No.4 , pp.163-168 .
- 5) Dixit V.N.,Kumar R. and Ajmal N., 1991, Fuzzy Ideals and Fuzzy Prime Ideals of a Ring, Fuzzy Sets and Systems, vol.44, pp. 127 – 138.
- 6) Golan J.S. , 1989 , Making Modules Fuzzy , Fuzzy Sets and Systems, vol.32, pp. 91 – 94.
- 7) Kasch F. , Wallace D.A.R. , 1982 , Modules and Rings , A Subsidiary of Harcourt Brace Jovanich Publishers , London .
- 8) Katsaras A.K. and Liu D.B. ,1977 , Fuzzy Vector Spaces and Fuzzy Topological Vector Spaces, J. Math. Anal. , vol.58 , pp.135-146 .
- 9) Liu W.J., 1982, Fuzzy Invariant Subgroups and Fuzzy Ideal , Fuzzy Sets and Systems, vol.8, pp.133 – 139.
- 10) Martines L., 1995, Fuzzy Subgroup of Fuzzy Groups and Fuzzy Ideals of Fuzzy Rings , the journal of fuzzy mathematics, vol.3, No.4, pp.833 – 849.
- 11) Martinez L., 1996 , Fuzzy Modules Over Fuzzy Rings In Connection With Fuzzy Ideals Of Fuzzy Rings, the journal of fuzzy mathematics, vol.4, No.4, pp.843 – 857.
- 12) Mordeson J.N. , 1993 , Bases of Fuzzy Vector Spaces , Information Sciences , vol.67 , pp.87-92 .
- 13) Mordeson J.N. and Sen M.K. , 1995 , Basic Fuzzy Subgroups , Information Sciences , vol.82 , pp.167-179 .
- 14) Mukherjee T.K. and Sen M.K., 1987, On Fuzzy Ideals of a Ring I, Fuzzy Sets and Systems, vol.21, pp. 99 – 104.
- 15) Pan F. , 1993 , Finitely Fuzzy Value Distribution of Fuzzy Vector Spaces and Fuzzy Modules , Fuzzy Sets and Systems , vol . 55 , pp. 319 – 322 .
- 16) Rudin W. , 1973, Functional Analysis , Mc Graw-Hill B. Company .
- 17) Zadeh L.A., 1965, Fuzzy Sets, Information and Control, vol.8, pp.338 – 353.