

## **Clinical Evaluation Of Bovine - Derived Xenograft In Treatment Of Human Periodontal Class II Molar Furcation Defects.**

Dr.Zainab Muhi Hameed Al- Fatlawi  
B.D.S,M.Sc.(Periodontology).  
Babylon University

### **Abstract :-**

The aim of this study is clinical evaluation of bovine derived xenograft in the treatment of human periodontal Class II molar furcation defects. Eighteen patients(8 males, 10 females) suffering from chronic periodontitis,with Grade II molar furcation defect, were treated using bovine derived xenograft , completed the study with the following clinical parameters were assessed immediately before the surgical procedure(Baseline) and at 6 months after treatment.These parameters were, Plaque index Score(PII),gingival index (GI) periodontal pocket depth(PPD),Clinical attachment level (CAL),Gingival Recession(GR) and Closed horizontal probing depth (CHPD).This therapy resulted in significant PPD reductions and no significant improvement was seen (CAL),(GR) and(CHPD). For the mean values for the of (PPD)(5.47±1.39),(CAL)(3.1±1.4), (GR)(0.21±0.51) and(CHPD)(3.5±1.3) at baseline ,while these parameters at 6 months after surgical treatment were(3.8±1.52),(2.35± 1.34), (0.3±0.5)and(2.2±1.3) respectively.

The findings of this study suggest improvement in clinical results with bovine derived xenograft treatment in class II furcation molar defects.

**Key words:** periodontal; regeneration; furcations; graft, bone,bovine- derived xenograft .

### **الخلاصة**

الهدف من هذه الدراسة هو لتقييم القياسات السريرية الناتجة عن استخدام الطعم العظمي المشتق من الابقار في علاج مجموعة من الناس لديهم اضرار فيها منطقة تشعب او تفرع جذور مصابة سريريا (درجة ثانية) و هذه القياسات تمثلت بعمق الجيب اللثوي وفقدان التعلق السريري و انحسار اللثة و عمق الجيب العرضي و مقارنة هذه النتائج مع نفس القياسات ولكن بعد مرور ستة اشهر من العلاج باستخدام التجريف الجراحي.العينة تكونت من 18 شخصا مصابين بالتهاب لثوي مزمن يرافقه اضرار فيها منطقة تشعب او تفرع الجذور مصابة سريريا(درجة ثانية) . التحليل الاحصائي للقراءة الاولى و القراءة الثانية بعد ستة اشهر اوضحت النتائج ان نقصان عمق الجيب اللثوي وفقدان التعلق السريري و انحسار اللثة و عمق الجيب العرضي مما يدل على تحسن سريري واضح في علاج هذه الاضرار التي فيها منطقة تشعب او تفرع جذور المصابة سريريا باستخدام التجريف الجراحي و الطعم العظمي المشتق من الابقار .

### **Introduction:-**

Molar furcation involvement is one of the most common dento-alveolar sequelae of periodontal disease. The term furcation involvement :refers to the involvement of bifurcation or trifurcation of the multirrooted teeth by the periodontal diseases. The bacterial plaque is the causative agent of the gingivitis and destructive periodontal disease causes destruction of the periodontal tissues.Grade II molar furcation defects mean bone destruction on one or more aspects of the furaction ,but a portion of the alveolar bone and periodontal ligament remains intact.or :-horizontal loss of the periodontal tissue support not exceeding 1/3 of the width of the tooth(1). The application of a specific treatment method for furcation involvement requires a thorough understanding of tooth anatomy, etiologic factors, and the biologic basis for treatment modalities.(2)

Contributing factors to furcation involvement include systemic factors ,such as diabetes and smoking(3-6) and local factors, such as cervical enamel projections(7,8) furcation entrance width(9) ,furcation and root concavities(10),bifurcational ridges(8,11),accessory pulpal canals (12-14) ,enamel pearls(15) and furcation restorations(16).

These factors must be assessed thoroughly to ensure a correct diagnosis leading to effective management of furcation involvement(17).

Historically, various treatment methods have been proposed to treat molar furcation defects. These methods ranged from conservative therapy, such as curettage and open flap debridement, to surgical treatment procedures, such as gingivectomy, root amputation, hemisection, or tunneling. Regardless of the treatment method used, most longitudinal studies have shown that molars are at higher risk for tooth loss than non molar teeth.(18-20).

The introduction of bone grafting methods ( 21-23), and the concept of tissue regeneration(24-26) offered new hope for improved and more predictable treatment of furcation involvement.

There are different methods of regenerative treatments like GTR (Guided Tissue Regeneration) technique, using bone graft and synthetic materials, and combination methods. Auto graft, Allograft and Xenograft bone and Alloplastic materials have been used in periodontal defects. However, there is no material for complete regeneration until now. The main reasons of using this material are improvement of bone regeneration and accessibility for patients, to remove microbial plaque, prevention of disease progress and losing tooth.(27) Many researches have performed successful results using Xenograft bone in periodontal defects. Some of these researches are in histological evaluation of Xenograft in human periodontal lesion(28,29) .The combination therapy of a membrane and xenograft for molars has resulted in successful closure of furcations.(28,30-33). Class II furcations have been shown to be the best candidates for regenerative treatment(34-36)

In assessing the success of these treatment methods, complete closure of the defect is desirable.(2)

Therapeutic results can be measured by periodontal probing depth (PPD) and clinical attachment level (CAL) improvements, bone regeneration, and evidence of histologic periodontal regeneration. Although histologic evaluation is most accurate, surgical closure of the furcation defect and improvements in PPD and CAL serve as suitable and practical outcome measures.(37,38).

The aim of this study is clinical evaluation of bovine derived xenograft in the treatment of human periodontal Class II molar furcation defects

## **Material and Methods**

### **Population screening**

Potential patients (18 patients) were selected from those referred to the periodontal department clinic of the college of dentistry (Babylon university) &/or private clinic. All patients received a complete periodontal examination, including a full-mouth periodontal probing ( Fig 1), radiographic examination (Fig 2). The study inclusion criteria were (i) diagnosis of chronic periodontitis (according to the criteria of the 1999 international classification;)( 39); (ii) presence of class-II furcations, presenting PD $\geq$ 5mm and bleeding on probing, after nonsurgical therapy; (iii) good general health. Patients who (i) were pregnant or lactating; (ii) required antibiotic pre-medication for the performance of periodontal examination and treatment; (iii) suffered from any other systemic diseases (cardiovascular, pulmonary, liver, cerebral, diseases or diabetes); (iv) had received antibiotic treatment in the previous 3 months; (v) were taking long-term anti-inflammatory drugs; (vi) were smokers were excluded from the study).

### **Non-surgical treatment**

All the subjects received a full-mouth periodontal treatment 2 months before the surgical procedure. All these treatments were performed by the same operator, with an ultrasonic device. At the same time the subjects underwent motivation session, during which oral hygiene instructions were given, to ensure that the subjects could maintain a proper level of oral hygiene before the surgical procedure.

### **Clinical parameters**

The following clinical parameters were assessed immediately before the surgical Procedure (Baseline) and at 6 months after treatment. These parameters were, Plaque index Score (PII) (40), gingival index Score (GI) (41) periodontal pocket depth (PPD), Clinical attachment level (CAL), Gingival Recession (GR) and Closed horizontal probing depth (CHPD) were calculated when probing with manual probe. (Fig 1)

- 1- Probing pocket depth (PPD): the distance from the free gingival margin to the base of the pocket. In each tooth, probing was applied in 3 areas and the highest Deep in the furcation was registered.
- 2- Clinical attachment level (CAL): the distance from CEJ to the base of the pocket.
- 3-Gingival Recession(GR): gingival margin position to the CEJ in 3 points of each Surface of the tooth.
- 4-Closed horizontal probing depth (CHPD): the distance from deepest area of probe Penetration vertically to buccal or lingual surfaces to connection line of hight of contour of mesial and distal roots.

### **Surgical procedures**

Following local anesthesia, sulcular incisions were made and mucoperiosteal flaps were raised at furcation area. Carefully, the tissue was reflected, preserving the maximum of interproximal soft tissue. Granulation tissue as well as the visible calculus were removed with hand currettes and with an ultrasonic device. The bone graft was applied from the farthest end of the involved furcation until the buccal surface of the tooth was covered with it. The surgical flaps were then replaced at their initial position and sutured. A black silk suture(3-0) using vertical mattress or an interrupted technique. After suturing, slight pressure on the facial and lingual flaps is applied to minimize the clot beneath the flap. It is optional to place a surgical dressing to protect the wound. Placement of a dressing must be a accomplished, however, without displacement of the graft or compromise of the blood supply to the gingival flaps for a period of 7 to 10 days, at which time the dressing and sutures were removed. No further dressings were placed unless special conditions dictated otherwise.

### **Postoperative management/periodontal maintenance**

The administration of antibiotics beginning immediately post-surgery is thought to aid in plaque control (42) Amoxicillin 500 mg t.i.d was prescribed for the 10-day immediate postsurgical period. NSAIDs were begun one hour prior to surgery and continued t.i.d for 3 days postsurgically. Patients should also be placed on achlorhexidine mouthwash to further aid in this process(43). For the first 4–6 weeks, patients should refrain from brushing the surgical area to prevent disturbance of the blood clot (44). Sutures are retained as long as they maintain closure and do not contribute to plaque accumulation and inflammation .Postoperative visits include plaque removal (mechanically), selective stain removal and reinforcement of oral hygiene. Periodontal probing should not be done prior to 6 months, since probing force may damage the healing site, thereby diminishing the regenerative outcome.

### **Statistical Analyses**

The changes of pretreatment and post-treatment (PII,GI ,PPD,CAL,GR and CHPD) surements were the basis for data analysis with the use of a statistical software program. The paired-samples t-test was used to compare the mean values of presurgical and postsurgical treatment .

### **Results**

Subject recruitment started in January 2008 and was completed by the end of June 2011. The first surgical procedure was carried out in March 2008, and all the 6-month follow-up visits were completed in December 2011. Data entry of all information and statistical analysis were performed by the end of December 2011.

Eighteen patients completed the study with clinical data collected at baseline and after 6 months post-treatment. (8 males, 10 females).

Table 1. showed Patient characteristics at baseline. The mean age was 50±10.5 years, including a majority of females.

Table 1.Patient characteristics at baseline.

Age	50±10.5
Female(%)	10(55.6)

The means of full-mouth plaque and gingival index scores at surgical site are shown in Table 2. The plaque and gingival index scores were maintained at low throughout the study, without statistical difference from baseline to 6 months for both parameters at the surgical site.

Table 2.Changes in clinical parameters(PII. And GI.)

	mean±S.D
<b>Plaque index</b>	
Baseline	0.52±0.3
6 months	0.41±0.2
P-value	N.S
<b>Gingival index</b>	
Baseline	1.62±0.3
6 months	1.51±0.28
P-value	N.S

PPD,CAL,GR and CHPD means are shown in Table 3.A reduction in PPD was observed ( $p<0.05$ ). The periodontal pocket depth was reduced to ( $3.8\pm1.52$  mm) 6 months after surgery(Fig 3&4),where it was ( $5.47\pm1.39$ mm) at baseline(presurgical treatment).

The amount of CAL presurgical treatment was ( $3.1\pm1.4$ ),while after 6 months was( $2.35\pm 1.34$  mm). It was reduced but the analysis showed that there was no significant difference in change of means before and 6 months after surgery ( $P>0.05$ ).

There was an increase in GR after 6 months ( $0.31\pm0.54$  mm),GR was ( $0.21\pm0.51$ )before surgical treatment .The difference between pre & post-surgery Wasn't significant ( $P>0.05$ ).

CHPD pre surgical treatment (Baseline) was( $3.5\pm1.3$  and  $2.2\pm1.3$ ) mm after 6 months.

They showed more reduction after treatment ,but there's non-significant difference between them. Table 3.Mean values and Standered deviations(±SD) For Presurgical(baseline) and Postsurgical treatment(after 6 months) of Periodontal pocket depth(PPD),clinical attachment level(CAL).gingival recession(GR)and closed horizontal pocket depth (CHPD) and Significant Differences between them.

	PPD	CAL	GR	CHPD
Baseline	5.47±1.39	3.1±1.4	0.21±0.51	3.5±1.3
6 months	3.8±1.52	2.35± 1.34	0.3±0.5	2.2±1.3
P-value	S.	N.S.	N.S.	N.S.

## **Discussion**

The treatment of furcation-involved teeth, however, still represents a challenge for clinicians(45). This class of lesions presents a poor response to non-surgical treatment (46,47).

Thus, the present study aimed to evaluate the clinical response and defect closure in human periodontal Class II molar furcation defects by mucoperiosteal flap surgery with a bovine-derived xenograft. Evaluation the clinical outcome(Plaque index(PII),Gingival index(GI), Periodontal pocket depth(PPD),clinical attachment level(CAL),gingival recession(GR)and closed horizontal pocket depth (CHPD))and re-evaluate these clinical parameters 6 months after treatment without surgical re-entry for a better diagnosis of bone regeneration and defect repair, other methods are needed among which, re-entry is the best,since it investigates furcation directly.

In our society ,it's not easy to do surgical re-entry procedure ,from ethical point of view. The use of re-entry surgery for evaluation may be questioned, but such secondary revision surgery is often necessary in non-research bone graft therapy as the average response in osseous defects is defect fill. Often, this leaves a residual defect that requires further therapy (either additional grafting, debridement, or conservative osseous resection).(48).

Because a bovine-derived xenograft applications have demonstrated good clinical results in infraosseous and furcations(28,32,33,49-52) .The present study showed improvements in the clinical parameters evaluated.The furcations presented a PPD reduction ,this reduction was also similar to that obtained in previous studies. The results of measuring PPD were approximately the same as (28,31,32,51).They reported a reduction of packet depth after 6 months.

The amount of CAL showed a reduction after 6 months following treatment, which was similar to that reported by(31,32,51),they mentioned a significant improvement of CAL following treatment ,in addition (28,31,46)found a significant reduction of CAL following treatment with xenograft accompanied with absorbing Membrane(28,31-33,46,47) .

This study also showed no significant reduction of CHPD and GR compare to baseline(pre-surgical treatment)this results may be due to increase trauma to the soft tissue during surgical operation,the improvement of CHPD indicates the resistance of the treated defect to the probe penetration. these results also agreed with the results of (Ak.Khoshkhoo N. 2004)(51),while (Reddy 2006)(31) found the same results(reduction of CHPD and GR)but theres a significant differences when compared these results to baseline. Bovine derived xenograft can be a favorable graft material replacing bone because of its biocompatibility and osteoconductive properties, its chemical composition, and crystalline structure and porosity morphology similar to human bone.

## **Conclusion**

Bovine derived xenograft can be a clinically useful graft material in treatment of Class II Molar Furcation Involvement.

**REFERENCES**

- 1- Lindhe J.,Thorkild K.,Niklaus P. lang.Treatment of furcation-involved teeth.Text book of clinical periodontology and implant dentistry.Third edition,Munksgard,Copenhagen 1998;PP 682-709.
- 2-Bassam Michael Kinaia, Jacob Steiger, Anthony L. Neely, Maanas Shah, and Monish Bhola. Treatment of Class II Molar Furcation Involvement:Meta-Analyses of Reentry Results. J Periodontol 2011; 82:413-428
- 3-Seppa`la` B, Seppa`la` M, Ainamo J. A longitudinal study on insulin-dependent diabetes mellitus and periodontal disease. J Clin Periodontol 1993;20:161-165.
- 4-Emrich LJ, Shlossman M, Genco RJ. Periodontal disease in non-insulin-dependent diabetes mellitus. J Periodontol 1991;62:123-131.
- 5-Axelsson P, Paulander J, Lindhe J. Relationship between smoking and dental status in 35-, 50-, 65-, and 75-year-old individuals. J Clin Periodontol 1998;25:297-305.
- 6-Mullally BH, Linden GJ. Molar furcation involvement associated with cigarette smoking in periodontal referrals. J Clin Periodontol 1996;23:658-661.
- 7-Masters DH, Hoskins SW. Projections of cervical enamel molar furcations. J Periodontol 1964;35:49-53.
- 8-Everett FG, Jump EB, Holder TD, Williams GC. The intermediate bifurcational ridge: A study of the morphology of the bifurcation of the lower first molar. J Dent Res1958 ;37:162-169
- 9-Bower RC. Furcation morphology relative to periodontal treatment. Furcation entrance architecture. J Periodontol 1979;50:23-27.
- 10-Bower RC. Furcation morphology relative to periodontal treatment. Furcation root surface anatomy. J Periodontol 1979;50:366-374.
- 11-Hou GL, Tsai CC. Cervical enamel projection and intermediate bifurcational ridge correlated with molar furcation involvements. J Periodontol 1997;68:687-693.
- 12-Vertucci FJ, Williams RG. Furcation canals in the human mandibular first molar. Oral Surg Oral Med Oral Pathol 1974;38:308-314.
- 13-Lowman JV, Burke RS, Pelleu GB. Patent accessory canals: Incidence in molar furcation region. Oral Surg Oral Med Oral Pathol 1973;36:580-584.
- 14- Gutmann JL. Prevalence, location, and patency of accessory canals in the furcation region of permanent molars. J Periodontol 1978;49:21-26.
- 15-Moskow BS, Canut PM. Studies on root enamel (2).Enamel pearls. A review of their morphology, localization, nomenclature, occurrence, classification, histogenesis and incidence.J Clin Periodontol 1990;17:275-281.
- 16-Wang HL, Burgett FG, Shyr Y. The relationship between restoration and furcation involvement on molar teeth. J Periodontol 1993;64:302-305.
- 17-Mu`ller HP, Eger T. Furcation diagnosis. J Clin Periodontol 1999;26:485-498.
- 18-Goldman MJ, Ross IF, Goteiner D. Effect of periodontal therapy on patients main- tained for 15 years or longer. A retrospective study. J Periodontol 1986;57:347-353.
- 19-McFall WT Jr.Tooth loss in 100 treated patients with periodontal disease.A long-term study. J Periodontol 1982;53:539-549.
- 20-Hirschfeld L, Wasserman B. A long-term survey of tooth loss in 600 treated perio- Dontal patients. J Periodontol 1978;49:225-237.
- 21-Schallhorn RG, Hiatt WH, Boyce W. Iliac transplants in periodontal therapy. J Periodontol 1970;41:566-580.
- 22-SchallhornRG.Present status ofosseous grafting procedures JPeriodontol1977;48:570- 576.
- 23-Mellonig JT, Bowers GM, Bright RW, Lawrence JJ.Clinical evaluation of freeze- dried bone allografts in periodontal osseous defects. J Periodontol 1976;47:125-131.
- 24-Melcher AH. On the repair potential of periodontal tissues. J Periodontol 1976;47:256- 260.
- 25-Becker W, Becker BE, Berg L, Prichard J, Caffesse R,Rosenberg E. New attachment after treatment with root isolation procedures: Report for treated Class IIIand Class II furcations and vertical osseous defects.IntJ Periodontics RestorativeDent1988;8(3)8-23.

- 26-Bowers GM, Chadroff B, Carnevale R, et al. Histologic evaluation of new attachment apparatus formation in humans. Part I. *J Periodontol* 1989;60:664-674.
- 27-Nasr HF, Aichelmann-Reidy ME, Yukna RA. Bone and bone substitutes. *Periodontol* 2000 1999; 19: 74-86.
- 28-Houser BE, Mellonig JT, Brunsvold MA, Cochran DL, Meffert RM, Alder ME. Clinical evaluation of anorganic bovine bone xenograft with a bioabsorbable collagen barrier in the treatment of molar furcation defects. *Int J Periodontics Restorative Dent* 2001;21(2):161-169.
- 29-Johan C. Wohlfahrt, Anne M. Aass, Hans J. Rønold, Lars Heijl, Håvard J. Haugen, and Ståle P. Lyngstadaas. Microcomputed Tomographic and Histologic Analysis of Animal Experimental Degree II Furcation Defects Treated With Porous Titanium Granules or Deproteinized Bovine Bone. *J Periodontol* .2012,83:211-221.
- 30-Bowers GM, Schallhorn RG, McClain PK, Morrison GM, Morgan R, Reynolds MA. Factors influencing the outcome of regenerative therapy in mandibular Class II furcations: Part I. *J Periodontol* 2003;74:1255-1268.
- 31-Reddy KP, Nayak DG, Uppoor AS. A clinical evaluation of anorganic bovine bone graft plus 10% collagen with or without a barrier in the treatment of class II furcation defects. *J Contemp Dent Pract*. 2006 15;7(1):60-70.
- 32-Taheri M, Molla R, Radvar M, Sohrabi K, Najafi MH. An evaluation of bovine derived xenograft with and without a bioabsorbable collagen membrane in the treatment of mandibular Class II furcation defects. *Aust Dent J*. 2009; 54(3):220-7.
- 33-Simonpietri-C JJ, Novaes AB Jr., Batista EL Jr., Filho EJ. Guided tissue regeneration associated with bovine-derived anorganic bone in mandibular Class II furcation defects. 6-month results at re-entry. *J Periodontol* 2000;71:904-911.
- 34-Reynolds MA, Aichelmann-Reidy ME, Branch-Mays GL, Gunsolley JC. The efficacy of bone replacement grafts in the treatment of periodontal osseous defects. A systematic review. *Ann Periodontol* 2003;8:227- 265.
- 35-Murphy KG, Gunsolley JC. Guided tissue regeneration for the treatment of periodontal intrabony and furcation defects. A systematic review. *Ann Periodontol* 2003;8:266-302.
- 36-Mathew JJ, Arun Kumar MS, Bhat KM. Evaluation of a porous hydroxyapatite alloplast in the management of grade II furcation defects in molars; A case series. *J Clin Exp Dent*. 2012;4(2):e97-101.
- 37-Machtei EE. Outcome variables for the study of periodontal regeneration. *Ann Periodontol* 1997;2:229- 239.
- 38-Jepsen S, Eberhard J, Herrera D, Needleman I. A systematic review of guided tissue regeneration for periodontal furcation defects. What is the effect of guided tissue regeneration compared with surgical debridement in the treatment of furcation defects? *J Clin Periodontol* 2002;29:103-116.
- 39-Armitage, G. C. (1999) Development of a classification system for periodontal diseases and conditions. *Annals of Periodontology* 1999; 4 ,1-6.
- 40-Silness J. & Loe, H. Periodontal disease in pregnancy. II. Correlation between oral hygiene and periodontal condition. *Acta Odontologica Scandinavica*(1964) 122, 121-135.
- 41-Loe H., J. Silness: Periodontal disease in pregnancy. 1-Prevalence and severity. *Acta Odont .Scand* 1963;21:533-551.
- 42-Sanders JJ, Sepe WW, Bowers GM, Koch RW, Williams JE, Lekas JS, Mellonig JT, Pelleu GB Jr, Gambill V. Clinical evaluation of freeze-dried bone allografts in periodontal osseous defects. III. Composite freeze-dried bone allografts with and without autogenous bone grafts. *J Periodontol* 1983; 54: 1-11.
- 43-Carols Solís, Antonio Santos, José Nart, and Deborah Violant. 0.2% Chlorhexidine Mouthwash With an Antidiscoloration System Versus 0.2% Chlorhexidine Mouthwash: A Prospective Clinical Comparative Study. *J Periodontol* 2011, 82, 80-85.
- 44-Garrett S, Bogle G. Periodontal regeneration. A review of flap management. *Periodontol* 2000 1993: 1: 101-108.

- 45-Huynh-Ba G, Kuonen P, Hofer D, Schmid J, Lang NP, Salvi GE. The effect of Periodontal therapy on the survival rate and incidence of complications of multirooted teeth with furcation involvement after an observation period of at least 5 years:a systematic review.J Clin Periodontol 2009; 36: 164–176.
- 46-Del Peloso Ribeiro, E., Bittencourt, S., Nociti,F. H. Jr., Sallum, E. A., Sallum, A. W. & Casati, M. Z. Comparative study of ultrasonic instrumentation for the nonsurgical treatment of interproximal and noninterproximal furcation involvements. Journal of Periodontology 2007 ;78, 224–230.
- 47-Cristiano Tomasi and Jan L. Wennström .Locally Delivered Doxycycline as an Adjunct to Mechanical Debridement at Retreatment of Periodontal Pockets: Outcome At furcation Sites. J Periodontol 2011, 82, 2, 210-218.
- 48-Yukna RA. Clinical evaluation of HTR polymer bone replacement grafts in human mandibular Class II molar furcations. J Periodontol 1994;65:342-349.
- 49-Kothiwale SV, Anuroopa P, Gajiwala AL. A clinical and radiological evaluation of DFDBA with amniotic membrane versus bovine derived xenograft with amniotic membrane in human periodontal grade II furcation defects. Cell Tissue Bank. 2009;10(4):317-26.
- 50-E.Todd Scheyer ,Diega Velasquez-Plata ,Michael A. Brunsvold ,David J. Lasho ,James T. Mellonig: A Clinical Comparison of a Bovine-Derived Xenograft Used Alone and in Combination With Enamel Matrix Derivative for the Treatment of Periodontal Osseous Defects in Humans. J Periodontol 2002, 73,4,423-432
- 51-Ak.Khoshkhoo Nejad, SH. Mohseni Salehi Monfared, M. Rooeintan. Bio-oss in Treatment of Furcation Class II Deffects and Comparison with Coronally Positioned Flap. Journal of Dentistry, Tehran University of Medical Sciences 2004; 1,26-31.
- 52-Sachin S. Shivanaikar, Mohamed Faizuddin.Treatment of periodontal bony defect with bovine derived xenograft and in combination of platelet rich plasma-A case report Archives of Oral Sciences & Research AOSR 2012;2(2):98-102.





Fig 1.pre-op periodontal pocket depth




Fig 2.Pre-op radiograph



Fig 3.post-op periodontal pocket depth(after 6 months)



Fig 4.post-op(after 6 months) radiograph.