

Semi - α - $T_{\frac{1}{2}}$ - Space in Bitopological Spaces

Qays H. I. Al-Rubaye

Al-Muthanna University, College of Education, Department of Mathematics

E-mail : alrubaye84@yahoo.com

Abstract :

The objective of this research is to study some properties of ij - semi - α - $T_{\frac{1}{2}}$ - spaces in bitopological spaces and their relationship with ij - semi - α - T_0 - spaces and ij - semi - α - T_1 - spaces .

Key words : Bitopological space , ij - semi - α - open set , ij - semi - α - $T_{\frac{1}{2}}$ - space .

الخلاصة :

الهدف من هذا البحث هو دراسة بعض خصائص فضاءات ij - شبہ الگا $T_{\frac{1}{2}}$ فی الفضاءات ثنائية التبولوجی و علاقتها مع فضاءات ij - شبہ الگا T_0 و فضاءات ij - شبہ الگا T_1 .

1. Introduction :

In 1963 Kelly [2] defined : a set equipped with two topologies is called a bitopological space , denoted by (X,τ_1,τ_2) where (X,τ_1) and (X,τ_2) are two topological spaces . This notion was studied in different senses , one of these is the ij - semi - α - open sets ,that suggested by Qays , 2012 [4] . In 1997 , Kumar Sampath , S . , [3] introduced the concept of ij - α -open sets in bitopological spaces . In 1981 , Bose , S. , [1] introduced the notion of ij - semi - open sets in bitopological spaces . In this paper we first (in section 2) ij - semi - α - open sets , and then (in section 3) we study especial case of ij - semi - α - $T_{\frac{1}{2}}$ - space in bitopological spaces in the sense of ij - semi - α - open sets .

2. Preliminaries :

Throughout the paper , spaces always mean a bitopological spaces , the closure and the interior of any subset A of X with respect to τ_i , will be denoted by $\tau_i-cl(A)$, and $\tau_i-int(A)$ respectively, for $i = 1,2$.

Definition 2.1 : [4]

Let (X,τ_1,τ_2) be a bitopological space , $A \subseteq X$. Then A is said to be ij - semi - α - open set if there exists an ij - α - open set U in X , such that $U \subseteq A \subseteq j-cl(U)$. The family of all ij - semi - α - open sets of X is denoted by ij - $S_\alpha O(X)$, where $i \neq j; i, j = 1,2$.

The following proposition will give an equivalent definition of ij - semi - α - open sets .

Proposition 2.2 : [4]

Let (X,τ_1,τ_2) be a bitopological space , $A \subseteq X$. Then A is an ij - semi - α - open set if and only if $A \subseteq j-cl(i-int(j-cl(i-int(A))))$.

Remark 2.3 : [4]

The intersection of any two ij - semi - α - open sets is not necessary ij - semi - α - open set as in the following example .

Example 2.4 :

Let $X = \{a,b,c\}$, $\tau_1 = \{X, \emptyset, \{a\}, \{b\}, \{a,b\}\}$, and $\tau_2 = \{X, \emptyset, \{a\}\}$. The family of all 12 -semi - α - open sets of X is : 12 - $S_\alpha O(X) = \{X, \emptyset, \{a\}, \{b\}, \{a,b\}, \{a,c\}, \{b,c\}\}$. Hence $\{a,c\}$ and $\{b,c\}$ are two 12 - semi - α - open sets , but $\{a,c\} \cap \{b,c\} = \{c\}$ is not 12 - semi - α - open set .

Proposition 2.5 : [4]

The union of any family of ij - semi - α - open sets is ij - semi - α - open set .

Remark 2.6 : [4]

(i) Every τ_i - open set is ij - semi - α - open set ,but the converse need not be true .

(ii) If every τ_i - open set is τ_i - closed and every nowhere τ_i - dense set is τ_i - closed in any bitopological space , then every ij - semi - α - open set is an τ_i - open set .

Definition 2.7 :

A set A is said to be an ij - semi - α - neighborhood of a point x if there exists an ij - semi - α - open set U such that $x \in U \subseteq A$.

Definition 2.8 : [4]

The complement of ij - semi - α - open set is called ij - semi - α - closed set . Then family of all ij - semi - α - closed sets of X is denoted by ij - $S_\alpha C(X)$, where $i \neq j; i, j = 1, 2$.

Remark 2.9 : [4]

The intersection of any family of ij - semi - α - closed sets is ij - semi - α - closed set .

Definition 2.10 : [4]

Let (X, τ_1, τ_2) be a bitopological space and $A \subseteq X$,the intersection of all ij - semi - α - closed sets containing A is called ij - semi - α - closure of A , and is denoted by ij - $S_\alpha cl(A)$; i.e ij - $S_\alpha cl(A) = \bigcap \{B \subseteq X : B \text{ is } ij\text{-semi-}\alpha\text{-closed set}, A \subseteq B\}$.

Theorem 2.11 : [4]

Let (X, τ_1, τ_2) be a bitopological space , and let $A \subseteq X$, then :

(i) ij - $S_\alpha cl(A)$ is the smallest ij - semi - α - closed set containing A .

(ii) A is ij - semi - α - closed set if and only if ij - $S_\alpha cl(A) = A$.

Remark 2.12 : [4]

Let (X, τ_1, τ_2) be a bitopological space , and let A, B be any subsets of X , then :

(i) $A \subseteq ij$ - $S_\alpha cl(A)$.

(ii) If $A \subseteq B$, then ij - $S_\alpha cl(A) \subseteq ij$ - $S_\alpha cl(B)$.

(iii) ij - $S_\alpha cl(ij$ - $S_\alpha cl(A)) = ij$ - $S_\alpha cl(A)$.

Lemma 2.13 :

If ij - $S_\alpha cl(\{x\}) \cap A \neq \emptyset$, for each $x \in ij$ - $S_\alpha cl(A)$, then $(ij$ - $S_\alpha cl(A)) - A$ does not contain a non empty ij - semi - α - closed set .

Proof :

Suppose $(ij$ - $S_\alpha cl(A)) - A$ contains a non empty ij - semi - α - closed set , say B , then $x \in B$ implies ij - $S_\alpha cl(\{x\}) \subseteq B \subseteq (ij$ - $S_\alpha cl(A)) - A$, and ij - $S_\alpha cl(\{x\}) \cap A = \emptyset$, which contradicts the hypothesis . ■

Definition 2.14 :

A subset A of a bitopological space (X, τ_1, τ_2) is said to be ij - semi - α - generalized closed set (briefly ij - semi - α - g - closed set) if ij - $S_\alpha cl(A) \subseteq U$ whenever $A \subseteq U$ and U is ij - semi - α - open set in (X, τ_1, τ_2) .

Lemma 2.15 :

A subset A of a bitopological space (X, τ_1, τ_2) is ij - semi - α - g - closed set if and only if ij - $S_\alpha cl(\{x\}) \cap A \neq \emptyset$ for each $x \in ij$ - $S_\alpha cl(A)$.

Proof :

Suppose that A is ij - semi - α - g - closed set, and for some $x \in ij$ - $S_\alpha cl(A)$, ij - $S_\alpha cl(\{x\}) \cap A = \emptyset$, then $A \subseteq X - (ij$ - $S_\alpha cl(\{x\}))$, where $X - (ij$ - $S_\alpha cl(\{x\}))$ is a

ij - semi - α - open set , so by definition (2.14) ij - S_α - $cl(A)$ $\subseteq X - (ij$ - S_α - $cl(\{x\}))$, hence $x \in X - (ij$ - S_α - $cl(\{x\}))$ i.e $x \notin X - (ij$ - S_α - $cl(\{x\}))$ which is a contradiction .

Conversely assume that for each $x \in ij$ - S_α - $cl(A)$, ij - S_α - $cl(\{x\}) \cap A \neq \emptyset$; if there is a ij - semi - α - open set U such that $A \subseteq U$ but ij - S_α - $cl(A) \not\subseteq U$ then there exists $x \in ij$ - S_α - $cl(A)$ and $x \notin U$, so $x \in X - U$ which implies ij - S_α - $cl(\{x\}) \subseteq X - U$ (since $X - U$ is ij - semi - α - closed) i.e ij - S_α - $cl(\{x\}) \cap A = \emptyset$, a contradiction . Therefore A is ij - semi - α - g - closed set . ■

3. Semi - α - $T_{1/2}$ - space in Bitopological Spaces :

In this section the notion of ij - semi - α - $T_{1/2}$ - space is introduced in bitopological spaces and their relationships with ij - semi - α - T_0 - space and ij - semi - α - T_1 - space are studied .

Definition 3.1 :

Let (X, τ_1, τ_2) be a bitopological space ;two subsets A and B of X are ij - semi - α - separated if each is disjoint from the other's ij - semi - α - closure . (i.e $A \cap ij$ - S_α - $cl(B) = \emptyset$ and ij - S_α - $cl(A) \cap B = \emptyset$) . Two points x and y in X are ij - semi - α - distinguishable if they do not have exactly the same ij - semi - α - neighborhoods (i.e there exists a ij - semi - α - open set containing x but not containing y or containing y but not containing x) . Two points x and y are ij - semi - α - separated if the singletons $\{x\}$ and $\{y\}$ are ij - semi - α - separated .

Definition 3.2 :

A bitopological space (X, τ_1, τ_2) is called ij - semi - α - T_0 - space if any two distinct points are ij - semi - α - distinguishable .

Remark 3.3 :

Every i - T_0 - space is ij - semi - α - T_0 - space.

Proof :

Follows from remark (2.6) (i) . ■

Remark 3.4 :

The converse of remark (3.3) is not true ,see the following example .

Example 3.5 :

Let $X = \{a, b, c, d\}$, $\tau_1 = \{X, \emptyset, \{a\}, \{a, d\}, \{b, c\}, \{a, b, c\}\}$, $\tau_2 = \{X, \emptyset, \{a\}\}$.

Then 12 - $S_\alpha O(X) = \tau_1 \cup \{\{a, b\}, \{a, b, d\}, \{a, c\}, \{a, c, d\}\}$ and it is clear that (X, τ_1, τ_2) is 12 - semi - α - T_0 - space but (X, τ_1) is not 1 - T_0 - space since the points b and c are not distinguishable .

Theorem 3.6 :

A bitopological space (X, τ_1, τ_2) is ij - semi - α - T_0 - space iff for each distinct points x, y in X , ij - S_α - $cl(\{x\}) \neq ij$ - S_α - $cl(\{y\})$.

Proof :

Suppose that (X, τ_1, τ_2) is ij - semi - α - T_0 - space and let $x \neq y$ be two points of X such that ij - S_α - $cl(\{x\}) = ij$ - S_α - $cl(\{y\})$, therefore $x \in ij$ - S_α - $cl(\{y\})$ and $y \in ij$ - S_α - $cl(\{x\})$. If U is ij - semi - α - open set such that $x \in U$ and $y \notin U$, then $y \in X - U$ (ij - semi - α - closed set) so ij - S_α - $cl(\{y\}) \subseteq X - U$ which means ij - S_α - $cl(\{x\}) \subseteq X - U$ and so $x \in X - U$ i.e $x \notin U$ a contradiction . Similarly the assumption that $x \notin V$ and $y \in V$ (for some ij - semi - α - open set V) leads to a contradiction , that is (X, τ_1, τ_2) is not an ij - semi - α - T_0 - space .

On the other hand, suppose that for each $x, y \in X$ and $x \neq y$ we have ij - S_α - $cl(\{x\}) \neq ij$ - S_α - $cl(\{y\})$,

therefore either $x \notin ij - S_\alpha - cl(\{y\})$ and so $x \in X - (ij - S_\alpha - cl(\{y\}))$ but $y \notin X - (ij - S_\alpha - cl(\{y\}))$, or $y \notin ij - S_\alpha - cl(\{x\})$ and so $y \in X - (ij - S_\alpha - cl(\{x\}))$ but $x \notin X - (ij - S_\alpha - cl(\{x\}))$ where $X - (ij - S_\alpha - cl(\{x\}))$ and $X - (ij - S_\alpha - cl(\{y\}))$ are ij -semi- α -open sets in (X, τ_1, τ_2) i.e x and y are ij -semi- α -distinguishable, hence (X, τ_1, τ_2) is ij -semi- α - T_0 -space. ■

Theorem 3.7 :

Every subspace of ij -semi- α - T_0 -space is ij -semi- α - T_0 -space.

Proof :

Let $(Y, \tau_{1Y}, \tau_{2Y})$ be a subspace of a bitopological space (X, τ_1, τ_2) and let X is ij -semi- α - T_0 -space. To prove that Y is ij -semi- α - T_0 -space, let $y_1 \neq y_2 \in Y$. Since $Y \subseteq X$, then $y_1 \neq y_2 \in X$ and X is ij -semi- α - T_0 -space. There exists ij -semi- α -open set G in X , such that $y_1 \in G$ and $y_2 \notin G$. So $G \cap Y$ is ij -semi- α -open set in Y and $y_1 \in G \cap Y$, $y_2 \notin G \cap Y$.

Hence $(Y, \tau_{1Y}, \tau_{2Y})$ is ij -semi- α - T_0 -space. ■

Definition 3.8 :

A bitopological space (X, τ_1, τ_2) is said to be ij -semi- α - $T_{\frac{1}{2}}$ -space if every ij -semi- α -g-closed set in (X, τ_1, τ_2) is ij -semi- α -closed.

Theorem 3.9 :

A bitopological space (X, τ_1, τ_2) is ij -semi- α - $T_{\frac{1}{2}}$ -space if and only if, for each $x \in X$, $\{x\}$ is ij -semi- α -closed or ij -semi- α -open.

Proof :

Assume that (X, τ_1, τ_2) is ij -semi- α - $T_{\frac{1}{2}}$ -space and $\{x\}$ is neither ij -semi- α -closed nor ij -semi- α -open then $X - \{x\}$ is not ij -semi- α -closed so $ij - S_\alpha - cl(X - \{x\}) = X \subseteq X$ i.e $X - \{x\}$ is ij -semi- α -g-closed set, by definition of ij -semi- α - $T_{\frac{1}{2}}$ -space, $X - \{x\}$ must be ij -semi- α -closed, a contradiction with the assumption.

On the other hand, suppose that for each x in (X, τ_1, τ_2) , $\{x\}$ is ij -semi- α -closed or ij -semi- α -open. Let A be ij -semi- α -g-closed set in (X, τ_1, τ_2) , then by lemma (2.13) and lemma (2.15) $(ij - S_\alpha - cl(A)) - A$ does not contain a non empty ij -semi- α -closed set, so if $x \in (ij - S_\alpha - cl(A)) - A$ then $ij - S_\alpha - cl(\{x\}) \subset (ij - S_\alpha - cl(A)) - A$ i.e $ij - S_\alpha - cl(\{x\}) \neq \{x\}$ which means $\{x\}$ is not ij -semi- α -closed, so it must be ij -semi- α -open, but $\{x\} \cap A = \emptyset$ implies $x \notin ij - S_\alpha - cl(A)$,

a contradiction, hence $(ij - S_\alpha - cl(A)) - A = \emptyset$. Therefore A is ij -semi- α -closed, and so (X, τ_1, τ_2) is

ij -semi- α - $T_{\frac{1}{2}}$ -space. ■

Theorem 3.10 :

If (X, τ_1, τ_2) is ij -semi- α - $T_{\frac{1}{2}}$ -space then it is ij -semi- α - T_0 -space.

Proof :

Suppose (X, τ_1, τ_2) is ij -semi- α - $T_{\frac{1}{2}}$ -space by theorem (3.9) every singleton is either ij -semi- α -closed or ij -semi- α -open. Let $x \neq y$ (in X), if $\{x\}$ is ij -semi- α -closed then $X - \{x\}$ is ij -semi- α -open set containing y but not containing x ; and if $\{x\}$ ij -semi- α -open then it is containing x but not containing y . So (X, τ_1, τ_2) is ij -semi- α - T_0 -space. ■

Journal of Kerbala University , Vol. 11 No.1 Scientific . 2013

Remark 3.11 :

The converse of theorem (3.10) is not true , see the following example .

Example 3.12 :

Let $X = \{a, b, c\}$, $\tau_1 = \{X, \emptyset, \{a\}, \{a, b\}\}$, $\tau_2 = D$ (the discrete topology on X) then $12 - S_\alpha O(X) = \tau_1$ and (X, τ_1, τ_2) is a 12 -semi- α - T_0 -space but not 12 -semi- α - $T_{\frac{1}{2}}$ -space (since $\{b\}$ is neither 12 -semi- α -closed nor 12 -semi- α -open) .

Theorem 3.13 :

Every i - $T_{\frac{1}{2}}$ -space is ij -semi- α - $T_{\frac{1}{2}}$ -space .

Proof :

It follows the fact that $\tau_i \subseteq ij - S_\alpha O(X)$ and theorem (3.9) . ■

Remark 3.14 :

The converse of theorem (3.13) is not true , see the following example .

Example 3.15 :

Let $X = \{a, b, c\}$, $\tau_1 = \{X, \emptyset, \{a\}, \{a, b\}\}$, $\tau_2 = I$ (the indiscrete topology on X) then $12 - S_\alpha O(X) = \tau_1 \cup \{\{a, c\}\}$ and (X, τ_1) is not 1 - $T_{\frac{1}{2}}$ -space but (X, τ_1, τ_2) is 12 -semi- α - $T_{\frac{1}{2}}$ -space .

Definition 3.16 :

A bitopological space (X, τ_1, τ_2) is said to be ij -semi- α - T_1 -space if any two distinct points in X are ij -semi- α -separated .

Proposition 3.17 :

A bitopological space (X, τ_1, τ_2) is ij -semi- α - T_1 -space iff every singleton subset $\{x\}$ of X is ij -semi- α -closed .

Proof :

Suppose (X, τ_1, τ_2) is ij -semi- α - T_1 -space , and $x \in X$, if $y \in ij - S_\alpha - cl(\{x\})$ but $x \neq y$ then $ij - S_\alpha - cl(\{y\}) \subseteq ij - S_\alpha - cl(\{x\})$. On the other hand by definition (3.16) we have $\{y\} \cap ij - S_\alpha - cl(\{x\}) = \emptyset$ which is a contradiction , so $ij - S_\alpha - cl(\{x\}) = \{x\}$ i.e $\{x\}$ is ij -semi- α -closed set .

Conversely if for each x , $\{x\}$ is ij -semi- α -closed then $ij - S_\alpha - cl(\{x\}) = \{x\}$ and any two distinct points of X are ij -semi- α -separated i.e (X, τ_1, τ_2) is ij -semi- α - T_1 -space . ■

Remark 3.18 :

Every i - T_1 -space is ij -semi- α - T_1 -space.

Proof :

It follows from the fact that in i - T_1 -space every singleton is an i -closed set in (X, τ_1) also any i -closed set in (X, τ_1) is ij -semi- α -closed set in (X, τ_1, τ_2) . ■

Remark 3.19 :

The converse of remark (3.18) is not true , see the following example .

Example 3.20 :

Let $X = \{a, b, c, d\}$, $\tau_1 = \{X, \emptyset, \{a\}, \{d\}, \{a, d\}, \{b, c\}, \{a, b, c\}, \{b, c, d\}\}$, and $\tau_2 = \{X, \emptyset, \{a\}, \{d\}, \{a, d\}\}$ then $12 - S_\alpha O(X) = \tau_1 \cup \{\{a, b\}, \{a, c\}, \{b, d\}, \{c, d\}, \{a, c, d\}, \{a, b, d\}\}$. So (X, τ_1, τ_2) is 12 -semi- α - T_1 -space since all singletons are 12 -semi- α -closed sets but (X, τ_1) is not a 1 - T_1 -space since $\{b\}$ and $\{c\}$ are not 1 -closed sets .

Theorem 3.21 :

Every subspace of ij -semi- α - T_1 -space is ij -semi- α - T_1 -space .

Proof :

Let $(Y, \tau_{1Y}, \tau_{2Y})$ be a subspace of a bitopological space (X, τ_1, τ_2) and let (X, τ_1, τ_2) is ij -semi- α - T_1 -space. To prove that Y is ij -semi- α - T_1 -space, let $y_1 \neq y_2 \in Y$. Since $Y \subseteq X$, then $y_1 \neq y_2 \in X$ and since X is ij -semi- α - T_1 -space. Then there exist two ij -semi- α -open sets G, H in X , such that $y_1 \in G$, but $y_2 \notin G$; and $y_2 \in H$, but $y_1 \notin H$. Then we obtain two sets $G_1 = G \cap Y, H_1 = H \cap Y$ are ij -semi- α -open sets in Y , we have $y_1 \in G_1$, but $y_2 \notin G_1$; and $y_2 \in H_1$, but $y_1 \notin H_1$. Hence $(Y, \tau_{1Y}, \tau_{2Y})$ is ij -semi- α - T_1 -space. ■

Remark 3.22 :

Every ij -semi- α - T_1 -space is ij -semi- α - T_0 -space, but the converse is not true.

Proof :

This follows directly from the definitions (3.2) and (3.16). ■

But the converse is not true as the following example.

Example 3.23 :

Let $X = \{a, b, c\}$, $\tau_1 = \{X, \emptyset, \{a\}\}$, and $\tau_2 = \{X, \emptyset, \{a\}, \{a, b\}\}$, the family of all 12 -semi- α -open sets of X is : $12\text{-}S_\alpha O(X) = \{X, \emptyset, \{a\}, \{a, b\}, \{a, c\}\}$. If we take a and b , $a \neq b$, then we can not find two 12 -semi- α -open sets, such that one of them contains a but not b and the other contains b but not a . Therefore (X, τ_1, τ_2) is not 12 -semi- α - T_1 -space, but it is clear that (X, τ_1, τ_2) is ij -semi- α - T_0 -space.

Theorem 3.24 :

If (X, τ_1, τ_2) is ij -semi- α - T_1 -space then it is ij -semi- α - $T_{1/2}$ -space.

Proof :

It follows from theorem (3.9) and proposition (3.17). ■

Remark 3.25 :


The converse of theorem (3.24) is not true, see the following example.

Example 3.26 :

Let $X = \{a, b\}$, $\tau_1 = \{X, \emptyset, \{a\}\}$, $\tau_2 = D$ (the discrete topology on X), then $12\text{-}S_\alpha O(X) = \tau_1$, and (X, τ_1, τ_2) is 12 -semi- α - $T_{1/2}$ -space (since $\{a\}$ is 12 -semi- α -open and $\{b\}$ is 12 -semi- α -closed), but not 12 -semi- α - T_1 -space since $\{a\}$ is not 12 -semi- α -closed.

Remark 3.27 :

The following diagram shows the relations between $i - T_k$ and ij - semi - α - T_k spaces , where $k = 0, \frac{1}{2}, 1$:



References :

- [1] Bose , S. , “ Semi - open sets , semi - continuity and semi - open mappings in bitopological spaces ” , Bull . Cal . Math . Soc . , 73 (1981) , 237 – 246 .
- [2] Kelly, J .C ., “ Bitopological Spaces ”, Proc . London Math . Soc . , 13 (1963) , 17 – 89 .
- [3] Kumar Sampath , S ., “ On a Decomposition of Pairwise Continuity ”, Bull. Cal . Math . Soc . , 89 (1997) , 441 – 446 .
- [4] Qaye , H . I. Al-Rubaye , “ Semi - α - Separation Axioms in Bitopological Spaces ” , Al-Muthanna Journal of Pure Sciences , Vol .1 , No .1 , September , (2012) , 190 – 206 .