

## **Modifying Plaque assay and Clearance test as tools in determination of phage typing for Escherichia coli bacterial interspecies**

Mohammed Abdul Rahmman Mohammed

Department of biology -College of science - Baghdad University - Baghdad - Iraq

### **Abstract :**

Bacteriophage of *Escherichia coli* interspecies from sewage samples were isolated , The phage particles were isolated from two different sewage samples , First sample was collected from sewage sample of Baghdad university and second sample was isolated from domestic sewage sample , first sample showed phages specialized for three *E. Coli* interspecies bacteria (first plate ) and two *E. Coli* interspecies bacteria (second plate ) , meanwhile second sample showed phage specialized for two *E. Coli*. interspecies

Study of appearance of *Escherichia coli* phages from first sample showed three types of *E. coli* phages with different size of inhibition zone 2 , 1.5,1 mm respectively ( first plate ) , meanwhile *E. Coli* interspecies bacteria showed phages related with two interspecies with size of inhibition zone 1.5 ,1 mm respectively ( second plate ) , on other hand , The second sample showed also two interspecies *E. coli* with inhibition zone (1,2)mm . We designed experimental method which showed the modifying method of phage assay to determine phage typing assay .We tested phage particles with different bacterial strains ( *E. coli* , *shegilla* and *Serratia* ) from different sources and the control was the host of each bacteriophages by taking the O.D for all the tests and controls , to setup new criteria for phage typing .:and this test is called ( Clearance Test )

The result showed that O.D for Test 1 , 2 , 3, was ( 1.6 , 1.2 . 1.7 ) for ( *E. coli* , *shegilla* and *Serratia* ) bacterial strains , meanwhile the control tests was ( 0.3 , 0.2, 0.4 ) for strains isolated from first sample (first and second plate ) and second samples with different interspecies respectively .

This result can predict high specificity of phage strain and this method can be used to determine interspecies strains .

So from this experimental we can identify only Clearance Test by measuring only O.D. of bacterial strain with different phages a instead of going through plaque assay .

**Key words :** phage typing , *E. Coli* interspecies , Plaque assay and clearance test

### **الخلاصة**

فاجات بكتريا ال *Escherichia coli* تحت النوع من عينات مياه المجاري ، العينة الاولى جمعت من عينة مياه المجاري لجامعة بغداد والعينة الثانية اخذت من مياه مجاري منزلية ، العينة الاولى كانت تحتوي على ثلاث انواع فاجات خاصة لل *Escherichia coli* تحت انواع مختلفة ( الطبق الاول ) ونوعين لفاجات ال *Escherichia coli* تحت النوع ( الطبق الثاني ) ، بينما العينة الثانية كانت تحتوي كذلك فاجين خاص لل *Escherichia coli* دراسة مظهر فاجات بكتريا ال *Escherichia coli* من العينة الاولى كانت تحتوي على ثلاث انواع من الفاجات مع احجام مختلفة لحلقات تثبيط ( 1.5 , 1 , 2 ) ، مليمتر للطبق الاول ، بينما فاجات ال *Escherichia coli* كانت تحوي نوعين من حلقات التثبيط ( 2 , 1 ) مليمتر نحن صممنا تجربة والتي عرضت طريقة محورة لتحليل الفاج لتحديد (نوع البكتريا باستخدام الفاج ) phage typing من مصادر مختلفة بينما المحلول القياسي تمثل بمضيف المتمثل ببكتريا التي عزل منها الفاج ومن خلال قياس الكثافة الضوئية لكل من الفحوصات والمحلول القياسي يتم تحديد نوع البكتريا وعائدتها الى فاج معين او لا وهذا الفحص سمي ( Clearancetest ) .

النتائج عرضت الكثافة الضوئية لفحص 1,2,3 كانت ( 1.6 , 1.2 , 1.7 ) لبكتريا ال *Escherichia coli* و ال *Serratia* وال *Shigella* على التوالي بينما فحص المحلول القياسي ( 0.3 , 0.2 , 0.4 ) للعينة الاولى للتطبيقين الاول والثاني والعينة الثانية على التوالي . هذه النتائج اظهرت خصوصية عالية للعائيات وهذه الطريقة ممكن استخدامها لتحديد تحت النوع باستخدام فحص ( Clearance test ) فقط

## **Introduction**

Phage typing is a rapid, economical, reliable, and reproducible technique, requiring no specialized equipment, for fingerprinting disease-causing agents for epidemiological investigation and surveillance. Intraspecies differentiation of bacteria can be based on taxonomic features, such as morphology, biochemical properties (biotyping), virulence (pathotyping), and antigenic structure (serotyping). In addition, Phage typing is a wide variety of genome-based taxonomic techniques (1) which have been developed such as pulsed-field gel electrophoresis (PFGE) (3), amplified-fragment length polymorphism (AFLP) (4), and amplification of repetitive bacterial DNA elements (5). Other typing systems are based on sensitivity to specific chemicals, including antibiotics, plasmid profiling, ribotyping, and the production of or sensitivity to bacteriocins and bacteriophages (lysotyping or phage typing) (6).

Phage typing provides long-term and internationally comparable surveillance data. Because of their recent introduction, such information is not available for molecular techniques. Phage typing of enteric pathogens has been and still is being used successfully to characterize disease-causing agents for epidemiological investigation and surveillance (6, 7)

A functional phage typing system includes the following characteristics:

- a) A panel of genetically and phenotypically stable temperate or lytic phages possessing broader rather than narrow host-range specificities.
- b) Results, which are obtained quickly and clear-cut and require limited training in interpretation.
- c) Method that can be standardized.
- d) Bacterial cells must display a stable phage type over time.
- e) Bacterial cells from a phage lytic reactions can be easily determined. (8)

The typing phages may be isolated either from the environment (sewage, river, lake water)

Phage isolation from the bacterial culture is possible directly from a rapidly growing in broth culture, after UV irradiation or mitomycin-c treatment

Biological features of the phages are a very important issue Lytic as well as temperate bacteriophages may be used for typing (9, 10)

### ***Goals of experiment :***

- 1 . Realize very efficient method to determine phage typing from modifying plaque assay .
- 2 . Set up criteria to show positive and negative result of phage typing method by using Clearance Test .
- 3 . The typing method goes through interspecies with high accuracy result

## **Material and Method**

### **Modifying plaque assay**

- 1 . Pour about 40 ml of sterile molten phage agar into each plate then Sub-culture each strain of the species to be phage typed onto an Luria Broth agar plate to obtain isolated colonies.
2. Incubate the LB plate at 37 °C overnight.
3. Pick, with a sterile inoculating needle, a small amount from the center of a smooth colony and suspend each pick in one tube of LB broth.
4. Incubate these culture tubes in a water bath shaker at 37 °C and 100 rpm for approximately 1 h or until the culture becomes slightly turbid.
5. Flood a dried phage agar plate with the phage broth culture using a disposable transfer pipette to produce a bacterial "lawn" of the test strain.
6. Dispense one drop of each phage on to the phage agar plate
- 7 Remove phage agar plate and allow phage drops to dry on the phage agar plate for a maximum of 15 min with the lid ajar.
8. Repeat the process with a plate that has not been inoculated with bacteria. This is the sterility check on the phage preparations. Invert and incubate the dried plates at 37 °C overnight. (8,10)

### **Clearance Test**

- 1 . Take 3 ml of phage buffer in two tubes for each strain to be tested , first tube as test and second tube as control .
- 2 . Suspend 0.1 of each bacterial strains in control and test tubes .
- 3 . Add to test tubes phage particles of host bacterial strains meanwhile the addition normal saline in control tubes .
- 4 . All tubes are incubated for 24 hours at 37 C to show lytic activity and reduction of growth in test tubes for each bacteriophage with it's bacterial strains hosts till get clearance (11 ,13)

### **ResultandDiscussion**

The modifying plaque assay technique showed incredible result of interspecies of *E. coli* bacterial strain from different samples

The appearance of *Escherichia coli* interspecies showed that there were three different isolated phages strains with zone inhibition ( 2 , 1.5 , 1 ) mm ( first plate ) meanwhile there were only two different isolated phage strain with zone of inhibition ( 2 , 1 ) mm respectively (second plate ) as shown in figure .(1,1.5)



**Figure (1) : show plaqued assay of bacteriophage of *Escherichia coli* interspecies for sample one ( first plate )**



**Figure (2) : show plaque assay of bacteriophage of *Escherichia coli* interspecies for sample one ( second plate )**

On other hand the *Escherichia coli* phage particle of second sample show also two interspecies with appearance ( 1,2) mm as in figure ( 3 ) .



**Figure (3) : show plaque assay of bacteriophage of *Escherichia coli* interspecies for sample two**

The modifying experimental plaque assay set up as test and control .

Test 1., 2, 3, with different bacterial strains *Escherichia colishegilla* and *Serratia* respectively meanwhile the control 1,2,3 *Escherichia coli* from sample 1 (first and second plates ) and sample 2 . We set up criteria to establish new method to evaluate the modifying plaque assay by clearance test through determine the O.D. of bacterial control and tests .

The results showed that positive sample with plaque assay showed Optical .Density. not more than 0.4 in which control 1,2,3 showed 0.3 and 0.32 and 0.38 respectively mean while the negative samples showed minimum O.D was 1 in which test 1,2,3,showed 1.6, 1.7 ,1.2 respectively

The result dramatically significantly to determine interspecies bacteria doing clearance test instead of using more sophisticate technique like plaque assay

The reduction of O. D for positive result ( control ) to (0.4) as maximum meanwhile the negative result was (1) as a minimum in due to the bacteriophage in control was specific to bacterial strain in sample therefore bacteriophage lyses bacterial strain and led to reduction in number of bacterial strain then reduction in O.D. of those strain , on other meaning the lytic activity of bacteriophage to bacterial strains in control led to reduction in turbidity of growth and clearance the sample therefore we call it clearance test

The tests samples showed O.D not less than (1) in due to the bacteriophage in tests was not specific to bacterial strains in sample therefore bacteriophage does not lyse bacterial strain and does not lead to reduction in number of bacterial strain .therefore the tests tubes were turbid with high value of O.D.

Our findings are in agreement with those Muyad and Khakhria in that clearance test gave good indicator for lytic process of phage particle by which the O.D. goes down to become less than 0.4 Therefore clearance test can be used as a tool for phage typing assay , by which we conclude that the modifying plaque assay is a incredible technique to isolate different phages from different samples and use those bacteriophages to identify not only the difference between bacterial strains but also the difference between interspecies of different strains in very specific manner by Clearance test .

## **References**

1. G'urtler V. and B.C. Mayall. 2001. Genomic approaches to typing, taxonomy and evolution of bacterial isolates. *International Journal of Systematic and Evolutionary Microbiology* 51:3–16.
2. Isaacs, S.; J. Aramini, B. ; Ciebin, J.A ; Farrar, R. ;Ahmed, D.; Middleton, A.U. and Chandran, L.J., *et al.* 2005. An international outbreak of salmonellosis associated with raw almonds contaminated with a rare phage type of *Salmonella enteritidis*. *Journal of Food Protection* 68:191–198.
3. Woodward, D.L.; C.G. Clark; R.A. Caldeira ; R. Ahmed; G. Soule; L. Bryden; H. Tabor; P. Melito, *et al.* 2005. Identification and characterization of *Shigella boydii* 20 serovar nov., new and emerging *Shigella* serotype. *Journal of Medical Microbiology* 54:741–748.
4. McLaughlin, J.; L.J. Castrodale; M.J. Gardner; R. Ahmed, and B.D. Gessner. 2006. Outbreak of multidrug resistant *Salmonella Typhimurium* associated with ground pork served at a school potluck. *Journal of Food Protection* 69:666–670.
5. Demczuk, W.; G. Soule; C. Clark; H.W. Ackermann; R. Easy; R. Khakhria; F. Rodgers, and R. Ahmed. 2003. Phage-based typing scheme for *Salmonella enterica* serovar Heidelberg, a causative agent of food poisonings in Canada. *Journal of Clinical Microbiology* 41: 4279–4284
6. Ikeda, Y.; H. Saito; K. Miura; J. Takagi; H. Aoki; Y. Ikeda; H. Saito; K.i. Miura *et al.* 2004. DNA base composition, susceptibility to bacteriophages and interspecific transformation as criteria for classification in the genus *Bacillus*. *Journal of General & Applied Microbiology* 50:353–362.

## **The First Scientific Conference the Collage of Education for Pure Sciences 2012**

7. Rabsch, W.; R.A. Helm, and A. Eisenstark. 2004. Diversity of phage types among archived cultures of the Demerec collection of *Salmonella enterica* serovar Typhimurium strains. *Applied & Environmental Microbiology* 70:664–669.
8. Gevers, D.; G. Huys, and J. Swings. 2007. Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiology Letters* 205:31–36.
9. Sambrook, J. and D.W. Russell. 2001. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Press, Cold Spring Harbor, New York.
10. Sambrook, J. & Russell, D. W. (2001) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Press, Cold Spring Harbor, New York).
11. Carlson, K. 2005. Working with bacteriophages: Common techniques and methodological approaches., In E. Kutter and A. Sulakvelidze (Eds.), *Bacteriophages: Biology and Applications*. CRC Press, Boca Raton, FL.
12. Ackermann, H.-W. 2005. Electron microscopy., In E. Kutter and A. Sulakvelidze (Eds.), *Bacteriophages: Biology and Applications*. CRC Press, Boca Raton, FL.
3. Adams, M.D. 1959. *Bacteriophages*. Interscience Publishers, Inc., New York.
13. Carlson, K. 2005. Working with bacteriophages: Common techniques and methodological approaches., In E. Kutter and A. Sulakvelidze (Eds.), *Bacteriophages: Biology and Applications*. CRC Press, Boca Raton, FL.
- 14 – Muyad M. M. (2009) . Molecular identification of *Pseudomonas auroginosa* and its phage effect on those bacterial strain .
- 15 - Khakhria R, Duck D, Lior H (1990) . Extended phage-typing scheme for *Escherichia coli* O157:H7.. Dec;105(3):511-20